

Short communication

**Unveiling the seed storage behavior for conservation of Indian olibanum:
Boswellia serrata Roxb. Ex Coleb.**

UNDER PEER REVIEW

Abstract

The study attempted to understand the storage behavior of *Boswellia serrata* Roxb. seeds; for which seeds were collected from three locations of Shivalik region of Himalayas. Across the locations, the freshly collected seeds had average moisture content (MC) of 10.06% and exhibited an average germination of 73%. When the seeds were further desiccated to MC 7%; the average germination was reduced to 51.33%. After this, desiccated seeds were stored hermetically at +5°C temperature for the period of 24 months, and periodic germination trials were conducted at an interval of 4 months. The average germination percentage was reduced from 73% to 13.67% during the storage period; seed deterioration process and low temperature-induced chilling injury could be attributed for reduced germination rate. Our results also revealed that the Pinjore seed source outperforms the other two sources in germination rate with adjusted margin mean of 38.13. The insights derived from the study can aid in designing appropriate protocols for the conservation of the studied species.

Keywords: Germination, Seed Moisture Content, Storage, Conservation, *Boswellia serrata* Roxb. Ex Coleb.

1. Introduction

The vagaries of climate change spread across the living biome of this planet, and kingdom Plantae is no exception to that. The studies reported that climate change will threaten the existence of many members of the biodiversity. In order, to protect biodiversity from the reported ill effects of climate change, the ex-situ conservation is often put forth. However, for proper implementation of the conservation programs, knowledge of seed storage behaviour is essential (Wyse and Dickie 2017). Based on seed storage behaviour, three types of seeds are recognized and often used: orthodox, recalcitrant, and intermediate or orthodox with restricted desiccation ability (Schmidt 2000). Baseline data for seed storage behavior is lacking for most of the forest tree species. *Boswellia serrata* is one such economically important tree species.

Boswellia serrata Roxb. Ex Coleb. (Burseraceae) is a moderate to large-sized deciduous tree endemic to India, Africa, and the Arabian Peninsula. It is the source of a most important gum-resin called 'salai guggul' or 'Indian olibanum' obtained from the bark after injury and is used for the treatment of rheumatoid arthritis, osteoarthritis, Crohn's disease, and asthma (Brendler et al. 2018; Mishra et al. 2020). Conventionally, the trees are propagated by seeds. However, poor fruit setting under open pollination (2.6 to 10%), inadequate production of viable seeds (Joshi et al. 1981), and scanty seed germination (10–20%) restrict the distribution and limits the presence of this species in its natural habitat (Ghorpade et al. 2010). Due to the difficulties in seed germination, regeneration, and destruction, the species' natural abundance has decreased.

Comment [M1]: Add the date of this study

Comment [M2]: Precise the site of study

Comment [M3]: and

Comment [M4]: write the reference correctly

Comment [M5]:

Comment [M6]: italic et al.,

Comment [M7]: write the reference correctly

Comment [M8]: write the reference correctly

Comment [M9]: write the reference correctly



Figure 1. Germinated seeds of *Boswelliaserrata*Roxb. Ex Coleb.



Figure 2. Dewinged and Winged seeds of *Boswelliaserrata*Roxb. Ex Coleb.

Comment [M10]: avoid putting photos in the introduction

For conservation biologists, foresters, and small industries, tree species preservation and sustainable exploitation have emerged as major issues. A major constraint for developing effective conservation strategies is the need for more information on their seed storage behavior. To the best of our knowledge, very little information is available on this species' seed storage behavior and germination. Thus, we investigated *Boswelliaserrata* seed germination (Figure 1) for different storage periods for two years to strengthen the strategies for conserving this species.

Comment [M11]: demonstrated the objective of this study

2. Materials and methods

The fruit is 1.3 cm long, swollen, and trigonous with three valves and 3-heart-shaped, 1-seeded pyrenes which are winged along the margins (Figure 2). Fruits were manually harvested during June 2021 from three sources of Shivalik Himalayas (i.e. Morni hills: N 30°40'50.5" and E 77°04'49.1"; Kalesar National Park: N 30°23'50.91" and E 77°30'23.41"; and Pinjore: N 30°44'51.0" and E 76°58'58.9") in Haryana, India. A total of 1000 fruits were collected from 8–10 trees (at least 100 m apart) at each location. Fruits were transported in gunny bags to the Forest Tree Seed Laboratory, FRI (Forest Research Institute), Dehradun (N 30°28'11.72" and E 78°03'36.99"). Seeds were extracted from fruits and cleaned. Seed samples without apparent physical damage or insect infestation were selected for the experiment.

Comment [M12]: where is figure 2?

2.1 Morphological analysis

For biometric determination 10 seeds were selected randomly in 4 replications for individual measurement of length and width using a digital caliper with a precision of 0.001 mm. The length was measured from the base to the apex, excluding the stalk and the width was measured in the midline of the fruits and seeds (Cheib 2009). After measurement of length and width, 100 seeds were weighed in 4 replications on an analytical balance accurate to 0.001 g. The quantitative characteristic data were submitted to descriptive analysis, calculated with Excel application, arithmetic mean and standard deviation of seeds.

Comment [M13]: correct

2.2 Germination test

To study the role of Seed Moisture content (SMC) in germination, we germinated seeds at initial moisture content, which varies from 9.25% to 11.60%. For medium to long-term storage of seeds, moisture content should be as low as possible (i.e., below 8-10%). Further periodic germination trials were carried out for two years in laboratory conditions, according to the International Seed Testing Association procedures (ISTA 2010). Seeds were considered as germinated upon emergence of radical (1 cm in length).

2.3 Desiccation study

About 2000 seeds (each seed source) were desiccated to a seed moisture content of 7% from initial moisture content of 9.25% (Pinjore), 9.34% (Kaleshar National Park) and 11.60% (Morni Hills) using silica gel drying method. The desired moisture content of each source example was monitored based on the below given equation (Hong and Ellis 1996).

Comment [M14]: correct

$$\text{Weight of seed lot (g) at MCd\%} = \frac{100 - \text{MCi\%}}{100 - \text{MCd\%}} \times \text{ISW (g)}$$

With MCd being the desired moisture content, MCi the initial moisture content, and ISW the initial seed lot weight. After this, whole seeds' Moisture content (MC) was verified following the low constant oven drying method (ISTA 2010). We randomly selected 1 gm seeds each in 4 replications and dried them at 103°C for 17h in a hot air oven. Seed MC was calculated on a fresh mass basis.

Comment [M15]: correct

2.4 Storage trials

Storage trials were conducted with desiccated seeds in an airtight plastic jar kept at +5°C in a walk in seed storage chamber. The viability of the stored seeds was monitored by germination tests, according to the International Seed Testing Association procedures (ISTA 2010), at intervals of 4 months for two years.

Comment [M16]: correct

2.5 Statistical analysis

To test the differences between the treatment and the collection source, we employed two factorial ANOVA (Afifi and Azen 1979). After running the ANOVA, by witnessing the statistical significance, we calculated adjusted marginal means for both treatment and source of seed collection to interpret meaningfully and robustly. The above-mentioned analysis was performed using STATA 17.0.

Comment [M17]: correct

3. Results and discussion

Table 1. Morphological parameters of *Boswellia serrata* seeds across different locations

Particulars	Seed Length (mm)	Seed Width (mm)	100 Seed weight (gm)
-------------	------------------	-----------------	----------------------

Morni Hills	8.6±0.15	7.06±0.06	4.24±0.02
Kalesar National Park	6.63±0.13	5.8±0.04	3.36±0.03
Pinjore	8.42±0.11	6.35±0.10	4.15±0.03
Average	7.88±0.13	6.4±0.07	3.92±0.03

The fruit is swollen and trigonous with three valves, and 3-heartshaped, 1-seeded pyrenes which are winged along the margins. From Table 1 we can observe that seed length, seed width and 100 seed weight was found to be highest for Morni Hills. However all these values were found to be lowest for Kalesar National Park. We determined that for Kalesar National Park, seed length, seed breadth, and 100 seed weight were 15.86%, 9.37%, and 14.28% less than average values. These seed dimensions can be considered as important traits for early selection of seed sources. Significant positive correlation between seed size and seedling growth has earlier been reported in Sweet Chestnut (*Castanea sativa Mill.*) (Tumpa et al. 2021).

Comment [M18]: correct




Figure 3. Effect of storage period on Germination and Mean Germination Time (MGT) of *Boswellia serrata* seeds collected from Morni hills, Haryana.

During hermetic storage at +5°C temperature, seed moisture content varied little. Whatever the period of storage, the extreme values found under experimental conditions remained close to the moisture content measured after desiccation, i.e. 7%. When *Boswellia serrata* seeds from Morni hills were stored hermetically at +5°C temperature, the rate of germination was decreased from 73% to 24% and Mean Germination Time increased from 4.48 to 5.20 days over the storage period of 24 months (Figure 3).




Figure 4. Effect of storage period on Germination and Mean Germination Time (MGT) of *Boswellia serrata* seeds collected from Kalesar NP, Haryana.

Whereas *B. serrata* seeds from Kalesar National Park did not survive for more than 12 months at +5 °C temperature. The rate of germination was decreased from 70% to 1% and Mean Germination Time also decreased from 6.41 to 4 days during storage period of 12 months. After 12 months the germination was found to be zero percent (Figure 4). However, *B. serrata* seeds from Pinjore survived for more than 24 months at +5 °C temperature. During this duration rate of germination decreased from 76% to 17% and Mean Germination Time increased from 3.7 to 6.11 days (Figure 5).




Figure 5. Effect of storage period on Germination and Mean Germination Time (MGT) of *Boswellia serrata* seeds collected from Pinjore, Haryana.

This decrease in germination trend could be attributed to seed deterioration process, which may be accelerated due to the disorganization of cell membranes, affecting the seed germination and vigor, mainly by increasing the leaching of essential constituents for germination, and loss of cell compartmentalization and repair mechanisms (Zhang et al. 2021). The current study discovered that there were significant variations in germination for each source as the storage period increased, suggesting that the germination of these seeds is negatively impacted by both seed age and storage duration. These findings also suggest that this species does not have seed dormancy; seeds can germinate as soon as the seed embryos imbibe water. Tropical seeds tend to have a shorter storage duration, because the higher temperatures of the tropics generates faster chemical reactions, such as respiration and photosynthesis, than in cooler climates. These trials demonstrate that *B. serrata* seeds drastically reduces germination with increase in storage period.

Comment [M19]: correct

Comment [M20]: Avoid abbreviation

The overall model was found to be statistically significant, which makes us interpret the model. The two-way factorial ANOVA results indicate that there exists a significant difference in germination among the storage period as well as the seed source. To interpret further, the adjusted margin means of germination rate were calculated for both treatments i.e., Storage period and Seed source. We discovered that with an adjusted margin mean of 38.13, the Pinjore seed source is found to be best in terms of germination percentage over the storage period, followed by Morni Hills and Kalesar National Park with an adjusted margin mean of 30 and 20.38 respectively. Interpretation based on marginal means is robust in comparison to actual means. Significant variation was found among the seed source for germination in different storage periods. This finding is in agreement with (Garomaet al. 2017) demonstrated that seed germination decreased with increased in seed storage period. It is also reported that genotype strongly influences the seed's vigor (Schmidt 2000).

Comment [M21]: correct

4. Conclusions

The present study revealed that *Boswellia serrata* seeds can tolerate desiccation below 10% moisture content, but with the advancement in storage period from 0 months to 24 months the germination percentage decreased drastically and MGT increased. The possible reason for decrease in germination with increase in storage period could be attributed to seed deterioration process and low temperature induced chilling injury. However further research should be carried out on storage behavior at temperature above +5°C to ascertain the role of temperature for decrease in germination. Our study also revealed that Pinjore seed source is superior among all other seed sources. Therefore, it is advisable that this seed source should be used for collection of bulk quantity of seeds to achieve better productivity.

Statements and Declarations

Data Availability

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

UNDER PEER REVIEW

Reference

- Afifi AA, Azen SP. Statistical Analysis: A computer-oriented approach. 2nded. New York: Academic Press. 1979; ISBN:9781483260716, 1483260712.
- Brendler T, Brinckmann JA, Schippmann U. Sustainable supply, a foundation for natural product development: The case of Indian frankincense (*Boswelliaserrata* Roxb. Ex Colebr.) J. Ethnopharmacol. 2018; 225:279–286. doi: 10.1016/j.jep.2018.07.017
- Cheib AL. Ecologia da germinação e potencial para formação de banco de sementes de espécies de *Arthrocareus* A. Berger (Cactaceae) endêmicas dos campos rupestres de Minas Gerais, Brasil. Programa de Pós-Graduação em Biologia Vegetal. Instituto de Ciências Biológicas. Universidade Federal de Minas Gerais. Belo Horizonte. 2009; 32.
- Garoma B, Chibsa T, Keno T, Denbi Y. Effect of storage period on seed germination of different maize parental lines. *Journal of Natural Sciences Research*. 2017; 7(4), 8-14.
- Ghorpade RP, Chopra A, Nikam TD. In vitro zygotic embryo germination and propagation of an endangered *Boswelliaserrata* Roxb., a source of boswellic acid. *Physiology and Molecular Biology of Plants*. 2010; 16(2):159-165. <https://doi.org/10.1007/s12298-010-0017-7>.
- Hong T, Ellis RH. A protocol to determine seed storage behavior. IPGRI Technical Bulletin No.1. International Plant Genetic Resources Institute, Rome, Italy. 1996. https://cropgenebank.sgrp.cgiar.org/images/file/learning_space/technicalbulletin1.pdf
- ISTA. International Rules for Seed Testing. The International Seed Testing Association, Zurichstr.50, Switzerland. 2010; ISBN-13978-3-906549-60-6.
- Joshi HB, Rashid MA, Venkataramany P. The Silviculture of Indian Trees. Controller of Publication, New Delhi, India. 1981.
- Mishram S, Bishnoi RS, Maurya R, Jain D. *Boswelliaserrata* Roxb.-A bioactive herbs with various pharmacological activities. *Asian J. Pharm. Clin. Res.* 2020; 13:33–39. <http://dx.doi.org/10.22159/ajpcr.2020.v13i11.39354>
- Schmidt LH. Guide to handling of tropical and subtropical Forest seeds. Danida forest Seed Centre. 2000.
- Tumpa K, Vidaković A, Drvodelić D, Šango M, Idžojić M, Perković I, Poljak I. The Effect of Seed Size on Germination and Seedling Growth in Sweet Chestnut (*Castanea sativa* Mill.). 2021. *Forests*. 12:858. <https://doi.org/10.3390/f12070858>
- Wyse SV, Dickie JB. Predicting the global incidence of seed desiccation sensitivity. *J Ecol*. 2017; 105(4):1082-1093. <http://dx.doi.org/10.1111/1365-2745.12725>
- Zhang K, Yin Z, Jing S, Jiasong M, Jun T. Deterioration of orthodox seeds during ageing: Influencing factors, physiological alterations and the role of reactive oxygen species.

Plant Physiology and Biochemistry, 2021; 158:475-485.
doi:10.1016/j.plaphy.2020.11.031.

UNDER PEER REVIEW

