

Prevalence of Gastrointestinal Parasite in Four Goat Breeds Slaughtered in Two Major Abattoirs in Akure, Ondo State, Nigeria

Abstract

Prevalence study of gastrointestinal parasites of different goat breeds was carried out on 240 faecal samples of goats comprising of 69 West African Dwarf, 55 Kano brown, Borno red 52 and 64 Red Sokoto breeds from two abattoirs in Akure, Ondo state, Nigeria. They were examined for intestinal helminth eggs using direct microscopic examination and sodium chloride floatation technique. Out of the 240 fecal samples examined, 56 (23.33%) were positive for gastrointestinal parasites. Sokoto red breed had a higher prevalence of 22 (34.38%) while West African dwarf breed had the lower prevalence of 12 (17.38%). Female goat had a higher prevalence of 35 (30.43%) while male had a prevalence of 21 (17.07%). The gastrointestinal parasites observed were *Strongyloidesstercoralis*, *Moniezia* spp., *Fasciola* spp., *Ascarislumbricoides*, *Ancylostomaduodenale*, *Diphyllobothriumlatum*, *Haemonchuscontortus* and *Trichostrongylus* spp. *Strongyloidesstercoralis* had the highest prevalence while *Moniezaspp*, *Haemonchuscontortus* and *Trichostrongylus* spp. had the lowest prevalence. Of the total 64 (26.67%) Sokoto red breeds, 22 (34.38%) had the highest gastrointestinal infection while of the total 69 (28.75%) West African dwarf goats, 12 (17.38%) had the lowest gastrointestinal infection prevalence. There is need for good and effective animal health management in the studied Abattoirs. Thus, the incidence of these parasites revealed in this study can be decreased by the use of effective management techniques, quick diagnosis and anthelmintic therapy, and education of animal owners on the importance of animal health.

Keywords: Gastrointestinal parasites, Abattoirs, Helminth eggs, and Anthelmintic drugs

Introduction

Goats are crucial species of ruminant livestock around the world, mostly in tropical and subtropical areas. It plays a crucial role in small-scale farming and the agricultural economy of developing

Comment [T1]: Add Study design, sampling technique (abattoir and goat), study period and conclusion to the abstract to make it to stand alone

Comment [T2]: How it was calculated? It should be scientific

Comment [T3]: This statement should be rephrase and merged with "Out of the 240 fecal samples examined, 56 (23.33%) were positive for gastrointestinal parasites. Sokoto red breed had a higher prevalence of 22 (34.38%) while West African dwarf breed had the lower prevalence of 12 (17.38%)." statement

Comment [T4]: It is shallow and not focused on the background of title

cultures by creating jobs and boosting household income. According to Nwosu et al. (2007), owning a goat may in some cases be a rural household's only source of realizable wealth due to the socioeconomic significance linked to goat ownership. Goats and other farm animals are valuable because there is a high demand for nutrient-dense animal protein (Anaeto et al. 2009). Due to their lower purchase price, improved fecundity, profusion, availability, and ease of domestication, goats, as small ruminants, have few benefits over large animals.

Comment [T5]: Add reference to support it

The management and rearing of goats are impacted by a variety of gastrointestinal parasites. Small ruminant production is still severely hampered by the high mortality rates brought on by parasitic diseases of the gastrointestinal tract (Perry et al. 2002). The inappropriate handling and unclean conditions in the tropics allow parasitism to flourish in animals. Nematodes (such *Ascaris lumbricoides* and *Trichuristrichiuria*), trematodes (like *Fasciola hepatica*), and cestodes (like tapeworms) are few gastrointestinal parasites that afflict ruminants. According to Osakwe and Anyigor (2007), goat intestinal parasites have a negative economic impact on underdeveloped farming communities. Ineffective digestion of the meal results in young animals growing more slowly, lighter animals at weaning, less milk being produced, increased susceptibility to other diseases, and subpar meat production (Barger, 1997). These parasites are a problem for both small- and large-scale farmers worldwide, but there is a greater incidence in sub-Saharan Africa in general and Nigeria in particular due to the availability of a wide range of agro-ecological factors suitable for diversified host and parasite species (Onaga et al. 2009).

The use of anthelmintics and antiprotozoal medicines to control intestinal parasite infections in ruminants has been successful over time (Gunathilaka et al. 2018). None of these, however, has been able to reduce the spread of parasitic illnesses. However, excessive anthelmintic use has resulted in anthelmintic residues in farm animal milk and meat, as well as anthelmintic-resistant parasites (Mohammed et al. 2019; Fissiha and Kinde 2021). In-depth knowledge of the pathophysiology and

epidemiology of these parasites, the context of immunity, and control over the host are essential for an anthelmintic and antiprotozoal approach to be effective. In order to prevent and treat illnesses in their earliest stages of infection in farm management methods, it is necessary to periodically study the prevalence of parasite species among farm animals. Thus, this research aims to evaluate the prevalence of gastrointestinal parasites and determine the influence of sex, breed and location on the prevalence of gastrointestinal helminths of goats in the study area.

Materials and Methods

Study area

Two major abattoirs in Akure, Ondo State, Nigeria were considered for the study. Animals were usually slaughtered early in the morning between the hours of 6.30 am and 8.00 am daily for a period of three months. Faecal samples were randomly collected from the intestines of slaughtered goats of both sexes (exotic and indigenous).

Study population

A total number of 240 goats were examined for the prevalence of gastrointestinal parasites. The study population consists of 125 male goats, 115 female goats brought to the abattoir for slaughter. The breeds of goats were Sokoto red, Kano brown, Borno red, West African dwarf.

Examination of collected faecal samples

The faecal samples of the goats were collected very early in the morning. Hand gloves were used to collect the samples from the rectum collected in clean labeled sterile universal vials, preserved in 10% formalin, and taken to the laboratory for microscopic examination using $\times 10$ and $\times 40$ magnification. Three diagnostic techniques were used for the faecal samples' examination, this includes formal ether concentration technique, sedimentation technique, and floatation techniques described by WHO (1994) with slightly modification.

Comment [T6]: What is your study design, sampling technique, your study period and sample size determination method for your study???

Comment [T7]: The study area were not described well.
1. Please include some information about situations of abattoirs.
2. Add origin of animal slaughtered at those abattoirs.
3. Longitude, latitudes, altitudes, mean temperature, rain fall of study area should also added

Comment [T8]: Add about age of animal whether it is similar or not, body condition and management system of animal

Identification of parasites

Identification of parasites was done using bench aid for intestinal parasites (WHO, 1994). Faecal egg counts (FEC) were determined by the modified McMaster technique (MAFF, 1986). Distinguishable nematode eggs, trematode and cestode eggs were identified directly.

Statistical analysis

The data generated were subjected to statistical analysis using statistical package for social sciences (SPSS) version 26. Chi-square analysis was used to determine the statistical difference between the variables such as gender and breed. Statistical difference was based on $p < 0.05$.

Result

Prevalence of gastrointestinal parasites from different goat breeds

The number and percentage of different goat breed samples collected from two different abattoirs was presented in Table 1, while the prevalence of gastrointestinal parasite infection among different goat breeds was represented in Table 2. Out of 240 examined faecal samples collected for gastrointestinal parasites study, 56 (92.81%) were positive of gastrointestinal parasite infection. The goat breed with the highest prevalence was Sokoto red breed (34.38%), followed by Kano brown breed (21.82%), then Borno red breed (19.23%) and the least gastrointestinal parasites prevalence was recorded with West African dwarf goat breed (17.38%). The distribution of parasite ova in all the goat breeds showed no significant difference ($\chi^2 = 6.284, p \text{ value} = 0.099$). The prevalence of gastrointestinal parasites infection between goat breeds in all faecal samples examined showed that Sokoto red breed had the highest parasites prevalence (34.38%) and West African dwarf breed had the least prevalence (17.38%) as shown in Table 2.

Table 1: Number and percentage of different goat breeds examined for gastrointestinal parasites from two abattoirs in Akure metropolis.

Comment [T9]:

Comment [T10]: How 56 (92.81%)??? If you divide $56/240 \times 100 = 23.33\%$

Formatted: Superscript

Comment [T11]: It is already mentioned, please remove redundant idea

Comment [T12]: Rephrase it by removing redundancy and don't write again table what was already written in text. Write only either in text or table.

Goat Breed	Abattoir 1 (%)	Abattoir 2 (%)	Total (%)
Sokoto red	32(26.67%)	32(26.67%)	64(26.67%)
Kano brown	29(22.92%)	26(22.92%)	55(22.92%)
Borno red	27(21.67%)	25(21.67%)	52(21.67%)
West African dwarf	32(28.75%)	37(28.75%)	69(28.75%)
Total	120 (50%)	120(50%)	240

Table 2: Prevalence of infection among different goat breeds in the study area

Goat breed	Number examined	Number infected	Prevalence (%)
Sokoto red	64	22	34.38%
Kano brown	55	12	21.82%
Borno red	52	10	19.23%
West African dwarf	69	12	17.38%
Total	240	56	23.33%

Chi-square (χ^2) = 6.284, p value = 0.099, degree of freedom (df) = 3

Prevalence of gastrointestinal parasites recovered from two different abattoirs from the study area

As shown in Table 3, abattoir II had the highest prevalence of parasites of 55.46%; while abattoir I had the least prevalence of 44.44%, there is a significant statistical difference (p value =0.013) in prevalence of gastrointestinal parasites comparing the two abattoirs (I and II). Eight (8) parasites of

Comment [T13]: Don't write again table what was already written in text. Write only either in text or table.

Comment [T14]: How you related prevalence of parasite with abattoir? Your materials and methods are not supporting this result. Rather you can relate with animal slaughtered at different abattoir if there is difference in origin of animal that brought to two abattoirs. It should be rewritten by avoiding those mentioned errors.

various genera were encountered in the faecal samples examined (Table 3 and Figure 1). These include; *Strongyloidesstercoralis*, *Ascarislumbricoides*, *Fasciola* spp., *Haemonchus contortus*, *Ancylostomaduodenale*, *Trichostrongylus*spp., *Diphyllobothriumlatum*, and *Moniezia*spp. In Abattoir I, *Strongyloidesstercoralis*was the most prevalent (25.00%) intestinal parasite encountered followed by *Ancylostomaduodenale* (17.86%), while *Moniezia*spp. (7.14%), *Haemonchus contortus* (7.14),and *Trichostrongylus* spp. (7.14%) were the least prevalent gastrointestinal parasites observed. In abattoir II, *Strongyloidesstercoralis*was the most prevalent(45.71%) intestinal parasite encountered followed by *Ancylostomaduodenale*(20.00%), and *Ascaris lumbricoides* (20.00%), *Fasciola* spp., *Haemonchus contortus*, *Moniezia*spp. and *Trichostrongylus*spp. was not encountered in abattoir II as shown in Table 3. The parasite was highest prevalence was *Strongyloidesstercoralis* and the least prevalent parasite were *Monieziaexpansa*, *Trichostrongylus*spp. and *Haemonchus contortus* (Figure 1).

Table 3: Prevalence of parasites recovered from the studied Abattoirs

Parasite	Number of goat and prevalence infections		Total (%)
	Abattoir I (%)	Abattoir II (%)	
<i>S. stercoralis</i>	7 (25.00)	16 (45.71)	23 (36.51)
<i>D. latum</i>	3 (10.71)	5 (14.29)	8 (12.70)

Comment [T15]: Relating prevalence of parasite with abattoir is not relevant. So that, try to relate with goat slaughtered at different abattoir.

<i>A. duodenale</i>	5 (17.86)	7 (20.00)	12 (19.05)
<i>A. lumbricoides</i>	4 (14.29)	7 (20.00)	11 (17.46)
<i>M. expansa</i>	2 (7.14)	0	2 (3.18)
<i>H. contortus</i>	2 (7.14)	0	2 (3.18)
<i>Fasciola</i> spp.	3 (10.71)	0	3 (4.76)
<i>Trichostrongylus</i> spp	2 (7.14)	0	2 (3.18)
Total	28 (44.44)	35 (55.46)	63




Figure 1: Prevalence of gastrointestinal parasites recovered from goat

Table 4: Prevalence of parasite infection male and female different goat breed

Goat breed	Male		Female		Total number examined	Total number infected (%)
	Number examined	Number infected (%)	Number examined	Number infected (%)		

Sokoto red	30	8 (26.67)	34	14 (41.18)	64	22 (34.38)
Kano brown	33	5(15.15)	22	7 (31.89))	55	12 (21.81)
Borno red	32	5(15.63)	20	5(25.00)	52	10(19.23)
West African dwarf	30	3(10.00)	39	9(23.07)	69	12(17.39)
	125	21(17.07)	115	35(30.43)	240	56(23.33)

Chi-square (χ^2) = 6.225, p value = 0.013, degree of freedom (df) = 1

Prevalence of gastrointestinal among goat gender from the study area

Table 4 present the prevalence of gastrointestinal parasites among the goat breed gender. Female goat (30.43%) had the highest prevalence of gastrointestinal parasites infection compare to their male counterpart (17.07%). There was a significant difference ($p = 0.013$) comparing the infection rate in relation to gender. Among the breed, female Sokoto red goat breed was found to have highest prevalence of gastrointestinal parasites of 41.18% followed by female Kano brown goat breed (21.81%). Likewise, among the male, the male Sokoto red goat breed has the highest gastrointestinal parasites of 26.67% followed by male Kano brown goat breed (15.15%) and male Borno red goat breed (15.15%).

Comment [T16]: Move this statement before table 4 and avoid redundancy of what is written in text in table

Discussion

With regard to the level of parasitic infection observed during this study; the prevalence of gastrointestinal parasites revealed that *Strongyloides* spp. was the highest prevalent parasite, this was similar to the reported of Odoi *et al.* (2007). The high prevalence of *Strongyloides* in abattoir II can be associated with the environment in which the goats were being reared, and also by poor animal management. Goats were reared in mixed crop-livestock systems, where a few numbers of goats were herded together in the same area during the dry and wet seasons. This system of rearing could

Comment [T17]: How? Not included in your methods and result.

cause high rate of parasitic infection due to possibilities of re-infection in contaminated pastures. Other factor that might likely associated with the prevalence of gastrointestinal parasitic infection in goats could be frequent access to communal grazing and browsing of forages. However, lower infections in abattoir I could be associated with improvement in management in terms of housing, feeding and healthcare for the animals.

This study revealed the presence of eight gastrointestinal parasites which include *Strongyloides* spp., *Ascaris* spp., *Haemonchus* spp., *Trichostrongylus* spp., *Moniezia* spp., *Ancylostoma* spp. and *Fasciola* spp. This agrees with the findings of Gadahiet *et al.* (2009) who reported that these parasites are the most pathogenic gastrointestinal parasites of small ruminants.

Prevalence of *Fasciola* spp. was low, this was in accordance with the report of Khanjari *et al.* (2014). However, infections may have been low in goats due to their foraging behaviour, which minimizes chances of ingesting the metacercaria which are found on plants closer to the ground. The only cestode observed in the study area was *Moniezia* spp. The occurrence of this parasite is associated with the ingestion of mites infected with cysts of *Moniezia* spp. (Diop *et al.*, 2015). High prevalence of *Haemonchus* and *Trichostrongylus* have been recorded in other studies (Tsoetsi and Mbat, 2003; Ayaz *et al.*, 2013; Tsoetsi *et al.*, 2013). The high and low in different gastrointestinal parasite species could be due to different geographical location, host factors and climatic conditions required for the development of free-living stages of these parasites.

The result of present study revealed that female had highest prevalence compared to male, this higher prevalence in females maybe due to the physiological condition of the animals during pregnancy and lactation (production activity) and also the absence of sufficient food required for production which may lead to the lowering of the body resistance of the females, this findings was similar to the findings of Emiruet *et al.* (2013) and Vieira *et al.* (2014) in Ethiopia and Brazil, where females were more susceptible to parasite infection than males, this was attributed to lowered resistance of female animals due to their reproductive events and insufficient/unbalanced diet against higher needs. However, these findings were contrary to the finding of Tariq *et al.* (2008), Ayaz *et al.* (2013) and Nabi *et al.* (2014), as

Comment [T18]: Communal grazing , browsing of forages and management in terms of housing, feeding and healthcare for the animals were not your study variables. How you can discuss about it?

well as Badaso and Addis (2015), who reported that males of all ages had high prevalence than females and attributed to the genetic predisposition and differential susceptibility owing to hormonal control. The disparity in both findings can be attributed to different in geographical locations where these studies were conducted.

Sokoto red goat breed had the highest prevalence while West African dwarf goat breed had the lowest prevalence, this finding is in agreement the report of Ovutoret *al.* (2014) and Adejinmiet *al.*(2015); who in their respective studies reported that helminth parasites are more prevalence in exotic breeds than indigenous breeds because of their efficient life cycle ranging from the very simple to the extremely complicated stage. The high prevalence in Sokoto red goat breed might be due to the system of management that these goats were subjected to as they were always left to wander about scavenging and feeding indiscriminately on anything they come in contact with and then return to their poorly kept and unhygienic ranches.

Conclusion

The prevalence rate of helminthiasis in livestock need to be put to check periodically. The result of this research will be of great assistance in understanding the epidemiology of the gastrointestinal parasites of goats in Akure and its environs. The result of this research revealed that gastrointestinal parasites are prevalence in small ruminants in Akure abattoirs. This study will assist veterinarians and livestock workers as they should certify that only healthy small ruminants examined are slaughtered. It will also help farmers in organizing animal husbandry system, maintenance of proper health, feeding and sanitary condition, deworming, towards maximum productivity goat meat in the study area. It is recommended that good management practices, prompt diagnosis, and treatment with anthelmintic and anti-protozoa drugs be implemented to reduce the risk of gastrointestinal infections of goats and other ruminants.

Comment [T19]: This variable was not statistically significant, so you have evidence to discuss and justify it.

Comment [T20]: You have to clearly conclude with main finding of your study rather than making general like this.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All analysed data involved in this study are included in this manuscript

Reference

- Adejinmi, O. O., Adejinmi, J. O., Falohun, O. O., Aderoju, O. R., Dauda, W. J., 2015. Prevalence of gastrointestinal parasites of goats in Ibadan, Southwest, Nigeria. *World J. Agric. Res.* 3, 49-51.
- Anaeto, M.B, Tayo, G.I., Chioma, G.O., Afolabi, A.A., 2009. Comparative study of Albendazole and *C. papaya* seed on the control of gastrointestinal nematodes in goats. *J.lif.Phy. Sci.* 3, 25-28.
- Ayaz, M.M., Raza, M.A., Murtaza, S., Akhtar, S., 2013. Epidemiological survey of helminths of goats in southern Punjab, Pakistan. *Trop. Biomed.*30, 62-71
- Badaso, T., Addis. M., 2015. Small ruminants haemonchosis: prevalence and associated risk factors in Arsi Negelle municipal abattoir, Ethiopia *Glob. Vet.* 15, 315-320
- Barger, I. A., 1997. Control by Management. *Vet.Parasitol*, 72, 493 – 506
- Diop, G., Yanagida,T., Hailemariam,Z., Menkir, S., Nakao, M., Sako, Y., Ba, C. T., Ito, A., 2015. Genetic characterization of *Moniezia* species in Senegal and Ethiopia. *Parasitol. Int.* 64, 256-260
- Emiru, B., Amede,Y., Tigre, W., Feyera,T., Deressa, B., 2013. Epidemiology of gastrointestinal parasites of small ruminants in Gechidistrict, Southwest Ethiopia. *Adv. Biol. Res.*7, 169-174

- Fissiha, W., Kinde, M. Z., 2021. Anthelmintic Resistance and Its Mechanism: A Review. *Infect. drug resist.* 14, 5403–5410
- Gadahi, J. A., Arshed, M. J., Ali, Q., Javaid, S. B., Shah, S. I., 2009. Prevalence of gastrointestinal parasites of sheep and goat in and around Rawalpindi and Islamabad. *Pak. Vet. World.* 2, 51-53.
- Gunathilaka, N., Niroshana, D., Amarasinghe, D., Udayanga, L., 2018. Prevalence of Gastrointestinal Parasitic Infections and Assessment of Deworming Program among Cattle and Buffaloes in Gampaha District, Sri Lanka. *BioMed Res.Inter.* 2018, 3048373.
- Khanjari, A., Bahonar, A., Alizadeh, A., Fallah, S., Bagheri, M., Fallah, M., Khanjari, Z., 2014. Prevalence of fasciolosis and dicrocoeliosis in slaughtered sheep and goats in Amol abattoir, Mazandaran, Northern Iran *Asian Pac. J. Trop. Dis.* 4, 120-124
- MAFF (Ministry of Agriculture, Fisheries and Food) (1986) *Manual of Veterinary Parasitological Laboratory Techniques*, ADAS, HMSO, UK.
- Mohammed, T., Tsegaw, F., Bushura, R., 2019. Veterinary Drug Residue: The Risk, Public Health Significance and its Management College of Veterinary Medicine and Animal Science. *Dairy and Vet Sci J.* 13, 555856
- Nabi, H., Saeed, K., Shah, S.R., Rashid, M.I., Akbar, H., Shehzad, W., 2014. Epidemiological study of gastrointestinal nematodes of goats in district Swat, Khyber Pakhtunkhwa, Pakistan *Sci. Int.*, 26, 283-286
- Nwosu, C., Madu, P., Richards, W., 2007. Prevalence and seasonal changes in the population of gastrointestinal nematodes of small ruminants in the semi-arid zone of north-eastern Nigeria. *Vet. Parasitol.* 144, 118-124

- Odoi, A., Gathuma, J.M., Gachui, C.K., Omere, A., 2007. Risk factors of gastrointestinal nematode parasite infections in small ruminants kept in smallholder mixed farms in Kenya *BMC Vet. Res.* 3, p. 6
- Onaga, T., Hara, N. and Shimizu, Y., 2009. Role of nitric oxide in the assessment of genetic variability of resistance to gastrointestinal nematode parasites in Creole goats in the humid tropics. *J. Ani. Sci.* 79, 1706-1712
- Osakwe, I. I., Anyigor, S. I., 2007. Prevalence of gastrointestinal helminths in west African dwarf (wad) goats in an Agrarian agro-ecosystem. *Ani. Res. Inter.* 4, 728 – 732
- Ovutor, O., Elele, K., Gboeloh, L. B. B., 2014. Prevalence of Gastrointestinal Helminths in Exotic and Indigenous Goats Slaughtered in Selected Abattoirs in Port Harcourt, South-South, Nigeria. *Chin. J. Biol.* 2014, 1 – 8
- Perry, B. D., Randolph, R. F., McDermott, J. J., Sones, K. R., Thomson, P. K., 2002. Investigating in Animal Health Research to Alleviate Poverty. International Livestock Research Institute, Nairobi, Kenya, 148 pp.
- Tariq, K., Chishti, M., Ahmad, F., Shauk, A., 2008. Epidemiology of gastrointestinal nematodes of sheep managed under traditional husbandry system in Kashmir valley. *Vet. Parasitol.*, 158, 138-143
- Tsotetsi, A., Mbatia, P., 2003. Parasitic helminths of veterinary importance in cattle, sheep and goats on communal farms in the Northeastern Free State, South Africa *J. S. Afr. Vet. Assoc.* 74, 45-48
- Tsotetsi, A.M, Njiru, S., Katsande, T.C., Mayo, G., Baloyi, F., 2013. Prevalence of gastrointestinal helminths and anthelmintic resistance on small-scale farms in Gauteng Province, South Africa. *Trop. Ani. Health Prod.* 45, 751–761.

Vieira, V.D., Feitosa, T.F., Vilela, V.L.R., Azevedo, S.S., Ribeiro, A.R.C., 2014. Prevalence and risk factors associated with goat gastrointestinal helminthiasis in the sertão region of Paraíba state, Brazil Trop. Anim. Health Prod. 46, 355-361

WHO, 1994. Bench Aids for the Diagnosis of Intestinal Parasites. World Health Organization, Geneva.

UNDER PEER REVIEW