

**Note:-**

**Elongate all the topics and write everything in a detailed manner**

**This article has everything in brief manner**

**Try to elongate it and add valuable points .**

**Also add references after 2020**

**Make sure every references you used here is present in the manuscript**

**Italicise all the scientific names**

**Comment [DCKP1]:** My Comments

**Influence of abiotic factors on efficacy of Entomopathogenic nematodes**

**Comment [DCKP2]:** Capital A - Abiotic

**Comment [DCKP3]:** If possible u can have minor changes in the title of the manuscript

**Abstract:**

Entomopathogenic nematodes (EPNs) are potential biocontrol agents against many insect pests. Abiotic factors such as temperature, moisture, soil texture can affect the survival, growth and development, and reproduction of entomopathogenic nematodes and ultimately their efficacy against insect pests.

**Comment [DCKP4]:** Add more points in the Abstract .  
Elongate it to about 250-300 words.

**Key words:** Entomopathogenic nematodes (EPNs) abiotic factors, biocontrol agents, survival, movement, reproduction, bioefficacy.

**Comment [DCKP5]:** Arrange it according to alphabetical order .

**Introduction**

Entomopathogenic nematodes in the genera *Steinernema* and *Heterorhabditis* are well known biocontrol agents against a wide range of insect pests. However, their efficacy greatly depends on their host range, host finding strategy, and above all their tolerance of environmental factors. EPN live in soil for short or long durations. The soil has a highly complex physical, chemical and biotic environment that responds to external environment as well. Thus soil environment influences the entomopathogenic nematode population.

**Comment [DCKP6]:** Genera

**Comment [DCKP7]:** Elongate it .it should be in concise manner .

**Response of Entomopathogenic nematodes to various abiotic factors**

**Temperature:**

The optimum temperatures for survival, movement, infection, and reproduction of EPNs vary among species and strains (Hazir *et al.*, 2001). IJs are non feeding, relying for energy on their internal lipid reserves, which vary in composition amongst EPN species. Dauers and IJs have lower levels of metabolic activity than other stages of their life-cycle. At higher temperatures, the infective juveniles have increased metabolic activity and deplete their energy reserves, shortening their life span. The low temperature induce inactivity in infective juveniles, such inactivity is characterized by decreased enzymatic activity and mobility, both reducing metabolic expenditures. Extended exposure to temperature extremes (below 0°C or above 40°C) is lethal to most species of entomopathogenic nematodes. In the soil environment, infective juveniles are normally buffered from temperature extremes. Extreme ambient temperatures can lead to desiccation, increased metabolic rate and consequently more rapid use of energy reserves by EPNs, which leads to shorter generation times, decreased infectivity and lowered virulence of such nematodes (Grewal *et al.*, 1994). For storage, the best longevity of infective juveniles is between 5 and 15°C. Temperature below 10-15°C can restrict their mobility, while temperatures higher than 30-40°C can inactivate them. It is generally known that steinernematid species are more active at lower temperatures than heterorhabditid species. The proteins which decreased in abundance to the greatest extent in *S. carpocapsae* after both 9°C and 20°C storage were those associated with metabolism, stress and the cytoskeleton. Oxidative phosphorylation produces the most energy of all metabolic processes, but it also produces detrimental reactive oxygen species. Free radicals damage cells and contribute towards aging. Catalase, the stress enzyme which breaks down hydrogen peroxide, and copper oxide dismutase were both increased in abundance after storage at low temperature. IJs may enhance their survival in colder conditions by reducing metabolic activities which produce these free radicals and increasing their stress response against them. However, some *Steinernema* species are well-adapted to warm temperatures. It has been observed that the original climatic locality of the nematodes correlated with their temperature range (Kung *et al.* 1991). The heat survival by IJs is related to the presence of heat shock proteins (HSP). *H. bacteriophora*(IS-5) which was isolated from the Negev Desert exhibits heat tolerance (Glazer, 2002). Shapiro *et al.* (1997) improved a heat tolerant hybrid strain from *H. bacteriophora*(IS-5) and *H. bacteriophora*(Hp88) using classical genetic methods. The hybrid strain killed *G. mellonella* larvae at 32°C. In general, freezing temperatures can be tolerated better by the IJs than high temperatures (above 37°C) (Shapiro-Ilan *et al.*, 2014). To survive at freezing

Comment [DCKP8]: ?????

Comment [DCKP9]: First use the full form and then use shortcut form .

Comment [DCKP10]: Italicise it

Comment [DCKP11]: Italicise and check spelling .

temperatures, IJs use three mechanisms; (1) freeze-avoiding: The sheath of *H. zealandica* prevents the IJ from supercooling (-32°C) (Wharton and Surrey, 1994). (2) freezing tolerant (Brown and Gaugler, 1997), or (3) cryoprotective dehydration (Wharton *et al.*, 2003). *S. feltiae* and *H. bacteriophora* can exhibit freezing tolerant and cryoprotective dehydration. Wright (1992) reported that reproduction rate of many entomopathogenic nematodes at low temperature was related to the growth rate of the bacterial symbiont. Tailliez *et al.* (2006) suggested that some *Xenorhabdus* species may have become adapted to tropical or temperate regions and/or influenced by the optimal growth and development temperatures of their nematode host.

Table.1. Effect of temperature on survival, reproduction and infectivity of Entomopathogenic nematodes

Entomopathogenic nematodes	Survival	Reproduction	Infectivity	References
<i>Steinernema glaseri</i>	15-30°C	28°C	15-35°C 10-32°C 0-25°C	Kung <i>et al.</i> , 1991; Grewal <i>et al.</i> 1994; Jagdale & Gordon, 1997
<i>S. carpocapsae</i> <i>S. carpocapsae</i> (Pocheon) <i>S. carpocapsae</i> (PDBCEM 6.11)	9 - 20°C	20-24°C	10-32°C 24°C 5-25°C 25-32°C	Grewal <i>et al.</i> , 1994, Saunders & Webster, 1999; Brown <i>et al.</i> , 2002; Yul <i>et al.</i> , 2002
<i>S. feltiae</i>	15°C	15-25°C	10-15°C 10°C	Grewal <i>et al.</i> , 1996; Hazir <i>et al.</i> , 2001; Chen <i>et al.</i> , 2003; Radova & Trnkova, 2010 Flores <i>et al.</i> , 2021
<i>S. anomali</i> (= <i>S. arenarium</i> ),	can tolerate sub-zero temperatures	23°C	23°C	Kakouli-Duarte & Hague, 1999; Ali & Wharton, 2013
<i>S. riobrave</i>	35 °C	20-35°C	10-39° C 29°C	Cabanillas <i>et al.</i> , 1994; Ali & Wharton, 2013
<i>S. siamkayai</i> Thailand strain <i>S. siamkayai</i> Tiruchirappalli strain	25-30 °C	25-30°C 20-35°C	25-30°C 37.5°C	Chongchitmatee <i>et al.</i> 2005; Tailliez <i>et al.</i> , 2006
<i>S. scapterisci</i>	In warm climates	20-32 °C	30°C	Hazir <i>et al.</i> 2003
<i>H. bacteriophora</i>	-19 -30°C	15 - 30°C	10°C- 35°C	Mason & Hominick 1995; Ali & Wharton, 2013;

				Milstead ,1981
<i>H. zealandica</i>	15-30°C	15-30°C	15-30°C	Waal <i>et al.</i> , 2018; Aryal <i>et al.</i> , 2022
<i>H. megidis</i>	9- 20°C	15-25°C	15°C Equal to or higher than 29°C	Dziegielewska <i>et al.</i> ,2023; Kung <i>et al.</i> , 1991; Grewal <i>et al.</i> , 1994; Berry <i>et al.</i> , 1997
<i>H. merelatus</i>	Warm climate	Equal to or higher than 29°C	Equal to or higher than 29°C	Kung <i>et al.</i> , 1991 Grewal <i>et al.</i> , 1994, Berry <i>et al.</i> , 1997

**Comment [DCKP12]:** If possible add reference after 2020

### Moisture:

Moisture is the most important factor influencing EPN populations (Shapiro-Ilan *et al.*, 2007; Campos-Herrera *et al.*, 2016). EPN require adequate soil moisture levels for their survival, locomotion and infectivity, which may vary among nematode species and isolates in different soil types (Glazer 2002). Moisture also influences aeration of soil. Some species develop survival strategies under water stress conditions, by reducing the body surface area exposed to the air and by lowering their rate of metabolism. This process, known as anhydrobiosis, allows the nematode to become resistant to desiccation. Anhydrobiosis can be reversed by wetting the soil, causing a recovery of nematode infectivity and virulence. Gradual water removal from the infective juveniles gives them time to adapt to the desiccating conditions. Thus, natural soils allow the infective juveniles to persist longer in dry soil. Infective juveniles may survive desiccating conditions by remaining inside the host cadaver until the soil moisture situation improves (Grewal 2000). Some species of the *Steinernema* have the ability to enter a state of anhydrobiosis when exposed to low moisture contents (Koppenhöfer *et al.*, 1995). Nematodes need a water film in the interstitial spaces of soil for effective movement. When this water film becomes too thin (i.e. in dry soil) or the interstitial spaces become completely filled with water (i.e. in saturated soil), oxygen levels decrease and nematode movement is restricted (Stuart *et al.*, 2015). Several studies indicate that soil moisture influence infectivity of entomopathogenic nematodes, Nematode infectivity is highest at moderate soil moistures (-10 to -100 kPa), and lower in wet (-1 kPa) and moderately dry (-1000 kPa) soil (Koppenhofer and Fuzy, 2007). *S. carpocapsae* and *S. glaseri* survival was good at low soil moisture of 2 and 4% respectively (Kung *et al.*, 1991) In general, a decrease in infectivity as soil moisture decreases is

that these nematodes have entering a state of anhydrobiosis, but it can be increased by rehydrating the soil using irrigation or rainfall (Alekseev *et al.*, 2006). Grant and Villani (2003) showed that infectivity of *H. bacteriophora*, *S. carpocapsae*, *S. feltiae*, and *S. glaseri* increased with soil moisture in sandy loam soils. Similarly, *H. bacteriophora* DI and *S. glaseri* KG (Molyneux and Bedding 1984) and *S. glaseri* NC and *S. carpocapsae* ALL (Koppenhöfer *et al.*, 1995) showed lower infectivity in extreme, low and high (near the saturation point) soil moistures. Studies have also shown that *H. bacteriophora* is less susceptible on *Popillia japonica* 3rd instar as the season changes from rainfall to spring (Paoli *et al.*, 2017). Under high humidity and moderate temperature, *S. virgalemense* is highly susceptible to female *Planococcus ficus* and mortality rate of *P. ficus* decreases with decreasing relative humidity levels (Platt *et al.*, 2018). Suboptimal moisture levels can be beneficial when formulating nematodes. Water activity in the formulation can gradually be reduced to a point where the nematode enters a state of partial anhydrobiosis where nematode's metabolism ceases which leads to a lowered use of energy reserves, increasing the life span of the nematode under suboptimal moisture levels. Pre-application of water rehydrates formulated nematodes and they resume function as virulent individuals, ready to be applied in the field.

#### **Soil texture and structure**

Texture of soil is decided by the proportion of sand, silt and clay inherited by it during its formation. Soil structure determines the number, size and distribution of pore spaces in soil crumbs. These pores are partly or wholly filled with soil moisture. Soil texture and structure can have various effects on EPNs and they should be evaluated for different species. Movement (Kapranas *et al.*, 2021), infectivity of IJs (Toledo *et al.*, 2009) and persistence (Koppenhöfer and Fuzy, 2006) are positively correlated if the space among soil particles is similar or greater than the diameter of an IJ (Stuart *et al.*, 2015). In general, the most favorable soil type for nematode movement is sandy soils, whereas clay soils have poorest survival rates because of low oxygen levels in the small soil pores (Hazir *et al.*, 2003). Besides soil mineral component, the proportion of organic media in the soil affects nematode movement. *H. bacteriophora*, *S. carpocapsae*, and *S. glaseri* survival and movement was more in sandy loam than silty clay loam or loamy soils. While survival of *H. bacteriophora* decreased, survival of *S. glaseri* increased with bulk density. Furthermore, the bulk density increased the infectivity of *S. carpocapsae*, but did not vary for *H. bacteriophora* and *S. glaseri*. Koppenhofer *et al.*, (1998) reported that establishment of *S.*

*scarabaei* was more in sandy loam soil compared to clay loam soil. However, this may vary with EPNs species and isolate. Campos-Herrera and Gutiérrez (2014) observed that infection dynamics varied among 14 different populations of *S. feltiae* in the same soil texture. Shelmithet *al.*, (2007) reported that survival of *S. kari* was 22-67%, with the highest occurrence being in clay soil, moderate in sandy clay and clay loam and the lowest being in the sandy clay soil. Campos-Herrera and Gutiérrez (2009) observed that heavy soils negatively affected the virulence of the Rioja isolate of *S. feltiae* on larvae of *Spodopteralittoralis* as an increase content of clay from 5 to 14%, caused an increase in the LC90 (to kill 90% of larvae in two days) from 220 to 4,178 IJs/cm<sup>2</sup>. Organic matter (OM) content of soil improves soil structure, lowering bulk density, increasing the available space through the distribution of aggregates and pore size, thus OM content and soil compaction may influence EPNs infection patterns (Kapranaset *al.*, 2017), and persistence in the soil (Shapiro and Lewis, 2018). Therefore, knowledge about soil requirements of an EPN species/isolate is critical to optimizing their performance as biological control agents in the field.

### **Salinity**

Soil salinity has a negligible effect on infective juvenile survival even at salinity above the tolerance levels of most crop plants. Seawater has no negative effects on the survival of several *Heterorhabditis* species/strains as they have been frequently isolated from soils near the seashore. However, tillage and herbicide applications suppress EPN numbers, the movement of EPNs is restricted and their ability to find and recognize their host is reduced. Bednarek (1998) observed that mineral fertilizers negatively affected EPN population density. EPNs appear to tolerate a wide range of salinities. CaCl<sub>2</sub> and KCl had no effect on the survival, penetration efficiency, or movement through a soil column for *H. bacteriophora* but moderate concentrations of these salts enhanced the virulence. Although seawater has no negative effect on IJs, high concentrations of NaCl, CaCl<sub>2</sub>, and KCl inhibit the infectivity and movement of *S. glaseri* (Grant and Villani, 2003).

### **Oxygen and pH**

Oxygen is also a limiting factor in water-saturated soils and soils with high organic matter content. Soil moisture and aeration affect the soil pH. Low oxygen level that can occur in water-saturated soils, clay soils (because of small pores) and soils with high organic matter content. IJs can survive in a wide range of pH. Soil pH between 4 and 8 did not have an adverse effect on IJ

survival, but a pH of 10 or higher was harmful. Kung *et al.*, (1990) observed in laboratory trials that pH 10 reduced steinernematid nematodes survival. *S.bicornutum* was found only in neutral soils, and *S. silvaticum* only in acidic environments (Kanga *et al.*, 2012). As far as EPNs as a formulated product is concerned, the presence of interstitial spaces in formulations is beneficial to such nematodes, as it supports gas exchange .

### UV radiation

The survival, virulence, or reproduction of EPNs can be affected by UV radiation. Shapiro-Ilan *et al.* (2015) investigated UV tolerance of nine different EPN species and fifteen strains for survival and virulence. Their results showed that there was a significant variation in UV tolerance among EPN strains and species. Steinernematids commonly displayed higher levels of UV tolerance compared with the heterorhabditids. Furthermore, it has been reported that shorter wavelengths (254 nm) had a negative effect on virulence compared to longer wavelengths (366 nm).

### Conclusion

A better understanding of ecological strategies of native nematodes for its efficiency in biological control in the local agro-climatic conditions is needed. Therefore, a means to lower the impact of temperature or moisture on EPN efficacy as biological agents should be enforced, by choosing the right EPN strain and applying these nematodes at the right time of the year, or postponing application until temperatures or moisture are optimal for the specific EPN strain used. Therefore isolation and description of new species and described species still need to be studied in detail for their potential use as biological control agent.

Comment [DCKP13]: Add more points .

### References

- Alekseev E., Glazer I., Samish M. 2006. Effect of soil texture and moisture on the activity of entomopathogenic nematodes against female *Boophilus annulatus* ticks. *Biocontrol*.51:507-518.
- Ali F., Wharton DA. 2013. Cold tolerance abilities of two entomopathogenic nematodes, *Steinernema feltiae* and *Heterorhabditis bacteriophora*. *Cryobiology*.66(1):24-9. doi:10.1016/j.cryobiol.2012.10.004.
- Aryal S., Nielsen UN., Sumaya NH., Wilson C., Riegler M. 2022. Effect of temperature on survival of Australian entomopathogenic nematodes and their virulence against the Queensland fruit fly, *Bactrocera tryoni*. *BioControl*.67:617-628.

- Bednarek A. 1998. The agricultural system, as a complex factor, effects the population of entomopathogenic nematodes (Rhabditida: Steinernematidae) in the soil. IOBC Bulletin 21:155-216.
- Berry RE, Liu J, Groth E 1997. Efficacy and persistence of *Heterorhabditismareletus* (Rhabditida: Heterorhabditidae) against root weevils (Coleoptera: Curculionidae) in strawberry. Environ Entomol 26: 465-470.
- Brown, I., Lovett, B., Grewal, P. S., & Gaugler, R. 2002. Latent infection: A low temperature survival strategy in steinernematid nematodes. Journal of Thermal Biology. 27: 531-539
- Brown, IM., Gaugler, R. 1997. Temperature and humidity influence emergence and survival of entomopathogenic nematodes. Nematologica. 43: 363-375.
- Cabanillas HE., Poinar GO., Raulston JR. 1994. Steinernemariobravisp. (Rhabditida: Steinernematidae) from Texas. Fundamental and Applied Nematology. 17:123-131.
- Campos-Herrera R., Gutiérrez C. 2009. Screening Spanish isolates of steinernematid nematodes for use as biological control agents through laboratory and greenhouse microcosm studies. Journal of Invertebrate Pathology. 100: 100-105.
- Campos-Herrera R., Gutiérrez C. 2014. Steinernema feltiae intraspecific variability infection process and sex-ratio. Journal of Nematology. 46: 35-45.
- Campos-Herrera R., Fahiem E., El-Borai., Martin JAR., Duncan LW. 2016. Entomopathogenic nematode food web assemblages in Florida natural areas. Soil Biology and Biochemistry. 93: 105-114.
- Chen, S., Li J., Han and Moens, X.M. (2003). Effect of temperature on the pathogenicity of entomopathogenic nematodes (Steinernema and Heterorhabditis spp.) to Delia radicum. Biocontrol., 48 : 713-724.
- Chongchitmate P., Somsook V., Hormchan P., Visarathanonth N. 2005. Bionomics of entomopathogenic nematode Steinernemasiamkayai Stock, Somsook and Reid (n.sp.) and its efficacy against Helicoverpa armigera Hubner (Lepidoptera: Noctuidae). Kasetsart J. 39:431-439.
- Dziegielewska M., Kaczmarek K., Kauk K. 2023. The influence of temperature on the biological activity of selected nematode species (Steinernematidae and Heterorhabditidae) under the conditions of their coexistence. Plant Protection Science 59(2): 193-201

- Flores P., Alvarado A., Paola Lax GL., Prodan S., Aballay E.2021. Morphological,molecular and ecological characterization of a native isolate of *Steinernemafeltiae* (Rhabditida: Steinernematidae) from Southern Chile. Parasites and Vectors.14:45.<https://doi.org/10.1186/s13071-020-04548-7>.
- Glazer I. 2002. Survival biology, p. 205-220. In: Gaugler R (ed) Entomopathogenic nematology. Wallingford, CABI Publishing UK, 400p.
- Grant JA and Villani MG 2003. Soil moisture effects on entomopathogenic nematodes. Environ Entomol.32:80-87.
- Grewal BPS, Gaugler R., Wang YI.1996. Enhanced cold tolerance of the entomopathogenic nematode *Steinernemafeltiae* through genetic selection. Ann Appl Biol.129:335-41.
- Grewal, PS., Selvan, S., Gaugler,R.1994. Thermal adaptation of entomopathogenic nematodes: Niche breadth for infection, establishment, and reproduction. Journal of Thermal Biology. 19(4): 245-253.
- Hazir S., Keskin N., Stock SP.,Kaya H., Ozcan S.2003.Diversity and distribution of entomopathogenic nematodes (Rhabditida: Steinernematidae and Heterorhabditidae) in Turkey. Biodiversity and Conservation.12:375-386.
- Hazir S., Stock SP., Kaya HK., Koppenhofer AM., Keskin N.2001.Developmental temperature effects on five geographic isolates of the entomopathogenic nematode *Steinernemafeltiae* (Nematoda: Steinernematidae). J Invertebr Pathol.77:243-50.
- Jagdale, G.B. and Gordon, R. (1997). Effect of recycling temperature on the infectivity of Entomopathogenic nematodes. Canadian J. Zool., 7: 2137-2141.
- Kakouli-Duarte T and Hague NG.1999. Infection, development , and reproduction of the entomopathogenic nematode *Steinernemaarenarium* (Nematoda: Steinernematidae) in the black vine weevil *Otiorhynchussulcatus* (Coleoptera: Curculionidae) . Nematology.1(2): 149-156.
- Kanga FN., Waeyenberge L., Hauser S., Moens M.2012. Distribution of entomopathogenic nematodes in Southern Cameroon.Journal of Invertebrate Pathology.109: 41-51.
- Kapranas A., Chronopoulou A., Lytra IC., Peters A., Milonas PG.2021.Efficacy and residual activity of commercially available entomopathogenic nematode strains for Mediterranean

- fruitfly control and their ability to infect infested fruits. *Pest Management Science*. 77: 3964-3969.
- Koppenhofer AM and Fuzy EM. 2007. Soil moisture effects on infectivity and persistence of the entomopathogenic nematodes *Steinernemascarabaei*, *S. glaseri*, *Heterorhabditiszealandica*, and *H. bacteriophora*. *Appl Soil Ecol* .35: 128-139.
- Koppenhöfer AM. Fuzy EM. 2006. Effect of soil type on infectivity and persistence of the EPNs *Steinernemascarabaei*, *Steinernemaglaseri*, *Heterorhabditiszealandica*, and *Heterorhabditisbacteriophora*. *J. Invertebr. Pathol.* 92: 11–22.
- Koppenhöfer, A.M., Kaya, H.K, Taormino SP., 1995. Infectivity of entomopathogenic nematodes (Rhabditida: Steinernematidae) at different soil depths and moistures. *Journal of Invertebrate Pathology*, 65: 193-199.
- Koppenhofer. A.M., J.F. Campbell, H. K. Kaya and R. Gaugler. 1998. Estimation of entomopathogenic nematode population density in soil by correlation between bait insect mortality and nematode penetration. *Fundamentals of applied Nematology*, 21(1): 95-102
- Kung, S.P., Gaugler, R., Kaya, HK. 1991. Effects of soil temperature, moisture and relative humidity on entomopathogenic nematode persistence. *Journal of Invertebrate Pathology*, 57(2):242-249
- Kung, S.P., R. Gaugler and H.K. Kaya. 1990. Influence of soil pH and oxygen on persistence of *Steinernema* spp. *Journal of Nematology*, 22: 440-445.
- Mason, J. M., & Hominick, W. M. (1995). The effect of temperature on infection, development and reproduction of heterorhabditids. *Journal of Helminthology*, 69: 337-345.
- Milstead, J.E. (1981). Influence of temperature and dosage on mortality of seventh instar larvae of *Galleria mellonella* and its bacteria associate *Xenorhabdus luminescens*. *Nematologica*., 27 : 167-170 Molyneux and Bedding 1984
- Paoli F., Marianelli L., Binazzi F., Mazza G., Benvenuti C., Peverieri GS., Roversi PF. 2017. Effectiveness of different doses of *Heterorhabditis bacteriophora* against *Popillia japonica* 3<sup>rd</sup> instars: Laboratory evaluation and field application. *Redia-Giornale di Zool.* 100:135-138.
- Platt T., Stokwe NF., Malan AP. 2018. Potential of local entomopathogenic nematodes for control of the vine mealybug, *Planococcus ficus*. *South African J Enol. Viticult.* 39(2):1-8.

- Radova, S. and Trnkova, Z. (2010). Effect of soil temperature and moisture on the pathogenicity of two species of Entomopathogenic nematodes (Rhabditida :Steinernematidae). J. Agrobiol., 27: 1-7.
- Saunders JE., Webster JM.1999. Temperature effects on *Heterorhabditismegidis* and *Steinernemacarpocapsae* infectivity to *Galleria mellonella*. J Nematol.31(3):299-304.
- Shapiro Ilan D., Lewis EE. 2018. Variable biotic and abiotic stress in entomopathogenic nematodes: Implications for biocontrol. Journal of Nematology
- Shapiro IlanD.,Glazer, I., Segal, D. 1997. Genetic improvement of heat tolerance in *Heterorhabditisbacteriophora* through hybridization. Biological Control, 8(2): 153-159
- Shapiro Ilan DI., Brown I., Lewis EE. 2014.Freezing and Desiccation Tolerance in Entomopathogenic Nematodes: Diversity and Correlation of Traits. Journal of Nematology. 46:27-34.
- Shapiro Ilan DI., Cottrell TE., Brown I., Gardner WA., Hubbard RK., Wood BW. 2007. Soil moisture and strain effects on entomopathogenic nematode suppression of the pecan weevil. Pecan Grower 19(1):32-38
- Shapiro-Ilan D, Hazir S ., Lete L. 2015. Viability and virulence of entomopathogenic nematodes exposed to ultraviolet radiation. Journal of Nematology 47: 184-189
- Shelmith, MW., Nderitu JH., Olubayo F., Kimenju JW.2007. Effects of soil texture on virulence and survival of *Steinernemakarii* and *Heterorhabditisindica*. Review of Nematology, 12:754-758.
- Stuart RJ, Barbercheck ME, Grewal PS. 2015.Entomopathogenic nematodes in the soil environment: Distributions, interactions and the influence of biotic and abiotic factors.In: Campos-Herrera, R. (eds) Nematode Pathogenesis of Insects and Other Pests. Sustainability in Plant and Crop Protection. Springer, Cham. [https://doi.org/10.1007/978-3-319-18266-7\\_4](https://doi.org/10.1007/978-3-319-18266-7_4)
- TailliezP.,Pages S., Ginibre N., Boemare N.2006. New insight into diversity in the genus *Xenorhabdus*, including the description of ten novel species. Internat J System EvolMicrobiol. 56:2805-2818.
- Toledo J., Williams T., Perez C., Liedo P., Valle JF., Ibarra JE.2009.Abiotic factors affecting the infectivity of *Steinernemacarpocapsae* (Rhabditida: Steinernematidae) on larvae of

*Anastrepha obliqua* (Diptera: Tephritidae). *Biocontrol Science and Technology*. 19:887-898.

De Waal JY., Addison MF., Malan AP. 2018. Potential of *Heterorhabditis zealandica* (Rhabditida: Heterorhabditidae) for the control of codling moth, *Cydia pomonella* (Lepidoptera: Tortricidae) in semi-field trials under South African conditions. *Int. J. Pest Manag.* 64: 102-109.

Wharton DA., Goodall G., Marshall CJ. 2003. Freezing survival and cryoprotective dehydration as cold tolerant mechanisms in the Antarctic nematode *Panagrolaimus davidi*. *Journal of Experimental Biology*. 206: 215-221.

Wharton, DA., Surrey, M. R. (1994). Cold tolerance mechanisms of the infective larvae of the insect parasitic nematode, *Heterorhabditis zealandica* Poinar. *Cryo Letters*, 15: 353-360.

Wright PJ. 1992. Cool temperature reproduction of steinernematid and heterorhabditid nematodes. *J. Invertebr. Pathol.* 60(2): 148-151.

Yul C, Wonn L, Sook Y, Myeong L, Thi H. 2002. Effects of temperature and nematode concentration on pathogenicity and reproduction of entomopathogenic nematode, *Steinernema carpocapsae* Pocheon strain (Nematode: Steinernematidae). *Korean J Appl Entomol.* 41: 269-277.

**Comment [DCKP14]:** Make sure every references you used here is present in the manuscript