

## Review Article

### **Plant Growth Promoting Rhizobacteria for Sustainable Production of Sugarcane and Rice**

#### **Abstract**

Sugarcane (*Saccharum officinarum* L.) and rice (*Oryza sativa* L.) are important cash and staple crop in the world respectively. In today's population explosion the lots of pressure on cropping land to mitigate the feed need of consumers with judicious use of chemical fertilizers without considering the health of soil and its sustainability. The rhizobacteria live region around roots (2-80 mm) of crop with diverse, dynamic and complex microflora having capacity with direct and indirect beneficial effects on crop health by availability of different nutrient, siderophores, rhizodeposits, auxins and role with bioremediation. There are different types of bacteria isolated from rhizosphere of sugarcane and rice having *Bacillus*, *Rhizobium*, *comamonas*, *cyanaobacteria*, *Nodosilinea*, *Levinella* and *Pseudomonas* genera with most efficient nitrogen producer, solubilizers of inorganic phosphate, potash and other macro and micronutrient. In rice microbiome as *Levinella*, *Pseudomonas*, *Anaeromyxobacter*, *Arenimonas*, *Arthrobacter*, *Bacillus*, *Bellilinea*, *Proteobacteria*, *Chloroflexi*, *Actinobacteria*, and *Acidobacteria*, and *cyanobacteria* with strains of *Nitrosomonas* and *Nitrobacter* are further divided in vegetative and reproductive growth stage of crop. The article give brief idea about the large arena of them.

**Key words**-Rhizobacteria, sugarcane, rice, sustainable production

#### **Introduction**

Rhizosphere give the term by Hiltner (1904) as narrow soil region near root zone having different secretion and associated soil microorganism as root microbiome communication. They feed on sloughed off plant cells called rhizodeposition and vice versa root released protein called root exudates. This symbiosis influence plant growth providing space to produce some *allochemicals* to control neighbors. Having the *rhizospheric* effect enhance the redox potential, ionic balance, osmotic pressure with formation of soil aggregates and prevent the crop roots

from desiccation in abiotic stress. Sugarcane (*Saccharum officinarum* L.) and rice (*Oryza sativa* L.) are important cash and staple crop in the world respectively. Sugarcane gives us tea sugar, brown sugar, alcohol, spirit, bagasse, press mud cakes, molasses with renewable energy sources for community transport with methanol blending in petrol. Every sugar factory enhances the socio economic status of the producing farmer in their locality with direct indirect employments opportunity generated by them. Now a days bio electricity generated by using the sugarcane wastes. For its sustainable production with soil management required use of different biofertilizers with bacterial inoculations consortia. In sugarcane various kinds of bacterial endophytes such as *Glucano acetobacter*, *Psudomonas*, *Azospirillum*, *Burkholderia*, *Raoultella*, *Ralstonia Mesorhizobium*, *Ochrobactrum*, *Sphingomonas*, *Novosphingobium*, *Pantoea* and *Bacillus* were found and they are perform various role like promote growth, including phosphate solubilization, iron carrier formation, saponin and glycosidase production, biological nitrogen fixation, and the production of 1-amino-cyclopropane-1-carboxylic acid deaminase and development throughout its life period.

Rice the main staple food for about half of the world's population, microorganisms exert positive influences on plant growth, production, and health of rice. A variety of microbial species as actinomycetes and Gram-positive bacteria reside in the rhizosphere and the phyllosphere of plants and also have multiple roles as symbiotic endophytes. They alter the morphology of host plants, enhance their growth, health, and yield, and reduce their vulnerability to biotic and abiotic stresses. Bacterial communities are usually dominated by the *Anaeromyxobacter*, *Arenimonas*, *Arthrobacter*, *Bacillus*, *Bellilinea* *Proteobacteria*, *Chloroflexi*, *Actinobacteria*, and *Acidobacteria*, and *cyanobacteria* which help in uptake of nitrogen and other growth promoting role in rice.

## 1. Sugarcane

### a) *Azospirillum* and sugarcane

*Azospirillum* is facultative endophytic, polymorphic, obligate can grow under low  $PO_2$  soil condition. Are micro-aerophilic, gram-negative rods and often associated with interior and surfaces roots of cereals and grasses (Grifoniet *al.*, 1995). It is very well known for its nitrogen-fixing ability and can fix at least 10 mg N g<sup>-1</sup> of carbohydrate and higher production of indole acetic acid. Three species of this genus namely, *Azospirillum amazonense*, *A. brasilense*, and *A.*

*lipoferum* have been isolated from sugarcane roots. Microbial consortia containing the species, *Azospirillumbrasilense*, and *Bacillus subtilis* have shown great potential to cycle nutrients from crop residues and restore soil fertility. The seed treatment enhances the root volume at 86% and 100% increase than untreated ones. The inoculation with *Azospirillumbrasilense* + *B. subtilis* associated with 45 kg ha<sup>-1</sup> of P<sub>2</sub>O<sub>5</sub> also increases sugar yield, resulting in a savings of 75% of the recommended P<sub>2</sub>O<sub>5</sub> rate (Rosa *et al.*, 2022).

#### *Enterobacteriaceae* family bacteria and sugarcane

The members of *Enterobacteriaceae* large family of bacteria are rod-shaped, gram-negative, facultative anaerobes, motile or non-motile and most of them reduce nitrate to nitrite. The dominating nitrogen-fixing bacteria isolated from sugarcane, rice, wheat, sorghum, grasses, and dicotyledonous plants roots. The three genera belong to the *Enterobacteriaceae* as *Enterobacter cloacae* and *Klebsiella pneumoniae* and *Pantoea* were isolated from all parts of the sugarcane. Three species of this genus, *Pantoeaananatis*, *P. herbicola*, and *P. stewartii* have been isolated from roots, stem, and leaves of Brazilian sugarcane plants (Mendes *et al.* 2007) capable of nitrogen fixation.

#### *Gluconacetobacter* and sugarcane

This genus belongs to the family *Acetobacteraceae* to produce acetic acid and nitrogen fixing and usually acid-tolerant and grow well below pH 5.0. They are gram-negative, aerobic, and rod-shaped bacteria. This organism has lack of nitrate reductase and only partial inhibition of nitrogenase activity by ammonium ions, enables it to fix nitrogen in the presence of soil nitrogenease is also a phytohormone producer. In sugarcane endophytic *Gluconacetobacterdiazotrophicus* and its abilities to fix nitrogen and disease resistance studied (Boddey, *et al.*, 2003). *G. dizotrophicus* inhibit *Xanthomonas albilineans* (leaf scald disease) (Blanco *et al.*, 2005). At Central Sugarcane Research Station Padegaon, Dist. Satara (Maharashtra) studied the effect of *Glucanoacetobacterdiazotrophicus* @ 10kg/ha (Set treatment) with other six treatment was carried from year 2014-2018 on pre-season sugarcane on COM 0265 variety with (75% P<sub>2</sub>O<sub>5</sub>, 100% K<sub>2</sub>O + Recommended FYM (Suru- 20 t/ha, Pre-season- 25 t/ha) and PSB application common to all treatments except T1 RDF, found that treatment with 50% N+ *Glucanoacetobacter* set treatment recorded highest tillering count at 120 DAP (167.05 thousands/ha) and no. of mill liable cane. For cane yield same treatment

recorded highest cane yield (57.54 t/ha), CCS%, CCS yield and sucrose %.(Khandagale *et al.*,2018).Same trend found in suru(Dec-Jan.Plantation) sugarcane as treatments with ( 50% N+ *Glucanoacetobacter* set treatment )recorded highest (162.00) tillering count at 120 DAP,NMC (53.31 thousand /ha) and cane yield. (52.17 t/ha. As regards CCS yield highest (7.98 t/ha), highest purity (97.90),sucrose %(21.20) found with same trends.The endophytic bacterial count found in same treatment as in root (7.10x10<sup>4</sup>, 11.20x10<sup>4</sup> and 7.58x10<sup>4</sup>), juice (16.53x10<sup>4</sup>,29.63x10<sup>4</sup> and 18.96x10<sup>4</sup>) and leaves (14.48x10<sup>4</sup>, 30.01x10<sup>4</sup> and 16.35x10<sup>4</sup>) at 4 months , 8 months and harvest stage of the crop, respectively.Having beneficial impact on sugar recovery , NMC and other yield attributing parameters.Some scientist found that a new role for *G. diazotrophicus* induce systemic resistance against *Xanthomonas albilineans*-cause leaf scald disease of sugarcane(Arencibia *et al.* 2006).

### ***Pseudomonas* spp.and sugarcane**

The genus belongs to the family *Pseudomonadaceae* with gram negative ,rod shaped bacteria secreted soluble greenish fluorescent pigment called *fluorescein* in low availability of iron are facultative anaerobic require NO<sub>3</sub> without O<sub>2</sub> and some with O<sub>2</sub> with multiple polar flagella as they are *chemolithotrophs*. Having large number of species with very well-known PGPR bacteria due to its ability to produce phytohormones, siderophores, antibiotics, phosphate solubilization and production of antifungal compounds. Some species also fix nitrogen in addition to above-mentioned characteristics. *Pseudomonas fluorescens*, *Pseudomonas putida* for PGPR and *Pseudomonas aurantiaca* , *Pseudomonas chlororaphis* for biocontrol agents in foliar spray due to the production of antifungal phenazine compounds. *Pseudomonas* spp. have been isolated from stem, root, leaves, and rhizosphere of sugarcane growing in Australia, Brazil, India, Pakistan, and South Africa.They are among most efficient solubilizers of inorganic phosphate ( Tayade *et al.*,2019).They produce phytohormones like indoleacetic acid, gibberellins, and cytokinin, iron-sequestering siderophores, phosphate-solubilizing enzymes, and 1-aminocyclopropane-1-carboxylate (ACC) deaminase (Yanni *et al.*,2001).Growth hormones produced by the bacteria enhance the development of lateral roots and improve the plant's nutrient uptake from the rhizosphere.

### **PGPR and biocontrol agents in sugarcane**

PGPR induce resistance in plants against fungal, bacterial and viral diseases, insects and nematodes. PGPR bring about Induced resistance through fortifying the physical and mechanical strength of the cell wall as well as changing the physiological and biochemical reaction of the host leading to the synthesis of defense chemicals against the challenge pathogen. PGPR provide different mechanisms for suppressing plant pathogens. They include competition for nutrients and space, antibiosis by producing antibiotics viz., pyrrolnitrin, pyocyanin, 2,4-diacetyl phloroglucinol and production of siderophores, viz., *Pseudobactin* which limits the availability of iron necessary for the growth of pathogens.

The *P. fluorescens* strains from sugarcane reported antifungal activity against *Fusarium oxysporum* and *Rhizoctonia bataticola* (Kumar *et al.* 2002). The strains *P. putida*, *B. subtilis*, and *Stenotrophomonas maltophilia* found control against local strains of red rot -*Colletotrichum falcatum* (Hassan *et al.* 2010 and Viswanathan and Samiyappan 2002). Antwerpen *et al.* (2002) checked the antifungal activity of *Burkholderia isolates* from the sugarcane rhizosphere, against *U. scitaminea* (sugarcane smut) and *Fusarium spp.* (stalk rot) (Hassan *et al.*, 2010)

## **Rice**

In rice vegetative stage bacterial population are differ from their reproductive stage. The ponding standing water in rice field rich in of Gram-negative bacteria plus algae. while in percolating water having only Gram-negative bacteriadominated by the *Anaeromyxobacter*, *Arenimonas*, *Arthrobacter*, *Bacillus*, *Bellilinea*, *Proteobacteria*, *Chloroflexi*, *Actinobacteria*, and *Acidobacteria*, and *cyanobacteria*.

## **Rice and *Pantoea* inoculant**

Bacteria belonging to the genus *Pantoea* are gram negative, highly diverse, aerobic or facultative anaerobic with peritrichous flagella in *Enterobacteriaceae* family though they are infecting rice with diseases some strains are till beneficial (*P. ananatis*, *P. agglomerans*, *P. rodosii*) showing a promising capacity to mediate the negative effects of arsenic stress as bioremediation and also act as phyto stimulants. They act as potential P solubilizers, N fixer, IAA and salt tolerance with siderophore production capacity. The rice endophyte *Pantoea agglomerans* (YS19) showed nitrogen-fixing activity *in vitro* produced four different phytohormones, including IAA and

promoted plant growth(Fenget *al.*,2006) They found ubiquitous in roots and reduced at the time of tillering with subsequent stages(Chebotaret *al.*,2015)

### **Rice and *Bacillus***

Microorganisms in the endospheric *Bacillus* are aerobic, gram positive, soil bacterium affect the root morphology, tiller biomass when was applied as an inoculant biofertilizer. They synergists affects on chlorophyll content, plant height, number of tillers, tiller biomass and yield attributes in rice. The combination of TUAT-1(Khim et al.,2018 *Bacillus pumilus* strain ) and 100% N (farmer recommended rate of N) resulted in the greatest tiller number and biomass at the maximum tillering stage, and positively affected other growth attributes and yield(4.89 ton ha<sup>-1</sup> ) and concluded that TUAT-1 promoted root development which increased nutrient uptake from the soil by promoting the growth and development of roots.

The strain *Bacillus subtilis sub sp. megatherium* acts as the PGPR and bio control agent in rice so that plant can overcome the negative effect of environmental stress .Plant increased micronutrients uptake affect Phyto hormone homeostasis, stabilized membrane integrity, decreased leaf transpiration ,increased nutrition and metabolic activity with maintaining cell turgor of rice leaves under drought stress because of production of hormone IAA(Abdelaziz *et al.*,(2018).

### **Rice and Siderophore-producing bacteria**

The genera *Sphingomonas*, *Pseudomonas*, *Pantoea* , *Burkholderia* and *Enterobacter* are primarily detected in rice plant tissues during plants' vegetative stage acts as siderophores, giving them greater access to iron and other elements in the soil. The *Pseudomonas fluorescens* at field application enhanced the number of tillers ,plant height and biomass with yield increase at 20-26 % over control( Seenivasan, 2011). Inoculation with *Beijerinckia indica* in field increased no. of tillers and plant height with 25-29% increased yield.( Biswas *et al.*,2000).

### **Rice and *Azospirillum* inoculant**

They are surface colonizing, gram negative, free living, nitrogen fixing, polar flagellar moment forming cyst in unfavorable climatic condition. The inoculation of rice seedlings with *Azospirillum brasilense* promoted early tillering and the better reproductive performance of rice plants. It was found to significantly increase the grain-filling rate and the grain weight per plant at harvest time. Isawa *et al.*, (2010) found that *Azospirillum* sp. (B510) strain in field condition enhanced no. of tillers and plant height with increased yield at 17-25%.

### **Rice and *Pseudomonas* inoculation**

In rice the inoculation with *P. dispersa* significantly improves the morpho-biochemical characteristics of the plants such as increased enzymatic activities and reduced arsenic uptake into rice tissues (Ghosh *et al.* (2021). A study by Patel *et al.* (2021) and (Oilvera *et al.*, 2019) indicates that *Pseudomonas fluorescens* produces a volatile organic compound (pyrazine) that can suppress infection and physiological damage can be reduced given by *Magnaporthe oryzae* in rice.

### **Rice and *Rhizobium* bacterial inoculant**

Though rice plants are not leguminous. So benefits of nitrogen not from N-fixation in nodules formed on plant roots because they can't have the *Sym* gene for endophytic colonization of rice root by them. But they having beneficial endophytic association (Chan and Zhu, 2013, Alam *et al.*, 2001). They enhance crop health and yield and reduced the harmful effect of fungicides, pesticides and synthetic chemicals. In Egypt, it was found that the association of *Rhizobium leguminosarum* with rice plants could significantly increase the shoot and root growth of these plants, their grain yield, and their nitrogen-use efficiency (Yanni *et al.*, 1997). The growth and yield responses of rice plants to inoculation with two bacterial strains (*R. leguminosarum* (E11) and *Rhizobium* sp. (IRBG74) showed significantly increased grain and higher straw yields at the maturity stage (Biswas *et al.*, 2009). A multinational collaborative study with three field experiments in Egypt reported that the *Rhizobium* inoculation of rice plants significantly increased their biomass, nutrient uptake (due to more robust root architecture), grain yield, fertilizer efficiency, harvest index, and grain nutritional value. Further experiments of selected *rhizobiaum* endophytes on rice showed that they produced cell-bound cellulase and polygalacturonase enzymes that hydrolyze glycosidic bonds in plant cell walls plus certain bacteriocins that can inhibit the growth of undesirable microbes.

## Rice and PGPR as disease control

Antifungal compounds produced by *B. subtilis* show a strong synergistic inhibitory effect on the hyphal growth of *Pyricularia grisea* and *R. solani* the respective causative agents for blast and sheath blight in rice (Leelasuphakulet *et al.*,2006).Another study that employed a strain of endophytic bacterium *Bacillus* sp. (EBPBS4) and *Serratia marcescens* used for managing sheath blight as antagonistic toward the fungal pathogen *R. solani* by decreasing its severity and incidences (Durgadevi *et al.*,2021,Yang, 2009,Somaya *et al.*,2005).*Bacillus oryzicola* reduced the bakanae disease (*Gibberella fujikuroi*) severity by 46-78% in field condition after its inoculation.The *Pseudomonas* spp. as biocontrol agents for resisting various soil-borne plant diseases (Stockwell and stock 2007) inhibits the mycelial growth of the sheath blight fungus *R. solani* by increasing chitinase and peroxidase activity in rice act as inducing systemic resistance.(Nandkumare *et al.*,2001).A talc-based bioformulation reduce the incidence of sheath blight in rice by up to 62%, with 12–21% more grain yield because of trigger the ISR against the microbes (Radiacommare *et al.*,2002,Vidyasakaran *et al.*,1997).

## Conclusion –

From above all various information we concluded that in sugarcane and rice PGPR promote plant growth directly and indirectly by increasing uptake and availability of macronutrients (N, P, K and essential minerals) and play a role by regulating plant hormone levels indirectly by reducing the inhibitory effects of various pathogens on plant growth and development in the form of biological control agents (Tariq *et al.*,2017). For future need of mankind with hygienic and residue free organic harvest we require careful management without burning the farm residue ,mulching ,heap management for harboring them in field. They are also act as the biocontrol agent against many disease in both the crop. By active application of microbial consortia at different growth stages of crop eliminate the dependency of big imports of potassic and phosphatic fertilizers with saving the incredible foreign currency .The harmonize

and symbiotic management of them must be emphasized for *satsvikaslakshy in swarnim Bharat* with biosphere and ecological conservation having sustainability for our mankind .

## References

- Abdelaziz S., Hemeda N.F., Belal E.E. and Elanberg R. (2018) Efficacy of facultative oligotrophic bacterial strain as PGPR and their potency against two pathogenic fungi causing damping off disease. *Applied microbial* .Open access.
- Alam M.S., Cui Z.J., Yamagishi T., Ishii R. (2001) Grain yield and related physiological characteristics of rice plants (*Oryza sativa* L.) inoculated with free-living rhizobacteria. *Plant Prod. Sci.* **4**:126–130.
- Antwerpen T.V., Rutherford R.S and Vogel J.L (2002) Assessment of sugarcane endophytic bacteria and *rhizospheric Burkholderia* species as antifungal agents. *Proc. S. Afr. Sug. Technol. Assoc.* **76**:301–304.
- Arencibia A.D., Vinagre F., Estevez Y., Bernal A., Perez J., Cavalcanti J., Santana I and Hemerly AS. (2006) *Gluconacetobacter diazotrophicus* elicits a sugarcane defense response against a pathogenic bacteria *Xanthomonas albilineans*. *Plant Signal Behav.* **1**(5):265–273
- Biswas J.C., Ladha J.K., Dazzo F.B., Yanni Y.G and Rolfe B.G. (2000) Rhizobial inoculation influences seedling vigor and yield of rice. *J. Agron.* **92**:880–886.
- Biswas J.C., Ladha J.K., Dazzo F.B. (2009) Rhizobia inoculation improves nutrient uptake and growth of lowland rice. *Soil Sci. Soc. Am. J.* **64**:1644–1650.
- Blanco Y., D. Pinon., Legaz M.E. and C. Vicente. (2005). Antagonism of *Gluconacetobacter diazotrophicus* (a sugarcane endosymbiont) against *Xanthomonas albilineans* (pathogen) studied in alginate-immobilized sugarcane stalk tissues. *J. Biosci. Bioeng.* **99**:366-371.
- Boddey R. M., Urquiaga, S., Alves, B.J.R., and V. Reis. (2003). Endophytic nitrogen fixation in sugarcane: present knowledge and future applications. *Plant Soil* **252**:139-149.
- Chebotar V.K., Malfanova N.V., Shcherbakov A.V., Ahtemova G.A., Borisov A.Y., Lugtenberg B., Tikhonovich I.A (2015) Endophytic bacteria in microbial preparations that improve plant development (review). *Appl. Biochem. Microbiol.* **51**:271–277.

- Chan C. and Zhu H.(2013) Are common symbiosis gene required for endophytic rice rhizobial interaction. *Plant signaling behavior*.8(9)25453.
- Durgadevi D., Harish S., Manikandan R., Prabhukarthikeyan S.R., Alice D andRaguchander T. (2021)Proteomic profiling of defense/resistant genes induced during the tripartite interaction of *Oryza sativa*, *Rhizoctonia solani* AG1-1A, and *Bacillus subtilis* against rice sheath blight. *Physiol. Mol. Plant Pathol.* **115**:101-669.
- Feng Y., D. Shen D. and W. Song.( 2006.) Rice endophyte *Pantoeaagglomerans* YS 19 promotes host plant growth and affects allocations of host photosynthates. *J. Appl. Microbiol.***100**:938-945.
- Grifoni A., BazzicalupoM.,Serio C.D., FarcelliS. and Fani R.(1995)Identification of *Azospirillum* strains by restriction fragment length polymorphism of 16 S rDNA and the histidine operon. *FEMS Microbiol Lett.***127**:85–91.
- Ghosh A., Pramanik K., Bhattacharya S., Mondal S., Ghosh S.K., Ghosh P.K and Maiti T.K.(2021)Abatement of arsenic-induced phytotoxic effects in rice seedlings by an arsenic- resistant *Pantoeadispersa* strain. *Environ. Sci. Pollut. Res.* **28**:21633–21649.
- Hassan M.N., Afghan S and Hafeez F.Y (2011) Biological control of red rot in sugarcane by native pyobutine producing *P.putida* strain NH 50 under field condition and its potential mode of action *Pest management Sci.*67(9):1147-1154..
- Isawa T., Yasuda M., Awazaki H., Minamisawa K., Shinozaki S.andNakashita H.(2010)*Azospirillum* sp. strain B510 enhances rice growth and yield. *Microbes Environ.* **25**:58–61.
- Kumar N.R., Arasu V.T.and Gunasekaran P.(2002) Genotyping of antifungal compounds producing plant growth-promoting rhizobacteria, *Pseudomonas fluorescens*. *Curr Sci* 82(12):1463–1466.
- Khandagale P.P.,Kesker M.M.,GhodkeS.K.,Rasker B.S. (2019) Study the effect of the *Gluconoacetobacterdiazotrophicus* and PSB formulation on quality parameter on suru sugarcane *.Int.J.Current Microbiology Applied sci.*8(5):210-215.

- Khin T.W., Odd A.Z., Ohstu N.O. and Yokohoma T. (2018) *Bacillus pumilis* strain TUAT - and nitrogen application in nursery phase promote growth of rice plants under fields condition. *Agronomy*. **8**(10):216-222.
- Mendes R., Pizzirani-Kleiner A.A., Araujo W.L and Raaijmakers J.M. (2007) Diversity of cultivated endophytic bacteria from sugarcane: genetic and biochemical characterization of *Burkholderiacepacia* complex isolates. *Appl Environ Microbiol* **73**:7259–7267.
- Nandakumar R., Babu S., Viswanathan R., Raguchander T. and Samiyappan R. (2001) Induction of systemic resistance in rice against sheath blight disease by *Pseudomonas fluorescens*. *Soil Biol. Biochem.* **33**:603–612.
- Oliveira M.I.D.S., Chaibub A.A., Sousa T.P., Cortes M.V., D.Souza A.C., Conceicao E.C.D and D.Filippi M.C. (2019) Formulations of *Pseudomonas fluorescens* and *Burkholderia pyrrocinia* control rice blast of upland rice cultivated under no-tillage system. *Biol. Control*. **27**:104-153.
- Pedraza R.O. (2008) Recent advances in nitrogen-fixing acetic acid bacteria. *Int. J. Food Microbiol.* **125**(1):25–35.
- Patel A., Kumar A., Sheoran N., Kumar M and Sahu K.P. (2021) Antifungal and defense elicitor activities of pyrazines identified in endophytic *Pseudomonas putida* BP25 against fungal blast incited by *Magnaporthe oryzae* in rice. *J. Plant Dis. Prot.* **128**:261–272.
- Radjacomare R., Nandakumar R., Kandan A., Suresh S., Bharathi M., Raguchander T and Samiyappan R. (2002) *Pseudomonas fluorescens* based bioformulation for the management of sheath blight and leaf folder in rice. *Crop Prot.* **21**:671–677.
- Rosa P.A.L., Galindo F.S., Olivera C.E.D., Jalal A., Martinao E.S., Fernadas G.C. and Morgana E.R. (2022) Inoculation with plant growth-promoting bacteria to reduce phosphate fertilization requirement and enhance technological quality and yield of sugarcane. *Microorganisms* **10**(1): 192-200.
- Stockwell V.O and Stack J.P. (2007) Using *Pseudomonas* spp. for integrated biological control. *Phytopathology*. **97**:244–249.

- Someya N., Nakajima M., Watanabe K., Hibi T., Akutsu K. (2005) Potential of *Serratia marcescens* strain B2 for biological control of rice sheath blight. *Biocontrol Sci. Technol.* **15**:105–109.
- Tayade A.S., Geetha, P., Anusha, S., Dhanapal, R. and Hari, K. (2019) Bio-intensive modulation of sugarcane ratoon rhizosphere for enhanced soil health and sugarcane productivity under tropical Indian condition. *Sugar Tech* **21**: 278–288.
- Seenivasan N. (2011) Efficacy of *Pseudomonas fluorescens* and *Paecilomyces lilacinus* against *Meloidogyne graminicola* infesting rice under system of rice intensification. *Arch. Phytopathol. Plant Prot.* **44**:1467–1482.
- Tariq M., Noman M., Ahmed T., Hameed A., Manzoor, N., and Zafar, M. (2017). Antagonistic features displayed by plant growth promoting rhizobacteria (PGPR): a review. *J. Plant Sci. Phytopathol.* **1**: 38–43.
- Vidhyasekaran P., Rabindran R., Muthamilan R., Nayar M., Rajappan K., Subramanian K., Vasumathi K. (1997) Development of powder formulation of *Pseudomonas fluorescens* for control of rice blast. *Plant Pathol.* **46**:291–297.
- Viswanathana R. and Samiyappan R. (2002) Induced systemic resistance by *Pseudomonas fluorescens* against red rot disease of sugarcane caused by *Colletotrichum falcatum*. *Crop Prot.* **21**:1–10
- Yanni Y.G., Rizk R.Y., El-Fattah F.K.A., Squartini A., Corich V. G.A., De Bruijn F., Rademaker J, Maya Flores J., Ostrom P., Vega-Hernandez M, Hollingsworth RI, Martinez-Molina E, Mateos P, Velazquez E, Wopereis J, Triplett E., Umali-Garcia M., Anarna J.A., Rolfe BG., Ladha J.K., Hill J., Mujoo R., Ng P.K. and Dazzo FB (2001) The beneficial plant growth-promoting association of *Rhizobium leguminosarum* bv. *trifolii* with rice roots. *Aust. J. Pl. Physiol.* **28**:845–870
- Yanni Y.G., Rizk R.Y., Corich V., Squartini A., Ninke K. and Philip-Hollingsworth S. (1997) Natural endophytic association between *Rhizobium leguminosarum* bv. *trifolii* and rice roots and assessment of its potential to promote rice growth. *Plant Soil.* **194**:99–114.

Yang D., Wang B., Wang J., Chen Y., Zhou M. (2009) Activity and efficacy of *Bacillus subtilis* strain NJ-18 against rice sheath blight and *Sclerotinia* stem rot of rape. *Biol. Control.* **51**:61–65.

UNDER PEER REVIEW