

Effect of continuous application of Nitrogen, Phosphorus and Potassium on growth parameters and yield of Rice

Abstract

A Long term (37 years) field experiment was conducted at Agronomical Research farm of Birsa Agricultural University, Kanke, Ranchi during Kharif season in 2020 to study the effect of Nitrogen, phosphorus and potassium on growth parameter and yield of rice. The experiment was conducted in Partially Confounded Design with nineteen treatments replicated four times. The rice variety used was Sahabhagi Dhan. Nitrogen, Phosphorus and potassium level used were 40, 80 and 120 kg N ha⁻¹, 0, 40 and 80 kg P₂O₅ ha⁻¹ and 0 and 40 kg K₂O ha⁻¹. Application of 120 kg N ha⁻¹, 80 kg P kg P₂O₅ ha⁻¹ and 40 kg K₂O ha⁻¹ significantly increased plant height, produced maximum numbers of tillers, dry matter and leaf area index at 30, 60, 90 DAS and at harvest. The maximum grain and straw yield was recorded under with application of 120 kg N ha⁻¹, 80 kg P kg P₂O₅ ha⁻¹ and 40 kg K₂O ha⁻¹.

Key word: growth parameters, yield, Kharif season, Soil fertility, crop growth, nitrogen level, Rice consumption

Introduction

“The success of Indian agriculture has received worldwide appreciation as food grain production increased from 50.8 million tons (mt) in 1950-51 to 182.57 mt in 2002-03, and this total food grain production comprised of 66.51 mt of rice, 69.32 mt of wheat, 26.22 mt of coarse cereals, and 11.31 mt of pulses” [1]. In India, rice plays a major role in diet, economy, employment, culture and history. It is the staple food for more than 65% of Indian population contributing approximately 40% to the total food grain production, thereby, occupying a pivotal role in the food and livelihood security of people. The country has the world's largest area under rice i.e., about 43 million hectare (Mha) and the second highest production i.e., about 110 Mt of milled rice at productivity of 2.56 t ha⁻¹ as per 2016-17 statistics. Global demand of rice needs to increase from the current 493 Mt to about 550 Mt in 2030. However, rice farming, particularly in the rainfed regions, faces multiple risks from uncertain climate, degraded soil, water shortage and underdeveloped markets. Introduction of high yielding varieties, irrigation and high analysis fertilizer accelerated the mining of nutrient other than supplied eternally from soil. To sustain the productivity it was essential to maintain the supply of nutrient. Since large amount of nutrient has to be applied to soil in chemical form which may have impact on soil properties and soil productivity in long term. The concept of balanced fertilization cannot be confined to N, P and K alone. Balanced fertilization includes application of all the plant nutrients essential for high agricultural productivity and health of the soil. Therefore, a paradigm shift is required for enhancing the rice productivity and sustainability. Hence the study focuses on the effect of continuous application of fertilizer on growth parameter and yield of rice.

Materials and method

A long term (37 year) field experiment was conducted at Agronomical Research farm of Birsa Agricultural University, Kanke, Ranchi during Kharif season in 2020. The experimental plot was a medium land having well drained soil and uniform topography. The soil was sandy loam in texture with bulk density 1.54 Mg m^{-3} and good water retention (FC 21.5 % and PWP 11.36 %) and water holding capacity (38.7%) was also observed in soil. Soil reaction was acidic in nature with low available N and P, medium in available K. The experiment was conducted in Partially Confounded Design with nineteen treatments replicated four times. The rice variety used was Sahabhazi Dhan. Nitrogen, Phosphorus and Potassium level used were 40, 80 and 120 kg N ha⁻¹, 0, 40 and 80 kg P₂O₅ ha⁻¹ and 0 and 40 kg K₂O ha⁻¹ respectively. The inorganic source of Nitrogen (N), Phosphorus (P) and Potassium (K) were applied through urea, Diammonium phosphate and muriate of potash as per treatment. The full dose of P and K along with one third of N were applied at the time of sowing and remaining N in 2 equal splits top dressed as per treatment at 30 and 60 DAS. The data on growth parameter were recorded at 30, 60, 90 DAS and at harvest.

Result and Discussion

Effect on growth parameter

In general optimum supply of N, P and K contributed to the plant growth and development. The study of data (Table 1 and 2) revealed that plant height, number of tillers and dry matter accumulation of rice differed markedly in different fertilizer doses at all growth stages of rice. The data shows consistent increase in plant height, number of tillers and dry matter accumulation with advancement of crop growth stages. It is also evident from the data that more than 50% height of rice and Number of tillers was attained by 60 DAS in all treatments.

The maximum plant height, number of tillers and dry matter accumulation was recorded in crop fertilized with 120 kg of N ha⁻¹ at 30 DAS, 60 DAS, 90 DAS and at harvest. The lowest plant height, number of tillers, leaf area index and dry matter accumulation is recorded in Control (no fertilizer) at all growth stages. The application of 120 kg of N increased plant height by 32.15%, 51.90%, 54.75% and 39.42% at 30 DAS, 60 DAS, 90 DAS and at Harvest respectively over control. Similarly, application of 120 kg N ha⁻¹ increase tillers by 21.5%, 20.92%, 63.34% and at 30 DAS, 60 DAS, 90 DAS and at harvest compared to control. [2] and [3] recorded similar result. N Application of 120 kg N ha⁻¹ produced maximum dry matter over other fertility level and was 89.23%, 115.33%, 125.21% and 138.62% higher over control at 30, 60, 90 and harvest respectively. [4] and [5] recorded similar findings for dry matter accumulation. However fertility level of 120 kg N ha⁻¹ remained at par with fertility level of 80 kg N ha⁻¹ for plant height

P ₀	13.68	62.64	71.00	83.91	59.67	293.00	244.75	219.58
P ₄₀	15.48	76.00	93.55	104.39	69.92	341.17	311.13	257.50
P ₈₀	15.72	81.86	101.95	109.69	71.00	362.50	331.50	271.67
SEm±	0.30	1.48	1.88	1.86	1.41	7.11	6.13	4.50
CD at 5%	0.84	4.19	5.33	5.27	4.00	20.18	17.40	12.77
K Level								
K ₀	14.75	71.63	84.97	96.82	66.08	323.44	285.39	242.22
K ₄₀	15.16	75.37	92.70	101.84	67.64	341.00	306.19	256.94
SEm±	0.24	1.21	1.54	1.52	1.15	5.81	5.01	3.68
CD at 5%	0.69	3.42	4.36	4.30	3.26	16.48	14.21	10.43
Control	11.60	50.73	60.45	74.25	56.58	282.00	186.50	150.42
CV%	10.06	10.30	10.87	9.50	10.56	10.72	10.72	9.37

Table 2: Effect of continuous application of Nitrogen, Phosphorus and Potassium dry matter accumulation, LAI and yield of Rice.

Fertilizer level	Dry matter accumulation (g m ⁻²)				Leaf area index			Grain yield (q ha ⁻¹)	Straw yield (q ha ⁻¹)
	30	60	90	harvest	30	60	90		
N Level									
N ₄₀	45.35	170.51	470.94	622.42	0.60	2.44	2.76	21.75	33.99
N ₈₀	47.02	178.25	495.92	667.01	0.65	2.61	2.92	25.79	39.73
N ₁₂₀	47.97	184.41	519.58	709.02	0.68	2.75	3.12	27.01	41.42
SEm±	0.96	2.98	9.84	13.16	0.01	0.05	0.06	0.44	0.77
CD at 5%	2.73	8.47	27.93	37.35	0.04	0.15	0.16	1.26	2.18
P Level									
P ₀	39.94	145.04	391.03	486.73	0.47	2.05	2.20	16.23	26.55
P ₄₀	48.12	185.20	517.50	707.45	0.71	2.67	3.12	26.83	40.86
P ₈₀	52.29	202.93	577.91	804.27	0.74	3.08	3.48	31.48	47.74
SEm±	0.96	2.98	9.84	13.16	0.01	0.05	0.06	0.44	0.77
CD at 5%	2.73	8.47	27.93	37.35	0.04	0.15	0.16	1.26	2.18
K Level									
K ₀	45.81	172.33	476.66	637.22	0.62	2.51	2.88	23.85	37.18
K ₄₀	47.76	183.11	514.30	695.08	0.66	2.68	2.99	25.84	39.58
SEm±	0.78	2.44	8.04	10.75	0.01	0.04	0.05	0.36	0.63
CD at 5%	2.23	6.91	20.80	30.49	0.03	0.12	NS	1.03	1.78
Control	25.35	85.64	230.70	297.13	0.40	1.82	1.90	13.71	22.90
CV%	10.88	8.78	10.16	10.26	10.94	10.24	9.93	9.34	10.39

Effect on grain and straw yield

The data of grain and straw yield (Table 2) revealed that application of full dose of nutrients nitrogen i.e., 120 kg N ha⁻¹ provided maximum grain and straw yield. “Application of 120 kg N ha⁻¹ increased grain and straw yield over imbalance and control, however remained at par with application of 80 kg N ha⁻¹. The increase in grain yield at higher nitrogen rates might be primarily due to increase in chlorophyll concentration in leaves leading to higher photosynthetic rate and might be due to the cumulative effect of increased translocation of photosynthates to sink resulting in the increase in yield compared to lower levels” [9, 10]. [11] recorded similar findings. [15] reported that “increased nitrogen levels significantly increased rice grain and straw yield”.

“Application Phosphorous at the rate 80 kg ha⁻¹ produced maximum grain and straw yield i.e., 31.48 q ha⁻¹ and 47.74 q ha⁻¹ respectively. Phosphorous level of 80 kg ha⁻¹ increase grain yield by 93.96% and straw yield by 79.81 % over application of 0 kg P₂O₅ ha⁻¹. The phosphorus fertilization improves various metabolic and physiological process in life cycle of plant. The higher yields associated at increased levels of phosphorus are due to better root growth and increased uptake of nutrients favoring better crop growth”. [12]

“Application of potassium exerted a positive influence on yield. Application of 40 kg K₂O ha⁻¹ recorded maximum grain and straw yield which was 8.34 % and 6.45% more over 0 kg ha⁻¹ application of P₂O₅. [13]. [18] also registered “26.97% more grain yield with application of 30 kg K₂O ha⁻¹ compared to control”. “Grain acts as physiological sink and the strength of this sink depends on the number of endosperm cells. Potassium application has a beneficial influence on the development of endosperm cells and on the grain weight of cereals” [14]. Optimum supply of potassium aid in supply of photosynthate to the developing sink.

Conclusion

Therefore, taking the findings of the present study into consideration, it may be concluded that Application of 120 kg N ha⁻¹, 80 kg P₂O₅ ha⁻¹ and 40 kg K₂O ha⁻¹ resulted in maximum plant height, number of tillers, dry matter accumulation and leaf area index at all growth stages. Fertility level of 120 kg N ha⁻¹, 80 kg P₂O₅ ha⁻¹ and 40 kg K₂O ha⁻¹ produced highest grain and straw yield ha⁻¹. This study indicated that application of 120 kg N ha⁻¹, 80 kg P₂O₅ ha⁻¹ and 40 kg K₂O ha⁻¹ contributed towards better vegetative growth and yield mainly due to higher absorption of nutrients which increased photosynthates accumulation and high biomass production.

Reference

- [1] Hindu Survey. 2004. The Hindu Survey of Indian Agriculture 2004, M/s Kasturi and Sons Ltd., Chennai, Pp. 5-6.
- [2] Jahan, M.S., Sultana, S. and Ali, M.Y. (2014). Effect of different nitrogen levels on the yield performance of aromatic rice varieties. *Bull. Inst. Trop. Agr., Kyushu Univ.* 27:47-56.

- [3] Hossain, M.B., Islam, M.O. and Hasanuzzaman, M. (2008). Influence of different nitrogen levels on the performance of four aromatic rice varieties. *International Journal of Agriculture and Biology* **10**:693-696.
- [4] Babu, Venkatesh. D., Sudhakar, P., Srikanth, B., Reddy, Sirinivasa. And Reddy, CH. Bhargava. Rami. (2022). Effect of different levels of nitrogen on growth and yield parameter in rice (*Oryza sativa L.*) varieties. *The Pharma Innovation*, **11**(10):1378-1382.
- [5] Bharti, C., Ram. Vishram. and Patidar, Rahul. (2022). Growth, yield attributes and yield of rice as influenced by nitrogen levels and its split application in plateau of north eastern hilly region. *The Pharma Innovation*, SP-**11**(7):3585-3588.
- [6] Tsukru, Seiekevino., Singh, P.K. and Pandey, Manoj. (2023). Effect of nitrogen and phosphorus levels on growth and yield of direct-seeded rice (*Oryza sativa L.*) under Nagaland conditions. *Annals of Plant and Soil Research*, **25**(1): 120-126.
- [7] Prajapati, K. and Modi, H.A. (2012). Importance of potassium in plant growth-A review. *Indian Journal of Plant Sciences*, **1**(2-3):177-186.
- [8] Mukherjee, D. and Sen, A. (2005). Influence of rice husk and fertility levels on nutrient content of rice (*Oryza sativa*). *Agricultural Science Digest* **25**:151-152.
- [9] Singh, M., Yadav, D.B., Kumar, N., Kakraliya, S.K. and Khedwal, R.S. (2017). Performance of different basmati rice on phenology, growth, and quality under different nitrogen scheduling as dry DSR sown condition in IGP. *International Journal of Current Microbiology and Applied Sciences*, **6**:73-80.
- [10] Rao, K.V., Surekha, K., Kundu, D.K., Prasad, A.S.R. (2004). Nutrient management to sustain productivity targets of irrigated rice. In *International Symposium on Rice: From Green revolution to gene revolution*, 4-6.
- [11] Singh, Anando. N., Sorokhaibam, Sakhen. Pramanik, Kalipada. And Nabachandra, N. (2017). Effect of planting time and nitrogen fertilization on yield, nutrient uptake and nitrogen use efficiency of hybrid rice under rainfed shallow land of North East India. *Int. J. Curr. Microbial. App. Sci.* **6**(11):2111-2120.
- [12] Archana, K., Reddy, Prabhakar. T., Anjaiah. T. and Padmaja. B. (2017). Effect of levels phosphorus and its time of application on soil nutrient status and yield of rice grown on P accumulated soil. *International Journal Of Current Microbiology and Applied Sciences*, Special issue-**4**:92-99.
- [13] Brohi, A.R., Karaman, M.R., Topbas, M.T., Aktas, A. and Savasli, E. (2000). Effect of potassium and magnesium fertilization on yield and nutrient content of rice crop

- grown on artificial siltation soil. *Turkish Journal of Agriculture and Forestry*, **24**(1) 429-435.
- [14] Schacherer, A. and Beinger, H. (1984). Number and size distribution of endosperm cells in developing cereal grains as an index for their sink capacity. *Ber.deutsch.bot.ges.* 97,183-185 In: Mengel, K. and Kirby, E.A. (1996) Principle of plant nutrition (4th ed.) Panima publishing corporation, New Delhi.
- [15] Maurya, Rakesh., Singh, Manoj. Kumar., Singh, Nikhil. Kumar., Singh, Manish. Kumar. and Singh, Anurag. Kumar. (2021). Effect of nitrogen levels on growth attributes, yields and nutrient uptake of different rice (*Oryza sativa* L.) varieties under the transplanted condition. *Journal of Experimental Biology and Agricultural Sciences*, **9**(Spl-3-NRMCSSA_2021):S336-S342.
- [16] Nadan, Neha., Roy, D.K., Kumar, Shashank. and Kumar, Ashok. (2020). Effect of nitrogen on growth, yield and nutrient uptake of rice under aerobic condition. *International Journal of Current Microbiology and Applied Sciences*, **9**(2)2025:2035.
- [17] Birla, Vinod., Vyas, M.D., Dubey, Megha., Waskle, Usha. and Mandre, Basant. Kumar. (2020). Effect of different doses of potassium on growth, yield attributing characters of rice in vertisol soil of Madhya Pradesh, India. *International Journal of Current Microbiology and Applied Sciences*, **9** (3)2629:2642.
- [18] Ola, Santosh., Sharma, Neetu., Sharma, B.C., Kumar, Anil., Chand, Gurdev. and Puniya, R. (2019). Optimization of phosphorus and potassium levels for productivity enhancement of fine rice in irrigated sub tropics of Jammu. *Journal of Pharmacognosy and Phytochemistry*, **8**(2): 1329-1332.