
A New Approach to Dual Jacobsthal Split Quaternions With Different Polar Representation



Abstract

In this paper, we introduce split quaternions with components including dual Jacobsthal and dual Jacobsthal-Lucas number sequences. By using Binet's formulas of these type split quaternions we give an explicit form of classic polar representations of them, after that we demonstrate a new polar representation by using Cayley-Dikson's notation of split quaternions which is based on two complex numbers. Some fundamental properties and identities for these type of split quaternions are studied. In further the current paper, it would be valuable to replicate similar approaches polar representation with dual Jacobsthal Split quaternions

Keywords: Dual Jacobsthal split quaternions, Dual Jacobsthal-Lucas split quaternions, Polar representation

2010 Mathematics Subject Classification: 11B83; 11R52; 15A66

1 Introduction

Quaternions have found extensive applications in the field of 3D computer graphics, robotics, and physics, particularly in the representation and computation of rotations and orientations [1]. They offer a compact and efficient way to represent 3D rotations without suffering from gimbal lock, an issue prevalent with Euler angles [2]. Moreover, quaternions are used in the analysis of spatial rotations in physics and in the control mechanisms of spacecraft and robotic arms due to their ability to interpolate rotations smoothly and efficiently [3].

Quaternions, extension of complex numbers was introduced by W. R. Hamilton as the following quadruple

$$Q = \gamma_0 + \gamma_1 i_1 + \gamma_2 i_2 + \gamma_3 i_3$$

where $\gamma_0, \gamma_1, \gamma_2, \gamma_3 \in \mathbb{R}$ and $i_1^2 = i_2^2 = i_3^2 = -1, i_1 i_2 = -i_2 i_1 = i_3$. Any quaternion can be write as $Q = (\gamma_0 + \gamma_1 i_1) + i_2(\gamma_2 - \gamma_3 i_1)$, where $\gamma_0 + \gamma_1 i_1$ and $\gamma_2 - \gamma_3 i_1$ are complex numbers, therefore quaternions are one of the hyper complex numbers,[4-8].

J. Cokle defined the split quaternion as $\hat{Q} = \delta_0 + \delta_1 i_1 + \delta_2 i_2 + \delta_3 i_3$ with $\delta_0, \delta_1, \delta_2, \delta_3 \in \mathbb{R}$ and quatenionic units i_1, i_2, i_3 satisfy

$$i_1^2 = -i_2^2 = -i_3^2 = -1, i_1 i_2 = -i_2 i_1 = i_3 \quad (1.1)$$

and can be shown as $\hat{Q} = S_{\hat{Q}} + V_{\hat{Q}}$, \hat{Q} with $S_{\hat{Q}} = 0$ is a pure split quaternion, [9-13].

Let $\hat{Q}_1 = S_{\hat{Q}_1} + V_{\hat{Q}_1}$ and $\hat{Q}_2 = S_{\hat{Q}_2} + V_{\hat{Q}_2}$ be split quaternions, then addition and multiplication are

$$\begin{aligned} \hat{Q}_1 + \hat{Q}_2 &= (S_{\hat{Q}_1} + S_{\hat{Q}_2}) + (V_{\hat{Q}_1} + V_{\hat{Q}_2}) \\ \hat{Q}_1 \hat{Q}_2 &= S_{\hat{Q}_1} S_{\hat{Q}_2} + \langle V_{\hat{Q}_1}, V_{\hat{Q}_2} \rangle + S_{\hat{Q}_1} V_{\hat{Q}_2} + S_{\hat{Q}_2} V_{\hat{Q}_1} + V_{\hat{Q}_1} \times V_{\hat{Q}_2} \end{aligned}$$

respectively, where $\langle \cdot, \cdot \rangle$ and \times are inner and cross products in Minkowsky space \mathbb{E}_1^3 . The conjugate and norm of $\hat{Q} = \delta_0 + \delta_1 i_1 + \delta_2 i_2 + \delta_3 i_3$ are respectively as $\bar{\hat{Q}} = \delta_0 - \delta_1 i_1 - \delta_2 i_2 - \delta_3 i_3$ and

$$N(\hat{Q}) = \sqrt{|\hat{Q} \bar{\hat{Q}}|} = \sqrt{|\delta_0^2 + \delta_1^2 - \delta_2^2 - \delta_3^2|} \quad (1.2)$$

if $N(\hat{Q}) = 1$, then \hat{Q} is a unit split quaternion, for any split quaternion \hat{Q} with $N(\hat{Q}) \neq 0$, $\frac{\hat{Q}}{N(\hat{Q})}$ is a unit split quaternion. Let $I_{\hat{Q}} = \delta_0^2 + \delta_1^2 - \delta_2^2 - \delta_3^2$, then \hat{Q} is space-like, time-like and light-like if $I_{\hat{Q}} > 0$, $I_{\hat{Q}} < 0$ and $I_{\hat{Q}} = 0$ respectively, the multiplicative inverse of \hat{Q} is $\hat{Q}^{-1} = \frac{\bar{\hat{Q}}}{N(\hat{Q})^2}$ and there is no inverse for light-like split quaternion. The Cayley-Dickson's form of a split quaternion \hat{Q} is $\hat{Q} = (\delta_0 + \delta_1 i_1) + (\delta_2 + \delta_3 i_1) i_2$ which is based on two complex numbers.

Dual numbers, extension of real numbers, first were defined by W. K. Clifford in 1873 as:

$$\mathbb{D} = \{A = a_1 + \varepsilon a_2 \mid a_1, a_2 \in \mathbb{R}, \varepsilon \neq 0, \varepsilon^2 = 0\}$$

This new number system is commutative and associative algebra over real numbers wich has dimension two. Kotelnikov initiated the study of the first applications of dual numbers. Eduard Study used these numbers and associated vectors in line geometry and kinematics,[14,15].

According to the following operation

$$\begin{aligned} (a_1 + \varepsilon a_2) + (b_1 + \varepsilon b_2) &= (a_1 + b_1) + \varepsilon(a_2 + b_2) \\ (a_1 + \varepsilon a_2)(b_1 + \varepsilon b_2) &= a_1 b_1 + \varepsilon(a_1 b_2 + a_2 b_1) \end{aligned}$$

algebra of dual numbers is a commutative ring, but not a field. The multiplicative inverse of A is

$$A^{-1} = \frac{1}{a_1} - \varepsilon \frac{a_2}{a_1^2}, a_1 \neq 0$$

and there is no inverse for pure dual numbers and hence this algebra of numbers is not field over real numbers. By using inverse of dual numbers we can define division operation of two dual numbers A_1 and B_1 as $A_1 B_1^{-1}$ where B_1 is not a pure dual number and $B_1 \neq 0$. Dual angle between lines d_1 and d_2 in \mathbb{R}^3 is defined as $\Phi = \phi + \varepsilon \phi^*$, where ϕ is angle and ϕ^* is the shortest distance between these lines. If $f(x + \varepsilon x^*)$ is a dual function, then by Taylor expansion we have

$$f(x + \varepsilon x^*) = f(x) + \varepsilon x^* f'(x) \quad (1.3)$$

where $f'(x)$ is the first derivative of $f(x)$, using Equality (1.3) we can write

$$\sqrt{a_1 + \varepsilon a_2} = \sqrt{a_1} + \varepsilon \frac{a_2}{2\sqrt{a_1}} \quad (1.4)$$

Dual split quaternion \mathbb{Q} is defined as

$$\mathbb{Q} = A + B i_1 + C i_2 + D i_3$$

where A, B, C, D are dual numbers and i_1, i_2, i_3 follows rules in (1.1), \mathbb{Q} can be write as $\mathbb{Q} = \hat{Q} + \varepsilon \hat{Q}^*$, where \hat{Q}, \hat{Q}^* are split quaternions and $\varepsilon^2 = 0$. Let $\mathbb{Q}_1 = \hat{Q}_1 + \varepsilon \hat{Q}_1^*$ and $\mathbb{Q}_2 = \hat{Q}_2 + \varepsilon \hat{Q}_2^*$ be dual split quaternions, then

$$\begin{aligned}\mathbb{Q}_1 + \mathbb{Q}_2 &= (\hat{Q}_1 + \hat{Q}_2) + \varepsilon(\hat{Q}_1^* + \hat{Q}_2^*) \\ \mathbb{Q}_1 \mathbb{Q}_2 &= \hat{Q}_1 \hat{Q}_2 + \varepsilon(\hat{Q}_1 \hat{Q}_2^* + \hat{Q}_1^* \hat{Q}_2)\end{aligned}$$

norm of \mathbb{Q} is defined as $N(\mathbb{Q}) = \sqrt{A^2 + B^2 - C^2 - D^2}$, and dual split quaternion with $N(\mathbb{Q}) = 1$ is called unit dual split quaternion. Dual split quaternion $\mathbb{Q} = \hat{Q} + \varepsilon \hat{Q}^*$ is spacelike, timelike or lightlike if and only \hat{Q} is spacelike, timelike and lightlike, respectively. Suppose $\Phi = \phi + \varepsilon \phi^*$ be dual angle then the polar representation for spacelike \mathbb{Q} is as

$$\mathbb{Q} = N(\mathbb{Q})(\sinh \Phi + \nu \cosh \Phi) \quad (1.5)$$

where $\nu = \frac{Bi_+Ci_2+Di_3}{\sqrt{-B^2+C^2+D^2}}$ is a unit pure dual split quaternion, $\sinh \Phi = \frac{D}{N(\mathbb{Q})}$ and $\cosh \Phi = \frac{\sqrt{|B^2-C^2-D^2|}}{N(\mathbb{Q})}$. The polar representation for timelike \mathbb{Q} with spacelike vector part is

$$\mathbb{Q} = N(\mathbb{Q})(\cosh \Phi + \nu \sinh \Phi) \quad (1.6)$$

where $\nu = \frac{Bi_+Ci_2+Di_3}{\sqrt{-B^2+C^2+D^2}}$ is a unit pure dual split quaternion, $\cosh \Phi = \frac{D}{N(\mathbb{Q})}$ and $\sinh \Phi = \frac{\sqrt{|B^2-C^2-D^2|}}{N(\mathbb{Q})}$ and the polar representation for timelike \mathbb{Q} with timelike vector part is

$$\mathbb{Q} = N(\mathbb{Q})(\cos \Phi + \nu \sin \Phi) \quad (1.7)$$

where $\nu = \frac{Bi_+Ci_2+Di_3}{\sqrt{B^2-C^2-D^2}}$ is a unit pure dual split quaternion, $\cos \Phi = \frac{D}{N(\mathbb{Q})}$ and $\sin \Phi = \frac{\sqrt{|B^2-C^2-D^2|}}{N(\mathbb{Q})}$, [16,17]. Any dual split quaternion can be written in the form $\mathbb{Q} = Ae^{B i_2}$, where $A = A_0 + A_1 i_1$ and $B = A_2 + A_3 i_1$ with A_0, A_1, A_2, A_3 dual numbers, [18].

The Jacobsthal J_n and Jacobsthal-Lucas j_n number sequences are defined by

$$\begin{aligned}J_n &= J_{n-1} + 2J_{n-2}, \quad J_0 = 0, J_1 = 1, n \geq 2 \\ j_n &= j_{n-1} + 2j_{n-2}, \quad j_0 = 2, j_1 = 1, n \geq 2\end{aligned}$$

The characteristic equation of these number sequences is $x^2 - x - 2 = 0$, with roots $\alpha = 2, \beta = -1$ and corresponding Binet's forms are

$$J_n = \frac{2^n - (-1)^n}{3} \quad (1.8)$$

$$j_n = 2^n + (-1)^n \quad (1.9)$$

For more details and relations about these number sequences, [19-22]. The n^{th} dual Jacobsthal \hat{J}_n and Jacobsthal-Lucas \hat{j}_n number sequences are defined by

$$\hat{J}_n = J_n + \varepsilon J_{n+1}$$

$$\hat{j}_n = j_n + \varepsilon j_{n+1}$$

where J_n and j_n are the n^{th} Jacobsthal and Jacobsthal-Lucas numbers, the Binet's formulas for \hat{J}_n and \hat{j}_n are

$$\hat{J}_n = \frac{2^n \alpha - (-1)^n \beta}{3} \quad (1.10)$$

$$\hat{j}_n = 2^n \alpha + (-1)^n \beta \quad (1.11)$$

where

$$\begin{aligned}\underline{\alpha} &= 1 + 2\varepsilon \\ \underline{\beta} &= 1 - \varepsilon\end{aligned}$$

Both sequences have applications and appear in various contexts in mathematics, including combinatorial identities, the study of geometric shapes, and connections to the Fibonacci and Lucas numbers. Their properties have been studied in depth, revealing fascinating aspects such as their relationships with binary representations and the golden ratio, though in a manner distinct from the Fibonacci and Lucas numbers.

The study of these sequences not only contributes to the field of number theory but also enhances our understanding of the elegant structures and patterns inherent in mathematics.

2 Main results

In this section we defined dual Jacobsthal and dual Jacobsthal-Lucas split quaternions, polar representations and some properties of these split quaternions are discussed.

Definition 2.1. The dual Jacobsthal and dual Jacobsthal-Lucas split quaternions are defined as

$$\mathcal{J}_n = \hat{J}_n + \hat{J}_{n+1}i_1 + \hat{J}_{n+2}i_2 + \hat{J}_{n+3}i_3 \quad (2.1)$$

$$\tilde{\mathcal{J}}_n = \hat{j}_n + \hat{j}_{n+1}i_1 + \hat{j}_{n+2}i_2 + \hat{j}_{n+3}i_3 \quad (2.2)$$

respectively, where \hat{J}_n is n^{th} dual jacobsthal, \hat{j}_n is n^{th} dual jacobsthal-Lucas number and i_1, i_2, i_3 follow the rules in (1.1).

From definition, the following recurrence relation can be prove easily

$$\mathcal{J}_n = \mathcal{J}_{n-1} + 2\mathcal{J}_{n-2}, \quad n \geq 2$$

and

$$\tilde{\mathcal{J}}_n = \tilde{\mathcal{J}}_{n-1} + 2\tilde{\mathcal{J}}_{n-2}, \quad n \geq 2$$

Theorem 2.1. The Binet's formulas for dual Jacobsthal and dual Jacobsthal-Lucas split quaternions are

$$\mathcal{J}_n = \frac{2^n \hat{\alpha} - (-1)^n \hat{\beta}}{3} \quad (2.3)$$

$$\tilde{\mathcal{J}}_n = 2^n \hat{\alpha} + (-1)^n \hat{\beta} \quad (2.4)$$

respectively, where

$$\begin{aligned}\underline{\alpha} &= (1 + 2\varepsilon)(1 + 2i_1 + 4i_2 + 8i_3) \\ \underline{\beta} &= (1 - \varepsilon)(1 - i_1 + i_2 - i_3)\end{aligned}$$

Proof. The proof can be done directly by using Equalities (1.10),eq1.11,2.1,(2.2). □

Proposition 2.1. The norm of dual Jacobsthal and dual Jacobsthal-Lucas split quaternions is

$$\begin{aligned}N(\mathcal{J}_n) &= \sqrt{(2^{2n+3} + \frac{1}{3}(j_n^2 - 1)) + \varepsilon(2^{2n+5} + j_{2n} + \frac{1}{3}(j_n^2 - 4))} \\ N(\tilde{\mathcal{J}}_n) &= 3 \sqrt{(2^{2n+3} + 3J_n^2 - \frac{1}{3}) + \varepsilon(2^{2n+5} + 3J_n^2 + j_{2n} - \frac{4}{3})}\end{aligned}$$

where J_n and j_n are n^{th} Jacobsthal and Jacobsthal-Lucas numbers, respectively.

Proof. From Equality (1.2), we have

$$N(\mathcal{J}_n) = \sqrt{|\hat{J}_n^2 + \hat{J}_{n+1}^2 - \hat{J}_{n+2}^2 - \hat{J}_{n+3}^2|}$$

and by using Equality (1.10), we obtain

$$\begin{aligned} \hat{J}_n^2 + \hat{J}_{n+1}^2 - \hat{J}_{n+2}^2 - \hat{J}_{n+3}^2 &= \left(\frac{2^n \underline{\alpha} - (-1)^n \underline{\beta}}{3} \right)^2 + \left(\frac{2^{n+1} \underline{\alpha} - (-1)^{n+1} \underline{\beta}}{3} \right)^2 \\ &\quad - \left(\frac{2^{n+2} \underline{\alpha} - (-1)^{n+2} \underline{\beta}}{3} \right)^2 - \left(\frac{2^{n+3} \underline{\alpha} - (-1)^{n+3} \underline{\beta}}{3} \right)^2 \\ &= -\frac{1}{3} (25 \cdot 2^{2n} (1 + 4\varepsilon) + (-1)^n 2^{n+1} (1 + \varepsilon)) \end{aligned}$$

Finally by using Equality (1.8) and doing basic calculations, we get

$$N(\mathcal{J}_n) = \sqrt{(2^{2n+3} + \frac{1}{3}(j_n^2 - 1)) + \varepsilon(2^{2n+5} + j_{2n} + \frac{1}{3}(j_n^2 - 4))}$$

On the other hand for $N(\tilde{\mathcal{J}}_n)$, by using Equality (1.11) we have

$$\begin{aligned} \hat{J}_n^2 + \hat{J}_{n+1}^2 - \hat{J}_{n+2}^2 - \hat{J}_{n+3}^2 &= (2^n \underline{\alpha} + (-1)^n \underline{\beta})^2 + (2^{n+1} \underline{\alpha} + (-1)^{n+1} \underline{\beta})^2 \\ &\quad - (2^{n+2} \underline{\alpha} + (-1)^{n+2} \underline{\beta})^2 - (2^{n+3} \underline{\alpha} + (-1)^{n+3} \underline{\beta})^2 \\ &= -3(25 \cdot 2^{2n} (1 + 4\varepsilon) + (-1)^{n+1} 2^{n+1} (1 + \varepsilon)) \end{aligned}$$

And by doing some necessary calculations, we get

$$N(\tilde{\mathcal{J}}_n) = 3 \sqrt{(2^{2n+3} + 3J_n^2 - \frac{1}{3}) + \varepsilon(2^{2n+5} + 3J_n^2 + j_{2n} - \frac{4}{3})}$$

□

Corollary 2.2. *The dual Jacobsthal and dual Jacobsthal-Lucas split quaternions are spacelike split quaternions with spacelike vector part.*

Theorem 2.3. *The classical polar representation of dual Jacobsthal split quaternion is*

$$\mathcal{J}_n = N(\mathcal{J}_n)(\sinh \Phi + \nu \cosh \Phi)$$

where

$$\nu = \frac{2^{n+1}(1 + 2\varepsilon)(i_1 + 2i_2 + 4i_3) + (-1)^n(1 - \varepsilon)(i_1 - i_2 + i_3)}{3 \sqrt{(J_{n+1}^2 + 2^{2n+3}) + \varepsilon(J_{n+1}^2 + J_{2n+2} + 2^{2n+5})}}$$

is a pure unit dual split quaternion and Φ is a dual angle such that

$$\Phi = \tanh^{-1} \left(\frac{J_n}{\sqrt{J_{n+1}^2 + 2^{2n+3}}} \right) + \varepsilon \frac{j_n J_{n+1}^2 + 2^{2n+4}(-1)^n - J_n J_{2n+2}}{\sqrt{J_{n+1}^2 + 2^{2n+3}} (2^{2n+4} + \frac{2}{3}(j_n^2 - 1))}$$

Proof. Since \mathcal{J}_n is a spacelike dual split quaternion, then by using Equality (1.5), the polar representation is

$$\mathcal{J}_n = N(\mathcal{J}_n)(\sinh \Phi + \nu \cosh \Phi)$$

where

$$\begin{aligned}\nu &= \frac{\hat{J}_{n+1}i_1 + \hat{J}_{n+2}i_2 + \hat{J}_{n+3}i_3}{\sqrt{-\hat{J}_{n+1}^2 + \hat{J}_{n+2}^2 + \hat{J}_{n+3}^2}} \\ \sinh \Phi &= \frac{\hat{J}_n}{N(\mathcal{J}_n)} \\ \cosh \Phi &= \frac{\sqrt{|\hat{J}_{n+1}^2 - \hat{J}_{n+2}^2 - \hat{J}_{n+3}^2|}}{N(\mathcal{J}_n)}\end{aligned}$$

therefore $\tanh \Phi = \frac{\hat{J}_n}{\sqrt{-\hat{J}_{n+1}^2 + \hat{J}_{n+2}^2 + \hat{J}_{n+3}^2}}$ and by using Equality (1.10) we get

$$\nu = \frac{2^{n+1}(1+2\varepsilon)(i_1 + 2i_2 + 4i_3) - (-1)^n(1-\varepsilon)(i_1 - i_2 + i_3)}{\sqrt{(J_{n+1}^2 + 2^{2n+3}) + \varepsilon(J_{n+1}^2 + 2^{2n+5} + J_{2n+2})}}$$

and

$$\tanh \Phi = \frac{J_n + \varepsilon J_{n+1}}{\sqrt{(J_{n+1}^2 + 2^{2n+3}) + \varepsilon(J_{n+1}^2 + 2^{2n+5} + J_{2n+2})}}$$

Using Equality (1.4) and doing necessary calculations we will have

$$\Phi = \tanh^{-1} \left(\frac{J_n}{\sqrt{J_{n+1}^2 + 2^{2n+3}}} + \varepsilon \frac{j_n J_{n+1}^2 + 2^{2n+4}(-1)^n - J_n J_{2n+2}}{2(J_{n+1}^2 + 2^{2n+3})^{\frac{3}{2}}} \right)$$

Finally by using Equality (1.3) we obtain

$$\Phi = \tanh^{-1} \left(\frac{J_n}{\sqrt{J_{n+1}^2 + 2^{2n+3}}} \right) + \varepsilon \frac{j_n J_{n+1}^2 + 2^{2n+4}(-1)^n - J_n J_{2n+2}}{\sqrt{J_{n+1}^2 + 2^{2n+3}} (2^{2n+4} + \frac{2}{3}(j_n^2 - 1))}$$

□

Corollary 2.4. *The classical polar representation of dual Jacobsthal-Lucas split quaternions is*

$$\tilde{\mathcal{J}}_n = N(\tilde{\mathcal{J}}_n)(\sinh \hat{\Phi} + \hat{\nu} \cosh \hat{\Phi})$$

where

$$\hat{\nu} = \frac{2^{n+1}(1+2\varepsilon)(i_1 + 2i_2 + 4i_3) - (-1)^n(1-\varepsilon)(i_1 - i_2 + i_3)}{3\sqrt{2^{2n+3} + (\frac{j_{n+1}}{3})^2} + \varepsilon(2^{2n+5} + J_{2n+2} + (\frac{j_{n+1}}{3})^2)}$$

is a pure unit dual split quaternion and $\hat{\Phi}$ is a dual angle such that

$$\hat{\Phi} = \tanh^{-1} \left(\frac{j_n}{\sqrt{j_{n+1}^2 + 9 \cdot 2^{2n+3}}} \right) + \varepsilon \frac{J_n j_{n+1}^2 - 3 \cdot 2^{2n+4}(-1)^n - j_n J_{2n+2}}{\sqrt{j_{n+1}^2 + 9 \cdot 2^{2n+3}} (2^{2n+4} + 6J_n^2 - \frac{2}{3})}$$

Proof. The Proof is similar to Theorem (2.3). □

Proposition 2.2. *Let $P = Ai_2 + Bi_3 = (A + Bi_1)i_2$ be an arbitrary dual split quaternion, if P is spacelike then its exponential form is*

$$e^P = \sinh |P| + \frac{A}{|P|} \cosh |P|i_2 + \frac{B}{|P|} \cosh |P|i_3 = \alpha_0 + \alpha_2 i_2 + \alpha_3 i_3$$

and if P is timelike, then

$$e^P = \cosh |P| + \frac{A}{|P|} \sinh |P| i_2 + \frac{B}{|P|} \sinh |P| i_3 = \beta_0 + \beta_2 i_2 + \beta_3 i_3$$

That is, it is a dual split quaternions which does not contain i_1 's term.

Proof. Suppose μ is a spacelike unit dual split quaternion, that is $N(\mu) = 1$, then from Equality (1.5) we have

$$e^{\mu\theta} = \sinh \theta + \mu \cosh \theta$$

if we rewrite $P = |P| \frac{P}{|P|}$, then by taking $\mu = \frac{P}{|P|}$ and $\theta = |P|$ we get the result, we can prove similarly for timelike P by using Equality (1.6). \square

Now we give the new polar representations for dual Jacobsthal and dual Jacobsthal-Lucas split quaternions by using Cayley-Dikson's form for split quaternions.

Theorem 2.5. *Every dual Jacobsthal split quaternion $\mathcal{J}_n = \hat{J}_n + \hat{J}_{n+1}i_1 + \hat{J}_{n+2}i_2 + \hat{J}_{n+3}i_3$ can be given in the form $\mathcal{J}_n = \mathbb{A}e^{\mathbb{B}i_2}$, where \mathbb{A} and \mathbb{B} are dual Jacobsthal complex numbers, that is*

$$\mathbb{A} = \frac{\hat{J}_n + \hat{J}_{n+1}i_1}{\sqrt{\hat{J}_n^2 + \hat{J}_{n+1}^2}}$$

$$\mathbb{B} = \tanh^{-1} \left(\frac{\sqrt{\hat{J}_n^2 + \hat{J}_{n+1}^2}}{\sqrt{\hat{J}_{n+2}^2 + \hat{J}_{n+3}^2}} \right) \frac{(\hat{J}_n \hat{J}_{n+2} + \hat{J}_{n+1} \hat{J}_{n+3}) + (\hat{J}_n \hat{J}_{n+3} - \hat{J}_{n+1} \hat{J}_{n+2})i_1}{\sqrt{(\hat{J}_n^2 + \hat{J}_{n+1}^2)(\hat{J}_{n+2}^2 + \hat{J}_{n+3}^2)}}$$

Proof. Suppose that $\mathbb{A} = a + bi_1$ and $e^{\mathbb{B}i_2} = \alpha_0 + \alpha_2 i_2 + \alpha_3 i_3$, then

$$\mathcal{J}_n = \mathbb{A}e^{\mathbb{B}i_2} = a\alpha_0 + b\alpha_0 i_1 + (a\alpha_2 - b\alpha_3)i_2 + (a\alpha_3 + b\alpha_2)i_3$$

if $\alpha_0 = 0$, then we can select $a = 1$ and $b = 0$, we will get $\mathbb{A} = 1$. For $\alpha_0 \neq 0$, we construct a complex number $\psi = a\alpha_0 + b\alpha_0 i_1 = \hat{J}_n + \hat{J}_{n+1}i_1$ and then $\mathbb{A} = \frac{\psi}{|\psi|}$, by using Equality (1.10) the explicit form of \mathbb{A} is

$$\mathbb{A} = \frac{2^n(1 + 2\varepsilon)(1 + 2i_1) + (-1)^{n+1}(1 - \varepsilon)(1 - i_1)}{\sqrt{j_n^2 + 2^{2n+2} + \varepsilon(j_n^2 + 9J_{2n} + 2^{2n+4})}}$$

Since \mathbb{A} is a unit dual complex number then $\mathbb{A}^{-1} = \bar{\mathbb{A}} = \frac{\hat{J}_n - \hat{J}_{n+1}i_1}{\sqrt{\hat{J}_n^2 + \hat{J}_{n+1}^2}}$, where $\bar{\mathbb{A}}$ is conjugate of \mathbb{A} and

$$e^{\mathbb{B}i_2} = \bar{\mathbb{A}}\mathcal{J}_n = \frac{(\hat{J}_n^2 + \hat{J}_{n+1}^2) + (\hat{J}_n \hat{J}_{n+2} + \hat{J}_{n+1} \hat{J}_{n+3})i_2 + (\hat{J}_n \hat{J}_{n+3} - \hat{J}_{n+1} \hat{J}_{n+2})i_3}{\sqrt{\hat{J}_n^2 + \hat{J}_{n+1}^2}}$$

The norm of $e^{\mathbb{B}i_2}$ is

$$|e^{\mathbb{B}i_2}| = \sqrt{\hat{J}_{n+2}^2 + \hat{J}_{n+3}^2 - \hat{J}_n^2 - \hat{J}_{n+1}^2}$$

Since $\frac{e^{\mathbb{B}i_2}}{|e^{\mathbb{B}i_2}|}$ is a unit spacelike dual split quaternion, then its classical polar form is

$$\frac{e^{\mathbb{B}i_2}}{|e^{\mathbb{B}i_2}|} = \sinh \hat{\Phi} + \hat{\mu} \cosh \hat{\Phi}$$

where $\hat{\Phi} = \phi + \varepsilon\phi^*$ is dual angle, then we can write

$$\begin{aligned}\sinh \hat{\Phi} &= \frac{\sqrt{\hat{J}_n^2 + \hat{J}_{n+1}^2}}{\sqrt{\hat{J}_{n+2}^2 + \hat{J}_{n+3}^2 - \hat{J}_n^2 - \hat{J}_{n+1}^2}} \\ \cosh \hat{\Phi} &= \frac{\sqrt{\hat{J}_{n+2}^2 + \hat{J}_{n+3}^2}}{\sqrt{\hat{J}_{n+2}^2 + \hat{J}_{n+3}^2 - \hat{J}_n^2 - \hat{J}_{n+1}^2}} \\ \hat{\mu} &= \frac{(\hat{J}_n \hat{J}_{n+2} + \hat{J}_{n+1} \hat{J}_{n+3})i_2 + (\hat{J}_n \hat{J}_{n+3} - \hat{J}_{n+1} \hat{J}_{n+2})i_3}{\sqrt{(\hat{J}_n^2 + \hat{J}_{n+1}^2)(\hat{J}_{n+2}^2 + \hat{J}_{n+3}^2)}}\end{aligned}$$

which gives

$$\tanh \hat{\Phi} = \frac{\sqrt{\hat{J}_n^2 + \hat{J}_{n+1}^2}}{\sqrt{\hat{J}_{n+2}^2 + \hat{J}_{n+3}^2}}$$

Finally from $\mathbb{B}i_2 = \hat{\mu}\hat{\Phi}$ we get the result and by using (1.10) the explicit form of \mathbb{B} can be write easily. \square

Corollary 2.6. Every dual Jacobsthal-Lucas split quaternion $\tilde{\mathcal{J}}_n = \hat{j}_n + \hat{j}_{n+1}i_1 + \hat{j}_{n+2}i_2 + \hat{j}_{n+3}i_3$ can be given in the form $\tilde{\mathcal{J}}_n = \mathbb{A}e^{\mathbb{B}i_2}$, where \mathbb{A} and \mathbb{B} are dual Jacobsthal-Lucas complex numbers, that is

$$\begin{aligned}\mathbb{A} &= \frac{\hat{j}_n + \hat{j}_{n+1}i_1}{\sqrt{\hat{j}_n^2 + \hat{j}_{n+1}^2}} \\ \mathbb{B} &= \tanh^{-1} \left(\frac{\sqrt{\hat{j}_n^2 + \hat{j}_{n+1}^2}}{\sqrt{\hat{j}_{n+2}^2 + \hat{j}_{n+3}^2}} \right) \frac{(\hat{j}_n \hat{j}_{n+2} + \hat{j}_{n+1} \hat{j}_{n+3}) + (\hat{j}_n \hat{j}_{n+3} - \hat{j}_{n+1} \hat{j}_{n+2})i_1}{\sqrt{(\hat{j}_n^2 + \hat{j}_{n+1}^2)(\hat{j}_{n+2}^2 + \hat{j}_{n+3}^2)}}\end{aligned}$$

Proof. The proof can be done similar to Theorem (2.5). \square

Example 2.7. Find the new polar representation for $\mathcal{J}_1 = (1 + \varepsilon) + (1 + 3\varepsilon)i_1 + (3 + 5\varepsilon)i_2 + (5 + 11\varepsilon)i_3$. We have $\mathcal{J}_1 = \mathbb{A}e^{\mathbb{B}i_2}$, where

$$\mathbb{A} = \frac{\sqrt{2}}{2} ((1 - \varepsilon) + (1 + \varepsilon)i_1)$$

and

$$\mathbb{B} = \frac{1}{\sqrt{17}} \tanh^{-1} \left(\frac{1}{\sqrt{17}} + \varepsilon \frac{\sqrt{17} - 35\sqrt{2}}{17} \right) \left(\left(4 + \frac{13}{17}\varepsilon\right) + \left(1 - \frac{52}{17}\varepsilon\right)i_1 \right)$$

Theorem 2.8 (Catalan's identities). For positive integers n and r with $n \geq r \geq 1$, we have

$$\begin{aligned}\mathcal{J}_{n+r}\mathcal{J}_{n-r} - \mathcal{J}_n^2 &= \frac{1}{3}(-1)^{n-r}2^{n-r}J_r(1 + \varepsilon)(j_r(1 - 13i_1 + i_2 - 13i_3) + 2(-1)^r(-1 + i_1 + 5i_2 + 7i_3)) \\ \tilde{\mathcal{J}}_{n+r}\tilde{\mathcal{J}}_{n-r} - \tilde{\mathcal{J}}_n^2 &= 3(-1)^{n-r}2^{n-r}J_r(1 + \varepsilon)(j_r(-1 + 13i_1 - i_2 + 13i_3) - 2(-1)^r(-1 + i_1 + 5i_2 + 7i_3))\end{aligned}$$

where J_n and j_n are n^{th} jacobsthal and Jacobsthal-Lucas numbers.

Proof. By using (2.3), we get

$$\begin{aligned}
\mathcal{J}_{n+r}\mathcal{J}_{n-r} - \mathcal{J}_n^2 &= \frac{1}{3} (2^{n+r}\underline{\alpha} - (-1)^{n+r}\underline{\beta}) \frac{1}{3} (2^{n-r}\underline{\alpha} - (-1)^{n-r}\underline{\beta}) \\
&\quad - \frac{1}{3} (2^n\underline{\alpha} - (-1)^n\underline{\beta}) \frac{1}{3} (2^n\underline{\alpha} - (-1)^n\underline{\beta}) \\
&= \frac{1}{9} 2^n (-1)^n (\underline{\alpha}\underline{\beta} (1 - 2^r(-1)^{-r}) + \underline{\beta}\underline{\alpha} (1 - 2^{-r}(-1)^r)) \\
&= \frac{1}{3} 2^n (-1)^n J_r ((-1)^{r+1}\underline{\alpha}\underline{\beta} + 2^{-r}\underline{\beta}\underline{\alpha}) \\
&= \frac{1}{3} 2^{n-r} (-1)^{n-r} J_r ((-1)^r\underline{\beta}\underline{\alpha} - 2^r\underline{\alpha}\underline{\beta})
\end{aligned}$$

from using

$$\underline{\alpha}\underline{\beta} = (1 + \varepsilon)(-1 + 13i_1 - i_2 + 13i_3) \quad (2.5)$$

$$\underline{\beta}\underline{\alpha} = (1 + \varepsilon)(-1 - 11i_1 + 11i_2 + i_3) \quad (2.6)$$

we get the result. Similarly, the proof for Jacobsthal-Lucas can be done. \square

Corollary 2.9 (Cassini's identities). *For positive integer n , the following identities hold*

$$\begin{aligned}
\mathcal{J}_{n+1}\mathcal{J}_{n-1} - \mathcal{J}_n^2 &= (-1)^n 2^{n-1} (1 + \varepsilon) (-1 + 5i_1 + 3i_2 + 9i_3) \\
\tilde{\mathcal{J}}_{n+1}\tilde{\mathcal{J}}_{n-1} - \tilde{\mathcal{J}}_n^2 &= 9(-1)^{n-1} 2^{n-1} (1 + \varepsilon) (-1 + 5i_1 + 3i_2 + 9i_3)
\end{aligned}$$

Proof. The proof can be done by taking $r = 1$ in Theorem 2.8. \square

Theorem 2.10 (d'Ocagne's identities). *For positive integers n and m with $n \geq m$ we have*

$$\begin{aligned}
\mathcal{J}_{m+1}\mathcal{J}_n - \mathcal{J}_m\mathcal{J}_{n+1} &= \frac{1+\varepsilon}{3} (2^n(-1)^m(-1 - 11i_1 + 11i_2 + i_3) - 2^m(-1)^n(-1 + 13i_1 - i_2 + 13i_3)) \\
\tilde{\mathcal{J}}_{m+1}\tilde{\mathcal{J}}_n - \tilde{\mathcal{J}}_m\tilde{\mathcal{J}}_{n+1} &= 3(1 + \varepsilon) (2^m(-1)^n(-1 + 13i_1 - i_2 + 13i_3) - 2^n(-1)^m(-1 - 11i_1 + 11i_2 + i_3))
\end{aligned}$$

Proof. By using Equality (2.3) we have

$$\begin{aligned}
\mathcal{J}_{m+1}\mathcal{J}_n - \mathcal{J}_m\mathcal{J}_{n+1} &= \frac{1}{3} (2^{m+1}\underline{\alpha} - (-1)^{m+1}\underline{\beta}) \frac{1}{3} (2^n\underline{\alpha} - (-1)^n\underline{\beta}) \\
&\quad - \frac{1}{3} (2^m\underline{\alpha} - (-1)^m\underline{\beta}) \frac{1}{3} (2^{n+1}\underline{\alpha} - (-1)^{n+1}\underline{\beta}) \\
&= \frac{1}{9} (-\underline{\alpha}\underline{\beta} 2^{m+1}(-1)^n - \underline{\beta}\underline{\alpha} (-1)^{m+1}2^n + \underline{\alpha}\underline{\beta} (-1)^{n+1}2^m + \underline{\beta}\underline{\alpha} 2^{n+1}(-1)^m) \\
&= \frac{1}{3} (\underline{\beta}\underline{\alpha} 2^n(-1)^m - \underline{\alpha}\underline{\beta} 2^m(-1)^n)
\end{aligned}$$

Finally by using Equalities (2.5) and (2.6) the result is clear. The proof for Jacobsthal-Lucas can be done similarly by using Equalities (2.4), (2.5) and (2.6). \square

3 CONCLUSIONS

In this paper, we define dual Jacobsthal and dual Jacobsthal-Lucas split quaternions. The polar representations of these split quaternions have been obtained similar to the real quaternions. For this, the modulus and argument have been calculated from an arbitrary split quaternion.

Our work opens up a plethora of avenues for future research. One particularly promising direction is the application of these concepts to dual split quaternions, incorporating the sequences of Jacobsthal and Jacobsthal-Lucas numbers. This approach is not just a mere extension of our current framework

but also a potential pathway to uncover new relationships and properties within the realm of quaternionic analysis. In conclusion, our study not only adds a new chapter to the field of quaternionic analysis but also sets the stage for a deeper and more comprehensive exploration of mathematical structures that intertwine number sequences with quaternionic frameworks. As we continue to unravel the mysteries of these fascinating mathematical entities, we anticipate that our work will inspire further research and collaboration in this dynamic and ever-evolving field.

References

- [1] Kuipers JB. *Quaternions and Rotation Sequences: A Primer with Applications to Orbits, Aerospace, and Virtual Reality*. NJ: Princeton University Press, Princeton, 1999.
- [2] Shoemake K. Animating rotation with quaternion curves. *Computer Graphics (SIGGRAPH '85 Proceedings)*. 1985;19(3):245-254.
- [3] Dam EB, Koch M, Lillholm M. *Quaternions, interpolation and animation*. Technical report, Department of Computer Science. University of Copenhagen, Denmark, 1998.
- [4] Altmann SL. *Rotations, Quaternions, and Double Groups*. Oxford University Press. 1986; Oxford.
- [5] Girard PR. The Quaternion Group and Modern Physics. *Eur. J. Phys.* 1984;5: 25–32.
- [6] Hamilton WR. On quaternions or on a new system of imaginaries in algebra. *Philos. Mag.* 1884;25(3):489-495.
- [7] Horadam, AF. Quaternion Recurrence Relations. *Ulam Quaterly*. 1993; 2: 23–33.
- [8] Iakin AL. Generalized Quaternions of Higher Order. *Fibonacci Quart.* 1977; 15: no. 4, 343–346.
- [9] Cockle J. On systems of algebra involving more than one imaginary; and on equations of the fifth degree. *Philosophical Magazine*. 1849; 35(238):434-437.
- [10] Ghadami R, Rahebi J, Yaylı Y. Spline split quaternion interpolation in Minkowski space. *Adv. Appl. Cliff. Algebr.* 2013;23(4):849–862
- [11] Kula L, Yaylı Y. Split quaternions and rotations in semi-euclidean space R_2^4 . *J. Korean Math. Soc.* 2007;44(6):1313–1327.
- [12] O'Neill B. *Semi-Riemannian Geometry*. Academic Press. New York. 1983.
- [13] Ozdemir M. Roots of a split quaternion. *Applied Mathematics in Science*. 2009; 22:258–263.
- [14] Hamilton WR. On quaternions or on a new system of imaginaries in algebra. *Philos. Mag.* 1884;25(3):489-495.
- [15] Study E. *Geometry der dynamen*, Leipzig. 1901.
- [16] Kula L, Yaylı Y. Dual Split quaternions and screw motion in Minkowski 3-space. *Iran. J. Sci. Technol. Trans.* 2006; A30(A3):1–14.
- [17] Sangwine SJ, Bihan NL. Quaternion polar representation with a complex modulus and complex argument inspired by the Cayley-Dickson form. *Adv. Appl. Clifford Alg.* 2010; 20(1):111–120.

-
- [18] Atasoy A, Ata E, Yaylı Y, Kemer Y. A New Polar Representation for Split and Dual Split Quaternions. *Adv. Appl. Clifford Algebras*. 2017; 27:2307–2319.
- [19] Catarino P, Vasco P, Campos H, Aires AP, Borges A. New families of Jacobsthal and Jacobsthal-Lucas numbers. *Algebra Discret. Math.*2015;20: 40–54.
- [20] Horadam AF. Jacobsthal representation numbers. *Fibonacci Quarterly*. 1996; 34: 40-54.
- [21] Horadam AF. Jacobsthal representation polynomials. *Fibonacci Quarterly*. 1997;35:137–148.
- [22] Horadam AF. Jacobsthal and pell curves. *Fibonacci Quart.*1988; 26:79–83.

©2011 Author1 & Author2; This is an Open Access article distributed under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/2.0>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.