

IPM Essentials: Combining Biology, Ecology, and Agriculture for Sustainable Pest Control

Abstract

Integrated Pest Management (IPM) represents a paradigm shift in pest control, moving away from heavy reliance on chemical pesticides to a more sustainable, environmentally friendly approach. This article explores IPM, an ecosystem-based strategy that integrates biological, ecological, and agricultural sciences to achieve long-term pest control in agriculture. IPM emphasizes understanding pest life cycles and their interaction with the environment, utilizing a combination of techniques including biological control, cultural practices, mechanical and physical barriers, and targeted chemical interventions. Regular monitoring and informed decisionmaking form the crux of this approach, focusing on economically viable and environmentally responsible pest control methods. The article highlights various success stories, the challenges faced in implementing IPM, and future directions including the incorporation of precision agriculture technologies and genetic advancements. Overall, IPM emerges as a crucial element in sustainable agriculture, promising to maintain ecological balance while ensuring effective pest management and provides an in-depth examination of Integrated Pest Management (IPM), a multifaceted approach to sustainable pest control that synergizes biology, ecology, and agricultural science. IPM represents a paradigm shift from traditional, chemically intensive pest control methods to a more holistic, environmentally conscious framework. The core of IPM lies in understanding the life cycles and ecological interactions of pests, employing a diverse array of strategies including biological control through natural predators, cultural practices like crop rotation, mechanical and physical barriers, and judicious use of chemical pesticides. The article underscores the importance of regular monitoring and decisionmaking based on established thresholds to maintain an effective, economically viable, and ecologically responsible pest management system. Case studies highlighting the successful implementation of IPM in various agricultural settings are discussed, alongside the challenges and prospects of IPM, particularly in the context of climate change and technological advancements. The article concludes that IPM is not only essential for sustainable pest control but also pivotal in ensuring long-term agricultural productivity and environmental health.

Keywords: pest management, agriculture, IPM, pesticides, chemicals

Introduction

Agriculture, the backbone of human civilization, has always faced the daunting challenge of pest control. In this perpetual battle, the introduction of Integrated Pest Management (IPM) has been a gamechanger, offering a ray of hope for sustainable agriculture. As we stand at the crossroads of increasing food demands and environmental conservation, understanding and implementing IPM is more crucial than ever [1-5]. This article aims to delve into the essence of IPM, exploring how this innovative approach combines biology, ecology, and agricultural science to create a sustainable and effective framework for managing pests. For centuries, farmers around the world grappled with pests using traditional methods that were often environmentally harmful and unsustainable in the long term. The postWorld War II era saw a significant shift with the widespread adoption of synthetic pesticides, which promised efficient pest control but eventually led to a host of problems including pest resistance, environmental pollution, and health hazards [6-9].

The Emergence of IPM: A Multidisciplinary Approach

Integrated Pest Management emerged as a response to these challenges, bringing forth a paradigm shift in how we approach pest control. It is not merely a set of practices but a philosophy that integrates various scientific disciplines. By leveraging knowledge from biology to understand pest behaviors and lifecycles, applying ecological principles to assess pest interactions with the environment, and utilizing agricultural science for implementing practical field strategies, IPM stands as a multifaceted approach to pest control [10].

The Principles of IPM

At its core, IPM is based on the principles of ecological balance and minimal environmental impact. It advocates for a judicious mix of biological control methods, such as using natural predators, cultural techniques like crop rotation, and mechanical means including traps and barriers. Chemical control is not eliminated but is used more judiciously and as a last resort. A critical aspect of IPM is continuous monitoring and decisionmaking based on scientific research and field observations [11].




Fig 1. IPM is a system of managing pests which is designed to be sustainable.

Source: <https://tracex.tech.com/integrated-pest-management-for-sustainable-agriculture/>

The Global Impact and Future Directions

IPM has not only revolutionized pest control but also played a pivotal role in shaping sustainable agricultural practices globally [12]. Its influence extends beyond farms, impacting policy decisions, research directions, and educational programs. The integration of emerging technologies such as precision agriculture, data analytics, and biotechnology in IPM promises further advancements in sustainable pest management. As we embark on this exploration of agricultural landscape, IPM is not just a set of techniques; it is a philosophy that respects and works with nature's balance, aiming for a sustainable future where agricultural productivity and ecological health go hand in hand [13-15].

The Genesis of IPM: A Response to Agricultural Challenges

The story of Integrated Pest Management (IPM) is deeply rooted in the agricultural history of humanity. Prior to the advent of IPM, farmers relied heavily on simple yet crude methods for pest control. These methods, often based on trial and error, ranged from the use of natural predators to the application of rudimentary pesticides like sulfur and lime. The agricultural revolution brought about a significant change in these practices, particularly with the

introduction of synthetic pesticides in the mid-20th century. The post-World War II era witnessed a dramatic increase in pesticide use, driven by the discovery and widespread adoption of synthetic chemicals like DDT. These chemicals offered an effective and quick solution to pest problems and were hailed as a revolution in agricultural productivity. However, this 'pesticide boom' soon revealed its downsides. Issues such as pest resistance, environmental pollution, and the decline of nontarget species, including beneficial insects and birds, started surfacing. The impact on human health also became a growing concern, as highlighted in Rachel Carson's seminal work, "Silent Spring," which documented the detrimental effects of indiscriminate pesticide use. The increasing awareness of the negative consequences of heavy pesticide reliance catalyzed the search for alternative pest control methods. This led to the conceptualization of Integrated Pest Management in the 1960s. IPM was born out of a necessity to balance the need for pest control with environmental conservation and human health concerns. It proposed a more scientific approach to pest management, one that integrated multiple methods and was grounded in ecological principles [16-19].

Initial adoption of IPM was driven by academia and forward-thinking agriculturalists. Universities began researching and promoting IPM techniques, focusing on understanding pest biology, environmental factors, and the ecological impact of pest control methods. The development of IPM was gradual, involving the integration of biological controls, such as the use of natural predators and parasites, alongside cultural practices like crop rotation and soil management. The evolution of IPM has been marked by a continuous integration of new technologies and scientific advancements. From the incorporation of pheromone traps and genetically modified crops to the use of data analytics and precision agriculture, IPM strategies have become increasingly sophisticated. This modern era of IPM not only addresses the immediate need for pest control but also focuses on long-term sustainability, ecological balance, and food safety [20].

Today, IPM is globally recognized as a key component of sustainable agriculture. Organizations like the Food and Agriculture Organization of the United Nations (FAO) and various governmental bodies worldwide promote IPM practices. However, challenges remain, including the need for widespread education among farmers, adaptation to climate change, and the economic implications of shifting from traditional methods to IPM practices. The genesis of Integrated Pest Management was a critical response to the challenges posed by traditional agricultural practices, particularly the overreliance on chemical pesticides. Its

evolution reflects a growing understanding of the intricate balance between agricultural productivity, environmental health, and human wellbeing. As the world continues to grapple with the dual challenges of feeding a growing population and preserving natural resources, IPM stands out as a beacon of sustainable agricultural practices [21]. Integrated Pest Management (IPM) is not just an alternative method for controlling agricultural pests; it's a holistic approach that encompasses a wide range of disciplines including biology, ecology, and agronomy. This multifaceted strategy transcends the traditional view of pest control, which primarily focuses on the elimination of pests, and instead emphasizes a balanced, sustainable interaction between agricultural practices and the environment.

The Foundations of IPM

The foundation of IPM lies in its comprehensive understanding of the ecological systems within which pests operate. This involves studying pest life cycles, their interactions with other organisms, and their responses to environmental conditions. Unlike conventional methods that often rely on reactive, blanket applications of pesticides, IPM promotes a proactive and targeted approach. This approach is based on scientific research and field observations, leading to more effective and sustainable pest management strategies [22-24].

Key Components of IPM

1. **Preventive Cultural Practices:** This involves modifying farming practices to reduce the conditions that are conducive to pest outbreaks. Techniques like crop rotation, intercropping, and selecting pest-resistant crop varieties can significantly reduce the vulnerability of crops to pests.
2. **Biological Control:** Utilizing natural enemies of pests, such as predators, parasitoids, and pathogens, is a cornerstone of IPM. This method harnesses the power of nature to maintain pest populations at lower levels.

3. Mechanical and Physical Controls: Implementing physical barriers, traps, and other mechanical devices can effectively reduce pest populations without the need for chemical interventions.

4. Chemical Control as a Last Resort: In IPM, chemical pesticides are used only when necessary and in a targeted, specific manner to minimize their impact on the environment and nontarget organisms.

5. Regular Monitoring and Decision Making: Continuous monitoring of pest and beneficial organism populations is integral to IPM. Decisions regarding pest control interventions are made based on thorough assessments and preestablished thresholds.

Beyond Pest Control: Environmental and Economic Benefits

The implementation of IPM extends its benefits beyond effective pest control. By reducing the reliance on chemical pesticides, IPM contributes significantly to environmental health, preserving biodiversity, and protecting natural resources like soil and water. Economically, it helps farmers reduce costs associated with pesticide purchases and application, while also minimizing potential health risks to farmworkers and consumers.

IPM in Practice: A Dynamic Approach

IPM is not a static set of practices but a dynamic approach that adapts to changing conditions and advances in agricultural science. It involves continuous learning, adaptation, and integration of new techniques and technologies. From precision agriculture to the development of genetically modified crops that are more resistant to pests, IPM continues to evolve, reflecting the changing needs and challenges of modern agriculture. Understanding IPM is crucial in recognizing its role as a comprehensive, sustainable solution to pest management. By balancing the immediate needs for pest control with longterm environmental and economic considerations, IPM offers a forwardthinking approach to

agriculture that is in harmony with the ecosystem. It's a paradigm that not only addresses the challenge of pest control but also contributes to the broader goals of sustainable agriculture and environmental stewardship [25-27].

The Pillars of IPM

The Pillars of Integrated Pest Management (IPM)

Integrated Pest Management (IPM) is founded on a series of core principles or "pillars" that guide its approach to sustainable pest control. These pillars represent a comprehensive strategy that balances the need for effective pest management with environmental stewardship and economic viability. Below, we explore these foundational elements of IPM:

1. Biological Control

Concept: Biological control involves using living organisms such as predators, parasites, and pathogens to regulate pest populations.

Examples: Introducing ladybugs to control aphid populations or using *Bacillus thuringiensis* (Bt), a naturally occurring bacterium, to manage caterpillar pests.

Advantages: This approach is environmentally friendly and helps maintain the ecological balance by leveraging nature's own mechanisms for pest control.

2. Cultural Control

Concept: Cultural control involves modifying agricultural practices to reduce the prevalence and impact of pests.

Examples: Crop rotation to disrupt pest life cycles, choosing pest-resistant crop varieties, or altering planting and harvesting times to avoid peak pest seasons.

Advantages: These practices can be cost-effective and reduce the reliance on chemical controls, while also contributing to soil health and crop diversity.

3. Mechanical and Physical Controls

Concept: This pillar encompasses the use of physical methods or mechanical devices to manage pests.

Examples: Using barriers like nets or row covers to protect crops, employing traps for rodents or insects, and implementing tillage to disrupt the life cycle of soil pests.

Advantages: Mechanical and physical controls can provide immediate results and are often a straightforward approach to pest management without the use of chemicals.

4. Chemical Control

Concept: Chemical control, while used more judiciously in IPM, involves the application of pesticides to manage pest populations.

Examples: Targeted spraying of pesticides when pest populations reach a critical threshold, or the use of baits and spot treatments instead of widespread application.

Advantages: When used as part of an IPM strategy, chemical controls can be effective in managing pests that are difficult to control through other means. The focus is on minimal and smart use to reduce environmental impact.

5. Monitoring and Decision Making

Concept: Continuous monitoring and informed decisionmaking are central to IPM, ensuring that interventions are timely, targeted, and necessary.

Examples: Regular scouting for pests, using pheromone traps for monitoring insect populations, and employing action thresholds to determine when intervention is required.

Advantages: This proactive approach minimizes unnecessary interventions, reduces costs, and enhances the effectiveness of pest control measures.

6. Education and Awareness

Concept: Educating farmers, agricultural workers, and the community about IPM principles is crucial for its successful implementation.

Examples: Training programs, workshops, and extension services that provide knowledge on pest identification, life cycles, and IPM practices.

Advantages: Increased awareness and understanding of IPM promote its adoption and lead to more sustainable pest management practices across the agricultural sector.

The pillars of Integrated Pest Management represent a comprehensive and balanced approach to pest control. By integrating biological, cultural, mechanical, and chemical strategies, along with continuous monitoring and education, IPM not only addresses the immediate concerns of pest management but also upholds longterm environmental health and economic sustainability. This multifaceted approach is key to the future of sustainable agriculture and ecological conservation [28-30].

Challenges and Future Directions

While Integrated Pest Management (IPM) has made significant strides in promoting sustainable agriculture, it faces several challenges that must be addressed. At the same time, the future of IPM holds promising directions, propelled by technological advancements and a growing emphasis on ecological balance. Let's explore these challenges and prospective developments:

Challenges in IPM

1. **Adoption and Implementation:** Despite its benefits, the widespread adoption of IPM remains a challenge. Many farmers are either unaware of IPM practices or reluctant to transition from conventional pest control methods due to perceived risks, costs, or lack of knowledge.

2. **Economic Constraints:** The initial investment in IPM can be higher than traditional pest control methods. Farmers may face economic barriers in accessing the resources, technology, and training needed for effective IPM implementation.

3. **Knowledge and Training:** IPM requires a deep understanding of ecological systems, pest biology, and sustainable agricultural practices. There is a significant need for comprehensive training programs and educational resources for farmers and agricultural professionals.

4. **Complexity and Labor Intensity:** IPM strategies can be more complex and laborintensive than conventional methods. Regular monitoring, decisionmaking based on ecological assessments, and the use of diverse pest control methods require more time and effort.

5. **Climate Change and Pest Dynamics:** Changing climate patterns are altering pest populations and behaviors, making pest management more challenging. IPM strategies need to adapt to these changes, requiring ongoing research and flexibility in practices.

Future Directions in IPM

1. **Advancements in Technology:** The integration of technology such as precision agriculture, drones, remote sensing, and AI can enhance the efficiency and effectiveness of IPM. These technologies can assist in accurate pest monitoring, targeted interventions, and datadriven decisionmaking [31].

2. **Genetic Advances:** The development of pestresistant crop varieties through genetic engineering or traditional breeding can significantly reduce the reliance on chemical controls, aligning with IPM principles [32].

3. **ClimateAdaptive Strategies:** Developing IPM strategies that are resilient to climate change will be crucial. This includes understanding how changing weather patterns affect pest biology and ecology and adjusting management practices accordingly.

4. Policy and Incentives: Governments and agricultural bodies can play a significant role in promoting IPM by providing incentives, creating supportive policies, and investing in research and extension services.

5. Community and Ecosystem Approaches: Adopting a broader ecosystem approach and encouraging community participation can enhance IPM effectiveness. This involves considering the agricultural landscape as a whole and fostering collaboration among farmers, researchers, and policymakers [33-37].

6. Integrated Approach with Other Sustainable Practices: Combining IPM with other sustainable agricultural practices like organic farming, agroecology, and conservation agriculture can lead to more holistic and sustainable agricultural systems. The future of IPM lies in overcoming its current challenges through education, technological innovation, policy support, and adaptive strategies. As we move forward, IPM will continue to evolve, integrating new scientific insights and technologies. Embracing these changes and challenges will be key to ensuring that IPM remains at the forefront of sustainable agriculture, contributing to food security, environmental health, and economic viability for future generations [38-42]

Conclusion:

Integrated Pest Management (IPM) stands as a testament to the ingenuity and resilience of modern agriculture. Balancing the complexities of pest control with the imperatives of environmental stewardship and economic viability, IPM has emerged as a critical component in the pursuit of sustainable agriculture. As we reflect on the journey of IPM, from its genesis as a response to the limitations of conventional pest control to its current status as a multifaceted and dynamic field, several key takeaways crystallize. IPM exemplifies a paradigm shift from reactive to proactive pest management. By prioritizing ecological balance, embracing biological control methods, and implementing practices that are in harmony with nature, IPM goes beyond mere pest control. It represents a deeper understanding of and respect for the intricate interplay between agricultural practices and the natural environment.

The strength of IPM lies in its integrated approach, combining diverse methods and strategies from biological, cultural, mechanical, and chemical controls. This integration not only addresses the immediate challenges of pest management but also fosters long-term sustainability. It's a holistic approach that considers the entire ecosystem, aiming to minimize negative impacts while maximizing agricultural productivity. The road ahead for IPM is not without challenges. Overcoming barriers to adoption, addressing economic constraints, and continually adapting to changing pest dynamics and climate conditions remain critical areas of focus. However, the future is bright with the promise of technological advancements, policy support, and continued innovation. The integration of precision agriculture, genetic advancements, and climate-adaptive strategies are just a few examples of the potential paths forward for IPM. The successful implementation and evolution of IPM hinge on continuous education, awareness, and collaboration. Farmers, researchers, policymakers, and the broader community must work together to share knowledge, develop and refine practices, and promote the widespread adoption of IPM principles. In conclusion, Integrated Pest Management is not just a set of techniques; it is a philosophy and an essential tool for sustainable agriculture. It offers a path forward that respects and preserves our environmental resources while ensuring food security and economic sustainability. As we face the challenges of a growing global population and environmental uncertainties, IPM stands as a beacon of balanced, responsible, and forward-thinking agricultural practice, pivotal in shaping a sustainable future for generations to come.

References

1. Baker, B. P., Green, T. A., & Loker, A. J. (2020). Biological control and integrated pest management in organic and conventional systems. *Biological Control*, *140*, 104095.
2. Barzman, Marco, Paolo Bàrberi, A. Nicholas E. Birch, Piet Boonekamp, Silke Dachbrodt-Saaydeh, Benno Graf, Bernd Hommel et al. "Eight principles of integrated pest management." *Agronomy for sustainable development* 35 (2015): 1199-1215.
3. Paul, EA and Robertson, GP (1989) Ecology and the agricultural sciences: a false dichotomy? *Ecology* 70, 1594–1597.

4. Perrin, RM and Phillips, ML (1978) Some effects of mixed cropping on the population dynamics of insect pests. *Entomologia Experimentalis et Applicata* 24, 585–593.
5. Peshin, R and Dhawan, AK (eds) (2009 *a*) *Integrated Pest Management, Volume 1: Innovation-Development Process*. Dordrecht, The Netherlands: Springer.
6. Peshin, R and Dhawan, AK (eds) (2009 *b*) *Integrated Pest Management, Volume 2: Dissemination and Impact*. Dordrecht, The Netherlands: Springer.
7. Peshin, R, Jayaratne, KSU and Sharma, R (2014) IPM extension: a global overview. In Abrol, DP (ed.), *Integrated Pest Management: Current Concepts and Ecological Perspective*. San Diego, California: Academic Press, pp. 493–529.
8. Philips, CR, Rogers, MA and Kuhar, TP (2014) Understanding farmscapes and their potential for improving IPM programs. *Journal of Integrated Pest Management* 5, C1–C9. <http://dx.doi.org/10.1603/IPM13018>.
9. Pickett, CH and Bugg, RL (eds) (1998) *Enhancing Biological Control: Habitat Management to Promote Natural Enemies of Agricultural Pests*. Oakland, California: University of California Press.
10. Singh, Arvind Kumar, Neera Yadav, Ajai Singh, and Arpit Singh. "Transcription factors that regulate gene expression under drought." In *Acta Biology Forum*, vol. 2, pp. 01-04. 2023.
11. Pilson, D and Prendeville, HR (2004) Ecological effects of transgenic crops and the escape of transgenes into wild populations. *Annual Review of Ecology, Evolution, and Systematics* 35, 149–174.
12. Unnisa, S. A., Rao, B. B., & Vattikoti, P. (2022). Biochemical parameters of selected plants as air pollution indicators. *Acta Botanica Plantae*, 43-50.
13. Pimentel, D (1961) Species diversity and insect population outbreaks. *Annals of the Entomological Society of America* 54, 76–86.
14. Salam, M. A., Islam, M. R., Diba, S. F., & Hossain, M. M. (2019). Marker assisted foreground selection for identification of aromatic rice genotype to develop a modern aromatic line. *Plant Science Archives*
15. Touseef, M. (2023). Exploring the Complex underground social networks between Plants and Mycorrhizal Fungi known as the Wood Wide Web. *Plant Science Archives*. V08i01, 5.
16. Pimm, SL (1991) *The Balance of Nature? Ecological Issues in the Conservation of Species and Communities*. Chicago, Illinois: University of Chicago Press.

17. Polis, GA, Myers, CA and Holt, RD (1989) The ecology and evolution of intraguild predation: potential competitors that eat each other. *Annual Review of Ecology, Evolution, and Systematics* 20, 297–330.
18. Poveda, K, Gómez, MI and Martínez, E (2008) Diversification practices: their effect on pest regulation and production. *Revista Colombiana Entomología* 34, 131–144.
19. Power, AG (1999) Linking ecological sustainability and world food needs. *Environment, Development and Sustainability* 1, 185–196.
20. Nweze, C. C., & Muhammad, B. Y. (2023). Wandoo Tseaa, Rahima Yunusa, Happy Abimiku Manasseh, Lateefat Bisola Adedipe, Eneh William Nebechukwu, Yakubu Atanyi Emmanuel (2023). Comparative Biochemical Effects of Natural and Synthetic Pesticides on Preserved *Phaseolus vulgaris* in Male Albino Rats. *Acta Botanica Plantae. V02i01*, 01-10.
21. Power, AG (2010) Ecosystem services and agriculture: tradeoffs and synergies. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365, 2959–2971. Okunlola, A. I., Opeyemi, M. A., Adepoju, A. O., & Adekunle, V. A. J. (2016). Estimation of carbon stock of trees in urban parking lots of the Federal University OF Technology, Akure, Nigeria (Futa). *Plant Science Archives*
22. Pretty, J and Bharucha, ZP (2015) Integrated pest management for sustainable intensification of agriculture in Asia and Africa. *Insects* 6, 152–182.
23. Prokopy, RJ (1993) Stepwise progress toward IPM and sustainable agriculture. *IPM Practitioner* 15, 1–4.
24. Prokopy, RJ and Croft, BA (1994) Apple insect management. In Metcalf, RL and Luckmann, WH (eds), *Introduction to Insect Pest Management*, 3rd Edn. New York, New York: Wiley, pp. 543–589.
25. Mydeen, Abdul Kapur Mohamed, Nikhil Agnihotri, Raj Bahadur, Wankasaki Lytand, Neeraj Kumar, and Sanjay Hazarika. "Microbial Maestros: Unraveling the crucial role of microbes in shaping the Environment." In *Acta Biology Forum*, vol. 2, pp. 23-28. 2023.
26. Qiang, CZ, Kuek, SC, Dymond, A and Esselaar, S (2011) *Mobile Applications for Agriculture and Rural Development*. Washington, DC: ICT Sector Unit, World Bank.
27. Risch, SJ and Carroll, CR (1982) Effect of a keystone predaceous ant, *Solenopsis geminata*, on arthropods in a tropical agroecosystem. *Ecology* 63, 1979–1983.

28. Risch, SJ, Andow, D and Altieri, MA (1983) Agroecosystem diversity and pest control: data, tentative conclusions, and new research directions. *Environmental Entomology* 12, 625–629.
29. Corpuz, Mary Cris, Hazel R. Balan, and Nick C. Panares. "Biodiversity of benthic macroinvertebrates as bioindicator of water quality in Badiangon Spring, Gingoog City." (2016). *Plant Science Archives*
30. Nanda, Rupali, Fraz Ahmed, and Renu Sharma. "NishaBhagat and Kewal Kumar (2022). Ethnobotanical Studies on Some Angiosperms of Tehsil Hiranagar of District Kathua (Jammu and Kashmir), India." *Acta Botanica Plantae*: 01-11.
31. Rogers, EM (1983) *Diffusion of Innovations*, 4th Edn. New York, New York: The Free Press.
32. Romeis, J, Bartsch, D, Bigler, F, Candolfi, MP, Gielkens, MM, Hartley, SE, Hellmich, RL, Huesing, JE, Jepson, PC, Layton, R, Quemada, H, Raybould, A, Rose, RI, Schiemann, J, Sears, MK, Shelton, AM, Sweet, J, Vaituzis, Z and Wolt, JD (2008) Assessment of risk of insect-resistant transgenic crops to nontarget arthropods. *Nature Biotechnology* 26, 203–208.
33. Root, RB (1973) Organization of a plant-arthropod association in simple and diverse habitats: the fauna of collards (*Brassica oleracea*). *Ecological Monographs* 43, 95–124.
34. Ashokri, H. A. A., &Abuzririq, M. A. K. (2023). The impact of environmental awareness on personal carbon footprint values of biology department students, Faculty of Science, El-Mergib University, Al-Khums, Libya. In *Acta Biology Forum. V02i02* (Vol. 18, p. 22).
35. Rosenheim, JA, Kaya, HK, Ehler, LE, Marois, JJ and Jaffee, BA (1995) Intraguild predation among biological-control agents: theory and evidence. *Biological Control* 5, 303–335.
36. Savidge, JA (1987) Extinction of an island forest avifauna by an introduced snake. *Ecology* 68, 660–668.
37. Savolainen, V, Cowan, RS, Vogler, AP, Roderick, GK and Lane, R (2005) Towards writing the encyclopaedia of life: an introduction to DNA barcoding. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 360, 1805–1811.
38. Schoenly, KG and Barrion, AT (2016) Designing standardized and optimized surveys to assess invertebrate biodiversity in tropical irrigated rice using structured inventory and species richness models. *Environmental Entomology* 45, 446–464.

39. Schoenly, K and Cohen, JE (1991) Temporal variation in food web structure: 16 empirical cases. *Ecological Monographs* 61, 267–298.
40. Schoenly, KG, Cohen, JE, Heong, KL, Arida, GS, Barrion, AT and Litsinger, JA (1996 *a*) Quantifying the impact of insecticides on food web structure of rice-arthropod populations in a Philippine farmer's irrigated field: a case study. In Polis, GA and Winemiller, KO (eds), *Food Webs: Integration of Patterns and Dynamics*. Boston, MA: Springer, pp. 343–351.
41. Schoenly, K, Cohen, JE, Heong, KL, Litsinger, JA, Aquino, GB, Barrion, AT and Arida, G (1996 *b*) Food web dynamics of irrigated rice fields at five elevations in Luzon, Philippines. *Bulletin of Entomological Research* 86, 451–466.

UNDER PEER REVIEW