

Productivity of Dry direct seeded rice as influenced by herbicide combinations

Abstract

A field experiment was conducted during Kharif, 2022, in Institute of Rice Research, Rajendranagar, Hyderabad in Telangana state (country?). Fourteen weed management practices were evaluated in a randomized complete block design, replicated thrice. Weed management in Dry DSR can be done by adopting mechanical/ Hand weeding at 20, 40, 60 DAS under labour available conditions. Under labour scarcity conditions PE application of Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix) followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE weed followed by mechanical weeding at 40-45 DAS is the best option. Other suitable Pre emergence herbicides that can be used are Pyrazosulfuron ethyl 10% WP 20 g/ha, Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix) and Pendimethalin 38.7 % CS 677.25 g/ha. Post emergence application of Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix) and Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix) at 2-4 weed leaf stage.

Key words: Grain yield, Herbicide combinations, Productivity, dry direct seeded rice, Weed density

Introduction

Rice (*Oryza sativa* L.) is one of the most important food crops in the world, forms the staple diet of billions of people and provides 70% direct employment to the rural India. Rice production symbolizes the single largest land use for food production on earth and worldwide, rice is grown on 164 M ha with an annual production of about 750Mt of paddy (FAO, 2019). To meet the global rice demand, it is estimated that about 114Mt of additional milled rice needs to be produced by 2035 (Singh *et al.*, 2015) and therefore, meeting ever increasing rice demand in a sustainable way with shrinking natural resources is a great challenge.

India is the second largest producer of Rice after China. The area production and productivity of Rice in India are 44.79 million hectares, 112.41 million tones, 2578 kg/ha respectively. In India, major rice growing states are West Bengal, Punjab, Uttar Pradesh, Tamil Nadu, Andhra Pradesh, Telangana, Bihar, Odisha etc. The area and production of Rice in Telangana is 2 million hectares and 6.6 million tons respectively. [Reference](#)

In the traditional method of establishment of rice, both puddling and transplanting operations requires a large amount of labour (Chauhan and Yadav, 2013). The increasing demand for labour in non agricultural sectors and increasing labour costs resulting from the migration of rural labour to the cities, creates labour shortage at the critical time of transplanting (Chauhan, 2012). Late commencement of monsoon showers and drudgery involved in puddling and manual transplanting delay rice transplanting (Farooq *et al.*, 2006). There is a need to look for direct seeding of rice that not only cope up with existing limitations but also are affordable and economically viable and secure future food demand.

Most significant biological constraint in DSR is weed infestation, which is very severe when compared to transplanted rice. The simultaneous emergence of crop and weed also poses a severe loss and weed control becomes a cumbersome procedure. In dry seeded rice, the critical period of weed competition has been reported to be 15-60 days after

seeding (Chauhan and Mahajan, 2014) and if weeds are controlled by any means during this period, minimal yield losses can be observed. Weed management using herbicides has become an integral part of modern agriculture. Herbicides have greater flexibility of operation and are often cost effective compared to any other method of weed management. **Reference**A number of herbicides at low doses are giving effective weed control. The present investigation was carried out to find out the best possible pre and post emergence herbicides for weed management in Dry direct seeded rice.

Material Methods

The investigation was carried out during *kharif* 2022 at Institute of Rice Research, Professor Jayashankar State Agricultural university, Rajendranagar, Hyderabad, situated at an altitude of 542.3 m above MSL at 17°19' N latitude and 78°23' E longitude. It is in the Southern Telangana agro-climatic zone of Telangana state. According to Troll's climatic classification, it falls under semi-arid tropics (SAT). During the cropping period rainfall of 861.1 mm was received. RNR 29325(Rajendranagar 5) is short duration rice variety matures in 125 days. It has long slender grains and moderate resistance to BPH and leaf blast. Experiment consist of 14 treatments consisting of pre emergence application of Pendimethalin 38.7 % CS 677.25 g/ha, Pyrazosulfuron ethyl 10% WP 20 g/ha and Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix) followed by post emergence application of Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix), Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix), Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) and Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix), followed by mechanical weeding at 40-45 DAS in herbicide treated plots, weed free treatment i.e., hand weeding twice at 25 and 45 DAT and weedy check in RBD replicated thrice. **Description of the measurements that was done to evaluate the results of this treatment ?.**

Weed density (No. m⁻²)

Weed density was taken at 20, 40 and 60 DAS in randomly selected quadrat (0.5 x 0.5 m²) in each plot. At every sampling, three categories of weeds viz., grasses, sedges and broad leaved weeds were separated and expressed as number m⁻². The sampling was done outside the net plot but within the gross plot.

Results and discussion

Effect on weed density

Grass weed density (No.m⁻²) at 20 DAS

At 20 DAS (Table1) the significantly lower grass weed density was observed with T₁₃: Mechanical/ hand weeding at 20, 40, 60 DAS (2.8), it was statistically on par with T₉- pendimethalin 38.4% + pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by triafamone 20% + ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (3.1), T₁- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (3.2) it was inturn on par with T₃- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE

followed by mechanical weeding at 40-45 DAS (3.5), T₆- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(3.6) and was further followed by the treatment T₈ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (3.7), T₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(3.8), T₁₀ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (4.0) and it was inturn on par with T₂- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(4.2), T₇ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS(4.2) T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS (4.2), T₄- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (4.4) it was followed by T₅ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (5.1) and significantly highest grass weed density was recorded with T₁₄- Unweeded control (5.4).(Table1)**Reference that gives support to the observed results ?**

Sedges weed density (No.m⁻²) at 20 DAS

At 20 DAT (table 1), the lower sedge weed density was observed with T₉- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (1.4), it was on par with T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS(1.7), it was inturn on par with T₁- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS(1.8) and it was inturn on par with T₃ Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (2.0), T₆- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(2.0), T₇ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS(2.0), T₈ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD

25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (2.1), T₁₄- Unweeded control (2.1).Further followed by T₂- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(2.2),T₄- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (2.2), T₅- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (2.2), T₁₀ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (2.2), T₁₃: Mechanical/ Hand weeding at 20, 40, 60 DAS (2.3) and significantly higher sedge weed density recorded with T₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(2.6). (Table1) **Reference that gives support to the observed results ?**

Broad leaved weed density (No.m⁻²) at 20 DAS

At 20 DAS (Table 1), the lower broad leaved weed density was observed with T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS(2.0) it was statistically on par with T₁₃- Mechanical/ Hand weeding at 20, 40, 60 DAS (2.6), it was inturn on par with all the treatments in the order of T₉- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (2.8),T₇ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS(2.8),T₁- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS(3.0),T₁₀ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (3.0),T₆- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(3.2), T₂- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(3.5),T₃ Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (3.5), T₄- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (3.5),T₅- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS

(3.8), T8 - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (3.8), T₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(3.9) except T₁₄- Unweeded control (5.8) which recorded significantly highest broad leaved weed density. (Table1) **Reference that gives support to the observed results ?**

Grass weed density(No.m⁻²) at 40DAS

At 40 DAS (Table 2), significantly lower grass weed density was observed with T₁₃- Mechanical/ Hand weeding at 20, 40, 60 DAS (4.2) and was at par with T₁- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS(1.4), it was followed by T₉- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (2.0), T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS(2.0) it was statistically on par with T₃- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (2.2), T₅- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (2.2), T₄- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (2.3) and T₇ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS(2.1) it was further followed by T₂- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(2.4) and was on par with T₆- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(2.5), T₈ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (2.4), T₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(2.4), and T₁₀ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (2.6). The significantly highest grass weed density was observed with T₁₄- Unweeded control (5.5). **Reference that gives support to the observed results ?**

Sedge weed density(No.m⁻²) at 40DAS

At 40 DAS (Table 2), significantly lower sedge weed density was with T₁₃- Mechanical/ Hand weeding at 20, 40, 60 DAS (1.9) it was statistically on par with T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS(2.1),T₅- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (2.2), T₇ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS(2.2), T₉- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (2.3), T₃- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (2.4), T₂- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(2.4) and T₆- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(2.4) it was followed by T₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(2.6), T₁₀ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (2.7), T₄- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (2.8), T₁- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS(3.0), T₈ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (3.2), The significantly highest sedge weed density was observed with T₁₄- Unweeded control (6.0). **Reference that gives support to the observed results ?**

Broad leaved weed density(No.m⁻²) at 40 DAS

At 40 DAS (Table 2), significantly lower density of broad leaved weed density was observed with T₁₃- Mechanical/ Hand weeding at 20, 40, 60 DAS (1.7) it was statistically on par with T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS(1.7) it was inturn on par with T₅- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (2.0), T₆- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(2.0), T₂- Pendimethalin 38.7 % CS 677.25

g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(2.1), T3- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (2.3) and T₄- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (2.4), T₉- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (2.3), T₁₀ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (2.3) and T₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(2.3), it was followed byT₁- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS(2.6), T₇ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS(2.7) andT₈ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (3.0), Significantly highest sedge weed density was observed with T₁₄- Unweeded control (7.3). **Reference that gives support to the observed results ?**

Grass weed density(No.m⁻²) at 60DAS

At 60 DAS (Table 3), significantly lower grass weed density was observed with T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS(1.6), it was due to effective reduction in weed density by pendimethalin + Pyrazosulfuron ethyl during initial period and by POE and mechanical weeding at later periods(Pavithra *et al.* , 2021). it was statistically on par with T₁₃- Mechanical/ Hand weeding at 20, 40, 60 DAS (1.7), T₉- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (1.7),T₉- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS T₂- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(2.1) and T₅- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (2.1) it was followed by T₆- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(2.9) it was statistically on par with T₇ - Pyrazosulfuron

ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS(3.0), T3- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (3.0)andT4- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (3.6), further followed by T₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(3.9),T₁₀- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (4.3) andT1- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS(4.4) which was inturn on par withT8 - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (5.0). Significantly highest grass weed density was observed with T14- Unweeded control (6.8). **Reference that gives support to the observed results ?**

Sedge weed density(No.m⁻²) at 60DAS

At 60 DAS (Table 3), significantly lower sedge weed density was observed with T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS(1.6) and T₁₃- Mechanical/ Hand weeding at 20, 40, 60 DAS (1.6) it was statistically on par with T₂- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(1.7), T9- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (1.8), T5- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (2.0) and T3- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (2.1) it was followed byT6- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop - butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(2.5), T7 - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS(2.7)andT₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(2.8)further followed by T₄- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (3.3) andT1-

Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS(3.6) it was followed by T₁₀ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (4.3)and followed by T₈ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (5.1). the significantly highest total sedge weed density was observed with T₁₄- Unweeded control (8.8).**Reference that gives support to the observed results ?**

Broad leaved weed density(No.m⁻²) at 60 DAS

At 60 DAS (Table 3) the significantly lower broad leaved weed density was observed with T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS(1.7) it was statistically on par with T₁₃- Mechanical/ Hand weeding at 20, 40, 60 DAS (1.8), T₉- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (1.8) and T₅- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (2.0) and was followed by T₂- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(3.0), T₃- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (3.6), T₁₀ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (3.6) it was inturn on par with T₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(3.9),T₇ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS(3.9), T₆- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(4.0) ,T₄- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (4.9) which was inturn on par with T₁- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS(5.0) and was followed by T₈ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (6.2) and T₁₄- Unweeded control (6.3).**Reference that gives support to the observed results ?**

Table 1: Weed density (No/m²) at 20 DAS as influenced by different weed management practices in Dry direct seeded rice during kharif, 2022

Treatments	Weed density (No/m ²) 20 DAS		
	Grass	Sedge	BLW
T1:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	3.2 (9.0)	1.8 (2.3)	3.0 (8.0)
T2:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS	4.2 16.3	2.2 (3.7)	3.5 (11.0)
T3:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	3.5 (11.3)	2.0 (3.0)	3.5 (11.0)
T4:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	4.4 (18.0)	2.2 (4.0)	3.5 (11.3)
T5:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	5.1 (25.3)	2.2 (4.0)	3.8 (13.3)
T6:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40-45 DAS	3.6 (11.7)	2.0 (3.0)	3.2 (9.0)
T7:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	4.2 (16.3)	2.0 (3.0)	2.8 (6.7)
T8:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	3.7 (13.0)	2.1 (3.3)	3.8 (13.7)
T9:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	3.1 (8.7)	1.4 (1.0)	2.8 (7.0)
T10:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	4.0 (15.0)	2.2 (4.0)	3.0 (8.0)
T11:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS	4.2 (17.0)	1.7 (2.0)	2.0 (3.0)

T12:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	3.8 (13.3)	2.6 (5.7)	3.9 (14.0)
T13:Mechanical/ Hand weeding at 20, 40, 60 DAS	2.8 (6.7)	2.3 (4.3)	2.6 (5.7)
T14:Unweeded control	5.4 (28.3)	2.1 (2.3)	5.8 (33.0)
S. Em±	0.15	0.11	0.78
CD (P=0.05)	0.4	0.3	2.3

Original data in parentheses was subjected to square root transformation

Table 2: Weed density (No/m²) at 40 DAS as influenced by different weed management practices in Dry direct seeded rice during kharif, 2022

Treatments	Weed density (No/m ²) 40 DAS		
	Grass	Sedge	BLW
T1:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	2.7 (6.3)	3.0 (8.3)	2.6 (5.7)
T2:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS	2.4 (4.7)	2.4 (4.7)	2.1 (3.7)
T3:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	2.2 (4.0)	2.4 (4.7)	2.3 (4.3)
T4:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	2.3 (4.3)	2.8 (6.7)	2.4 (5.0)
T5:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	2.2 (4.0)	2.2 (3.7)	2.0 (3.0)
T6:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40-45 DAS	2.5 (5.3)	2.4 (4.7)	2.0 (3.0)
T7:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	2.1 (3.7)	2.2 (3.7)	2.7 (6.3)
T8:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	2.4 (5.0)	3.2 (9.0)	3.0 (8.0)
T9:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% +	2.0 (3.3)	2.3 (4.3)	2.3 (4.7)

Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS			
T10:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	2.6 (6.0)	2.7 (6.3)	2.3 (4.3)
T11:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS	2.0 (3.0)	2.1 (3.7)	1.7 (2.0)
T12:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	2.4 (5.0)	2.6 (5.7)	2.3 (4.3)
T13:Mechanical/ Hand weeding at 20, 40, 60 DAS	1.4 (1.0)	1.9 (2.7)	1.7 (2.0)
T14:Unweeded control	5.5 (29.0)	6.0 (35.7)	7.3 (51.7)
S. Em±	0.1	0.2	0.23
CD (P=0.05)	0.3	0.6	0.7

Original data in parentheses was subjected to square root transformation

Table 3: Weed density (No/m²) at 60 DAS as influenced by different weed management practices in Dry direct seeded rice during kharif, 2022

Treatments	Weed density (No/m ²) 60 DAS		
	Grass	Sedge	BLW
T1:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	4.4 (18.7)	3.6 (12.0)	5.0 (24.3)
T2:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS	2.1 (3.7)	1.7 (2.0)	3.0 (8.3)
T3:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	3.0 (8.0)	2.1 (3.7)	3.6 (12.0)
T4:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	3.6 (12.3)	3.3 (10.3)	4.9 (23.0)
T5:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	2.1 (3.3)	2.0 (3.0)	2.0 (3.0)
T6:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40-45 DAS	2.9 (7.7)	2.5 (5.7)	4.0 (15.3)

T7:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	3.0 (8.3)	2.7 (6.7)	3.9 (14.7)
T8:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	5.0 (24.0)	5.1 (25.3)	6.2 (37.0)
T9:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	1.7 (2.0)	1.8 (2.3)	1.8 (2.3)
T10:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	4.3 (17.7)	4.3 (17.7)	3.6 (12.0)
T11:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS	1.6 (1.7)	1.6 (1.7)	1.7 (2.0)
T12:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	3.9 (14.3)	2.8 (7.0)	3.9 (14.7)
T13:Mechanical/ Hand weeding at 20, 40, 60 DAS	1.7 (2.0)	1.6 (1.7)	1.8 (2.3)
T14:Unweeded control	6.2 (37.7)	8.8 (76.0)	6.3 (38.7)
S. Em±	0.24	0.2	0.28
CD (P=0.05)	0.70	0.6	0.8

Original data in parentheses was subjected to square root transformation

Plant height (cm) at harvest

Data pertaining to plant height (Table 4) under different weed management practices was analysed statistically and presented in table 3. There was no significance difference in plant height was observed with different weed management practices. **Reference that gives support to the observed results ?**

Tillers /m² at harvest

Significantly higher number of tillers (Table 4) at harvest was observed with T₁₃- Mechanical/ Hand weeding at 20, 40, 60 DAS (364) it was statistically on par with T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS(92.5), T₅- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (332), T₉- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha

(Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (329) it was followed by T₂- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(325), T₃- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (317), T₆- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(308), T₈ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (305),T₁₀ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (302),T₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(298), T₄- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (298) it was followed by T₇ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS(274) it is inturn on par with T₁- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS(255). Significantly lower number of tillers/m² was recorded with T₁₄- Unweeded control (233). **Reference that gives support to the observed results ?**

Panicles /m² at harvest

Significantly highest number of panicles/ m² at harvest (Table 4) were observed with T₁₃- Mechanical/ Hand weeding at 20, 40, 60 DAS (327) it was statistically on par with T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS(319), T₅- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (314),T₉- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (307), T₂- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(300) and T₃- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (293), it was followed by T₆- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(279),T₈ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium

10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (305)T₄- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (273), T₁₀ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (272), T₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(264), T₇ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS(250) it was inturn on par with T₁- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS(247), significantly lowest number of panicles/m² recorded in T₁₄- Unweeded control (209).

Reference that gives support to the observed results ?

UNDER PEER REVIEW

Grain yield

The significantly higher grain yield was observed (Table 4) with T₁₃- Mechanical/ Hand weeding at 20, 40, 60 DAS (3959) it was statistically on par with T₁₁ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS(3795), T₅- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (3609),T₉- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40 -45 DAS (3570), T₂- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS(3568) and T₃- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (3502), This was mainly due to killing of germinating weed seeds as well as removal of established weeds, During the critical period of crop weed competition, weed density continuously remained increasing and crop growth affected due to competition from weeds for the natural resources (Spandana et al., 2017),it was followed byT₆- Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40 -45 DAS(3463), T₇ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS(3400), T₁₂- Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40- 45 DAS(3355),T₄- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS (3285), T₁₀ - Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop -butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS (3167), T₁- Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS(3114),T₈ - Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40 -45 DAS (3068). Significantly lower grain yield was observed with T₁₄- Unweeded control (1419). **Reference that gives support to the observed results ?**

Table4: Growth and yield attributes influenced by different weed management practices in Dry direct seeded rice during kharif, 2022

Treatments	Plant height (cm) at harvest	Tillers/m ² at harvest	Panicles/m ² at harvest	Grain Yield (kg/ha)
T1:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	79.9	255	247	3114
T2:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix)PoE at 2-4 leaf stage of the weed followed by mechanical weeding at 40-45 DAS	79.5	325	300	3568
T3:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	80.8	317	293	3502
T4:Pendimethalin 38.7 % CS 677.25 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	76.4	298	273	3285
T5:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	75.7	332	314	3609
T6:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 h/ha (Ready mix) PoE followed by mechanical weeding at 40-45 DAS	79.0	308	279	3463
T7:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75 g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	74.3	274	250	3400
T8:Pyrazosulfuron ethyl 10% WP 20 g/ha PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	79.9	305	275	3068
T9:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Triafamone 20% +	77.7	329	307	3570

Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS				
T10:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 h/ha (Ready mix)PoE followed by mechanical weeding at 40-45 DAS	82.0	302	272	3167
T11:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix)PE followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS	77.6	355	319	3795
T12:Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha(Ready mix) PE followed by Bispyribac Sodium 10% SC 25 g/ha + Penoxsulam 2.67 % OD 25g/ha (Tank mix) PoE followed by mechanical weeding at 40-45 DAS	77.5	298	264	3355
T13:Mechanical/ Hand weeding at 20, 40, 60 DAS	76.6	364	327	3959
T14:Unweeded control	75.6	233	209	1419
S. Em+	1.98	13.0	15.2	164.1
CD (P=0.05)	NS	38.0	44	480

Conclusion

Weed management in Dry DSR can be done by adopting mechanical/ Hand weeding at 20, 40, 60 DAS under labour available conditions. Under labour scarcity conditions PE application of Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix) followed by Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP 4 g/ha + Cyhalofop Butyl 10% EC 75g/ha (Tank mix)PoE weed followed by mechanical weeding at 40-45 DAS is the best option. Other suitable Pre emergence herbicides that can be used are Pyrazosulfuron ethyl 10% WP 20 g/ha, Pendimethalin 38.4% + Pyrazosulfuron ethyl 0.85% ZC 900 + 20 g/ha (Ready mix) and Pendimethalin 38.7 % CS 677.25 g/ha. Post emergence application of Triafamone 20% + Ethoxysulfuron 10% WG 44+22.5 g/ha (Ready mix) and Penoxsulam 1.02 % + Cyhalofop-butyl 5.1% OD 135 g/ha (Ready mix) at 2-4 weed leaf stage.

References

- Chauhan, B.S and Mahajan, G. eds. 2014. Recent advances in weed management. Springer New York.
- Chauhan, B.S.2012. Weed ecology and weed management strategies for dry-seeded rice in Asia. Weed Technology.26(1):1-13.

- Chauhan, B.S and Yadav, A.2013. Weed management approaches for dry-seeded rice in India: a review. *Indian Journal of Weed Science*.45(1):1-6.
- Farooq, M., Barsa, S.M and Wahid, A.2006. Priming of field-sown rice seed enhances germination, seedling establishment, allometry and yield. *Plant growth regulation*.49(2):285-294.
- Food and agriculture organization of the United Nations.2019. *Rice Market Monitor* 16(1). Trade and <http://www.fao.org/economic/est/publications/ricepublications/MarketsDivision>,
- Singh, M., Bhullar, M.S and Chauhan, B.S.2015. Influence of tillage, cover cropping and herbicides on weeds and productivity of dry direct-seeded rice. *Soil and Tillage Research*.147:39-49.
- Pavithra, R. Poonguzhalan, A.L. Narayanan and P. Saravanane 2021 Weed management in aerobic rice with sequential application of pendimethalin and bispyribac-sodium under coastal deltaic ecosystem. *Indian Journal of Weed Science* 53(1): 85–87.
- Spandana Bhatt.P, M. Yakadri, M. Madhavi, S. Sridevi and P. Leela Rani 2017 Productivity of transplanted rice as influenced by herbicide combinations. *Indian Journal of Weed Science* 49(2): 128–131.