

Assessment of different okra (*Abelmoschus esculentus* L.) varieties for yield and biomass production

Proposed title:

Evaluation of different okra (*Abelmoschus esculentus* L.) cultivars for yield and biomass production

Abstract

In a field experiment conducted at G. P. Koirala College of Agriculture and Research Center, Morang, Nepal, spanning from April to August 2022. ~~The study was conducted to~~ evaluate the yield and biomass production of different okra (*Abelmoschus esculentus* L.) varieties. ~~Employing a randomized complete block design with three replications, the study included varieties~~ such as ArkaAnamika, Chandani, Chiranjeevi F1, F1 Glory, Gunjan, JK1666, OH-940, Punjab Selection, R35 Selection, and Swastik-2. Results revealed significant variability among the varieties, with F1 Glory (0.91 Kg/m²) and Punjab Selection (0.96 Kg/m²) emerging as top performers in average yield per m². Conversely, Arka Anamika, Gunjan, and JK1666 exhibited the lowest yields. Notably, Punjab Selection (4.17 Kg/m²), OH-940 (4.28 Kg/m²), and Swastik-2 (3.86 Kg/m²) showcased the highest fresh biomass, while Gunjan had the lowest biomass. F1 Glory demonstrated the highest harvest index (1.14), followed by Chandani and Punjab Selection (both 0.88). These findings emphasize the substantial influence of varietal selection on okra productivity, highlighting the potential of Punjab Selection and F1 Glory for further exploration in vegetable cultivation and biomass production. The study provides valuable insights for optimizing okra cultivation strategies in Nepal and similar agro-ecological contexts.

Keywords

Abelmoschus esculentus, Okra biomass, Fresh plant weight, Okra performance, Okra Varieties

Introduction

The fibrous pods of okra (*Abelmoschus esculentus* L.), commonly referred to as "lady's finger," make it one of the most significant and nutrient-dense vegetables widely cultivated in tropical and warm temperate ~~parts~~ areas of the world (Jonah et al., 2019). Biomass is the leftover residue of agricultural plants like roots, shoots, and stems (Freidenreich et al., 2022; Lee et al., 2020). Biomass is classified into four types; they are woody biomass (it consists of wood plants and trees), non-woody biomass (plants parts like stems, roots, and leaves), process waste biomass (waste from various typed of agroindustry), and processed fuel biomass (producer gas, charcoal, etc.) (Verma et al., 2017). The biomass of the plant is determined by weighing the overall part of the plant individually and the root and shoot separately of an individual plant on the same day (Brocks & Bareth, 2018; Nath et al., 2021). ~~The maximum~~ Biomass can be highly obtained from the area where the leguminous crop ~~was~~ planted in the last planting season (Maclaren et al., 2019). ~~A measure of energy produced from~~ Green biomass technology is the most effective way to decrease the emission of greenhouse gases to meet the increasing energy demand and to reduce the problem of global warming (Antar et al., 2021).

Okra is ~~geographically African in origin which falls under the Malvaceae family and~~ is considered a rich source of vitamins, carbohydrate, minerals, proteins, and unsaturated fats (Sousa et al., 2020). It is an annual warm-season plant grown under a pH of 5 to 8, mainly for pods (Moosavi et al., 2018). Further, biomass production in okra is highly affected by the various constituents of soil, as these components are directly related to the plant's growth. The production of the biomass of okra varies with the application of fertilizers during cultivation. Higher integration of nitrogenous fertilizer results in higher biomass production as it increases the composition of dry matter and vice versa. ~~The~~ Maximum yield of pods and biomass can be obtained through the integrated application of fertilizers ~~more than recommendation ed chemical fertilizers~~ (Jonah & Kwaga, 2019). The biomass of okra can be used to synthesize fibers as it is biodegradable and eco-friendly compared to synthetic fibers (Gupta et al., 2021; Singha & Guleria, 2015; Verma et al., 2017). ~~As it is eco-friendly and solves the problems that arise from synthetic fibers,~~ Natural textile fibers are highly popular and have huge demand (Gupta et al., 2021; Singha & Guleria, 2015). Lignocellulosic fibers can be obtained from the stem of okra ~~through the~~ process known as retting, where fibers are separated from cementing and gummy components of okra. About 70% of α -cellulose fibers are present in biomass (stem), whose quality is similar to that of jute and can be used for the production of nets, various types of fabrics, carpets, etc. (Gupta et al., 2021). Second-generation fuels can be obtained from okra, which helps to decrease dependence on ~~the~~ nonrenewable sources or can be used for animal feed (Lee et al., 2020; Vadenbo et al., 2017). Cellulose in okra can be used as an adsorbent for wastewater treatment. Also, it can remove various types of ions and metals present in an aqueous solution. The biomass of okra is composed of hemicellulose (20 to 40 %), cellulose (55 to 65%), and lignin (15

Comment [Ma1]: Abstract needs improvement. Add the why the author design this study/ need of the research and also include conclusion or suggestions

Comment [Ma2]: Please give complete information first... then use abbreviation

Comment [Ma3]: This is a vague sentence please select the appropriate words.

Comment [Ma4]: Add the conclusion/suggestion which is applicable at the farmer levels

Comment [Ma5]: This section need rearrangement of the sentences for cohesiveness.

Comment [Ma6]: Shift this information in first paragraph

Comment [Ma7]: Repeated this information the is already provide in first paragraph

Comment [Ma8]: This is a vague sentence the author talking about fuels and suddenly add the information about livestock. Please split these two statements and also write in a better way.

to 25%). It consists of fibers obtained from stem parts (Singha & Guleria, 2015). Dried biomass is converted into powder form for a bed dryer. For the production of bioenergy, biomass is completely dried before cofiring for the improvement of the quality of bioenergy (Verma et al., 2017). Bio-flocculants produced from the biomass of okra are a must-suitable and eco-friendly alternative to chemical flocculants (Lee et al., 2018).

Biomass production emerges as a versatile solution to tackle diverse energy-related challenges, facilitating the generation of various types of energy. Despite its inherent benefits after production, the cultivation phase introduces challenges affecting arable land usage, competition with food and fiber crops, uneven soil distribution, nutrient depletion, weed proliferation, environmental impacts, and farmers' predominant focus on marketable pods. Unfortunately, the potential utilization of leftover plant biomass is often overlooked. This study uniquely centers on the exclusive assessment of biomass production and yield in different okra varieties. Given the limited existing research on okra varieties that simultaneously yield the highest biomass and yield, there exists a notable gap in knowledge on this topic. To bridge this gap and address challenges encountered during cultivation, including those related to environmental factors and farmers' practices, the research places a specific emphasis on evaluating ten okra varieties sourced from both domestic and foreign origins. This research aims to contribute valuable knowledge to an underexplored area, shedding light on okra varieties that can potentially maximize both yield and biomass production.

Materials and Methods

Site description

The experiment was conducted in the research field at G. P. Koirala College of Agriculture and Research Center located at Sundarharai, Morang, Nepal, to evaluate the yield and biomass of different okra varieties from April 2022 to August 2022. The geographical coordinates of this location are approximately 26° 40' 49.3" North latitude and 87° 21' 16.1" East longitude, and it sits at an elevation of 150 meters above sea level. The climate in this region is classified as tropical, characterized by an average annual temperature ranging from 25°C to 35°C. The soil characteristics of the experimental site were analyzed in qualitative measures with the help of a soil test kit box (Table 1).

Table 1: Soil characteristics of the experimental site

S. N.	Soil constituents	Properties
1	Soil pH	6.5
2	Soil texture	Silty loam
3	Nitrogen	Medium
4	Potassium	Medium
5	Phosphorus	Low
6	Surface soil wetness	0.648
7	Profile soil moisture	0.606
8	Root zone soil moisture	0.628

Meteorological data

~~The climate of the experimental area is tropical, with an average annual temperature ranging between 25 to 35 °C.~~ The average maximum temperature was 33.46 °C, and the minimum temperature was 22.86 °C. The total precipitation during the experimental period amounted to 1254.20 mm. The meteorological data during the whole experimental period is depicted in Figure 1.

Comment [Ma9]: Rearrange this information. Please arrange these lines in a meaning full ways

Formatted: Strikethrough

Comment [Ma10]: Already mention above

Comment [Ma11]: Relative humidity is present in the figure but missing in the text. Please add the information




Figure 1. Observational meteorological data throughout the growing period of okra in Gothgaun, Morang.

Cultural practices

The field experiment followed a Randomized Complete Block Design (RCBD) and involved ten okra varieties i.e., Arka Anamika, Chandani, Chiranjeevi F1, F1 Glory, Gunjan, JK 1666, OH-940, Punjab Selection, R35 Selection, and Swastik-2 with three replications. The ten different okra varieties in the experiment were Arka Anamika, Chandani, Chiranjeevi F1, F1 Glory, Gunjan, JK 1666, OH-940, Punjab Selection, R35 Selection, and Swastik-2. Each individual plot covered an area of nine square meters (3m x 3m), resulting in a total experimental area of 270 m². In each plot, there were 40 plants with a spacing of 0.6 m between rows and 0.35 m between individual plants. Fertilizer was applied at a rate of 120:60:60 kg NPK per hectare, and FYM was applied at a rate of 25 t/ha. The total amount of phosphorus (P) and potassium (K), as well as half of the nitrogen (N) fertilizer, were applied at the time of sowing. The remaining nitrogen doses were split into two applications at 22 and 40 days after sowing (DAS). Weeding was performed manually two times at 20 and 42 DAS. Irrigation was carried out in every two days in response to the soil moisture content.

Data collection

The data collection involved randomly selecting 12 plants from each plot, and data were collected from the first harvest to the last harvest. Harvesting was conducted based on maturity criteria, where okra pods were harvested when they reached a specific size of 12 cm or exhibited a bright green, fleshy appearance according to standard agricultural practices. Yield data (kg/m²) was recorded at weekly intervals throughout the growth period. Additionally, fresh biomass data, encompassing stem, shoots, roots, and leaves, were recorded collected at the end of the plant's growth period in August. The yield per m² was calculated by aggregating individual plant yields within each plot and then normalizing it to the plot area.

Statistical analysis

The obtained data on various parameters was subjected to statistical analysis using Rstudio (4.2.2 version). The significance of the results was assessed through an analysis of variance (ANOVA). Data entry was performed using MS Excel (2019), and the same software was utilized to create graphs and bar charts for the parameters under investigation. To compare the means of various parameters, Duncan's Multiple Range Test (DMRT) was employed, with a significance level of 5% (Gomez & Gomez, 1984).

Results and Discussion

Yield of different okra varieties

The experimental findings underscored the substantial variability in okra yield per m² across diverse varieties at distinct days after sowing (DAS), as detailed in Table 1. Statistically, the observed variations in each DAS were significant. These findings accentuate the temporal dynamics and significant varietal influences on okra yield, providing valuable insights for agronomic optimization and varietal selection. These findings indicate the dynamic growth patterns in okra crop output over time.

Notably, the overall mean at 45 DAS stood at 0.09 kg/m², exhibiting a subsequent progressive increase, peaking at 1.19 kg/m² by 75 DAS before exhibiting a decline. Chandani (1.13 Kg), Punjab selection (1.13 Kg), and Swastik-2 (1.11 Kg) manifested the highest yields per m², succeeded by F1 Glory (0.84 Kg) and OH-940 (0.82 Kg). In contrast, Gunjan and JK 1666 displayed the lowest yields (0.20 Kg/m²). Statistically, the observed

Comment [Ma12]: Give the reference here.

Comment [Ma13]: Great excellent. Please add the package.

Comment [Ma14]: I suggest the author start discussion with a separate heading because the author does not discuss their findings in here.

Comment [Ma15]: Discussion section need hard work. It should be improved. Discuss the findings with the prior studies. Here the author discuss the findings with single reported studies. Please try to include more information regarding your research

Formatted: Font: 10 pt, Bold

Formatted: Font: Bold

Formatted: Indent: First line: 0"

variations in each DAS were significant. These findings accentuate the temporal dynamics and significant varietal influences on okra yield, providing valuable insights for agronomic optimization and varietal selection. These findings indicate the dynamic growth patterns in okra crop output over time.

Comment [Ma16]: Rearrange these lines

Table 1: Yield per m² of different varieties of okra on various days after sowing

Varieties	45 DAS	60 DAS	75 DAS	90 DAS	105 DAS	Average
Arka Anamika	0.04 ^{de}	0.36 ^d	0.82 ^c	0.58 ^{cde}	0.45 ^{bcd}	0.45 ^d
Chandani	0.09 ^{bcd}	0.53 ^{cd}	1.17 ^{bc}	1.12 ^{ab}	1.13 ^a	0.80 ^{ab}
Chiranjeevi F1	0.01 ^e	0.56 ^{bcd}	0.97 ^{bc}	0.72 ^{bcd}	0.33 ^{cd}	0.52 ^{cd}
F1 Glory	0.12 ^b	0.92 ^{ab}	1.66 ^a	1.03 ^{ab}	0.84 ^{ab}	0.91 ^a
Gunjan	0.20 ^a	0.64 ^{abcd}	1.00 ^{bc}	0.39 ^e	0.20 ^d	0.49 ^{cd}
JK 1666	0.09 ^{bcd}	0.76 ^{abc}	1.15 ^{bc}	0.55 ^{de}	0.20 ^d	0.55 ^{bcd}
OH-940	0.11 ^{bc}	0.69 ^{abcd}	1.32 ^{ab}	0.96 ^{abcd}	0.82 ^{ab}	0.78 ^{ab}
Punjab Selection	0.05 ^{cde}	0.97 ^a	1.32 ^{ab}	1.31 ^a	1.13 ^a	0.96 ^a
R35 Selection	0.11 ^{bc}	0.57 ^{bcd}	1.24 ^{abc}	0.99 ^{abc}	0.72 ^{abc}	0.73 ^{abc}
Swastik-2	0.09 ^{bcd}	0.57 ^{bcd}	1.30 ^{ab}	1.11 ^{ab}	1.11 ^a	0.84 ^a
Mean	0.09	0.66	1.19	0.88	0.69	0.70
SEM±	0.010	0.047	0.055	0.064	0.076	0.039
CV (%)	32.54	30.00	20.18	25.83	35.46	19.66
LSD	0.05	0.34	0.41	0.39	0.42	0.23
F-Test	***	*	*	**	***	**

*Significant at 5% level of significance, **Significant at 1% level of significance, ***Significant at 0.1% level of significance, CV: Coefficient of variance, LSD: Least Significant Difference, SEM: Standard Error of Mean

Formatted: Els-body-text, Left, Indent: First line: 0.2"

Comment [Ma17]: Please cross check the table numbering

Formatted: Font: Bold, English (United States)

Fresh biomass and harvest index of different okra varieties

Table 3 illustrates the fresh biomass across various okra varieties. OH-940 and Punjab Selection exhibited the highest average fresh biomass at 4.28 and 4.17 Kg/m², respectively, which is statistically comparable to Swastik-2 (3.86 Kg) and Chiranjeevi F1 (3.50 Kg). In contrast, Gunjan recorded the lowest biomass at 1.63 Kg. The mean fresh biomass per square meter was calculated as 3.07 Kg, yielding highly significant results at the 0.1% level. This data underscores the variability in fresh biomass among okra varieties, emphasizing the importance of varietal selection in optimizing production outcomes. Furthermore, the investigation revealed noteworthy variability in the harvest index across various research-utilized varieties, as illustrated in Table 3. Variety F1 Glory exhibited the highest harvest index (1.14), followed by Chandani and Punjab Selection with indices of 0.88. Conversely, the lowest harvest index (0.47) was noted in variety Chiranjeevi F1. The statistical significance of these results was established at a 5% level of significance.

Formatted: Font: Bold

Comment [Ma18]: Please follow single style like use () in all the cases.

Comment [Ma19]: Rephrase this sentence

Comment [Ma20]: Again this is the vague sentence.

Table 3: Fresh biomass and harvest index of different varieties of okra

Variety	Fresh biomass (Kg/m ²)	Harvest index (HI)
Arka Anamika	2.01 ^e	0.55 ^{bc}
Chandani	3.27 ^{cd}	0.88 ^{ab}
Chiranjeevi F1	3.50 ^{bc}	0.47 ^c
F1 Glory	2.84 ^d	1.14 ^a
Gunjan	1.63 ^e	0.72 ^{bc}
JK 1666	1.93 ^e	0.79 ^{abc}
OH-940	4.28 ^a	0.68 ^{bc}
Punjab Selection	4.17 ^a	0.88 ^{ab}
R35 Selection	3.17 ^{cd}	0.78 ^{abc}
Swastik-2	3.86 ^{ab}	0.81 ^{abc}
Mean	3.07	0.77
SEM	0.173	0.043
CV (%)	10.26	24.24
LSD	0.54	0.32
F-Test	***	*

*Significant at 5% level of significance, **Significant at 1% level of significance, ***Significant at 0.1% level of significance, CV: Coefficient of variance, LSD: Least Significant Difference, SEM: Standard Error of Mean

Formatted: Font: Bold

The results revealed the significant difference variation in yield and biomass production among distinct okra varieties used in the research. The study revealed notable yield disparities among varieties, with results ranging from 0.45 to 0.91 Kg/m². In contrast, Biswas et al. (2016) observed a wider range of 0.57 to 1.4 Kg/m² in fruit yield, surpassing our findings. Possible factors contributing to this variation may include differences in cultivation practices, environmental conditions, or genetic factors, highlighting the multifaceted nature of yield outcomes in agricultural studies. Similarly, according to Bulo et al. (2019), the highest fruit yield per plant was 0.212 Kg and 0.202 Kg. This result seems to align with the yield given by Punjab selection (0.217 kg) and F1 Glory. Sibsankar et al. (2012) reported that fruit yield per plant ranges from 0.2562 to 0.2744 kg. The presented

Comment [Ma21]: This is a vague sentence please rephrase it

Comment [Ma22]: You should clearly discuss the varieties and conditions

result seems to be highly varied from the result obtained from the reported experiment. This can probably be due to the differences in genotype or management practices or climatic factors. According to Zareen et al. (2017) reported the highest pod yield value was 0.99 Kg/m² which is similar to our findings given by Punjab Selection 0.96 Kg/m². Imoloame & Usman (2018) reported that okra yields varied from 0.185 to 1.812 Kg/m². This result seems intensively low from the result obtained in our study. Disparities in agricultural yield might result from variances in soil quality, agronomic practices, or regional weather variables. According to Sharma & Singh (2023), the study revealed that the greatest biomass yield occurred in treatment T1, employing automatic drip irrigation with 100% soil field capacity, yielded 7.2 tons/ha (0.72 Kg/m²). This outcome appears notably lower than OH-940, it reported the highest yield of 4.28 Kg/m², suggesting potential factors such as robust growth conditions or genetic influences contributing to OH-940's superior fresh plant weight. The disparity underscores the complex interplay of environmental and genetic factors influencing biomass production, warranting further investigation into optimizing growth conditions and understanding genetic contributors to enhance yields. Likewise, the statistical similarity with Punjab selection (4.17 Kg) and OH-940 (4.28 Kg) suggests consistent performance. Gunjan's lower weight (1.63 Kg/m²) may result from environmental stress or genetic traits influencing reduced growth. Statistically, Gunjan, Arka Anamika and JK1666 exhibited comparable weights implying similar genetic potential or adaptation to common growing conditions. Systematically, Attarde et al. (2012) reported a fresh biomass ranging from 51.5 to 72.4g. Our research yielded significantly higher results, potentially attributed to variations in fertilizer doses or differences in the methodology procedures. Additionally, the discrepancies may arise from environmental factors such as soil quality, climate, and cultivation practices, influencing plant growth and overall productivity. Moreover, genetic variations among okra varieties utilized in the studies could contribute to the observed differences in fresh plant weights.

Conclusion

In conclusion, Punjab selection displayed superior performance among the ten examined okra varieties, boasting the highest average fresh biomass and overall yield. In conclusion, among the ten examined okra varieties, Punjab selection displayed superior performance, boasting the highest average fresh biomass and overall yield. Conversely, Arka Anamika, Gunjan, and JK1666 exhibited lower biomass, with Arka Anamika producing the least yield. This highlights the substantial impact of varietal selection on okra productivity. Future research should investigate specific traits contributing to high yields, guiding farmers toward optimal cultivar choices. Exploring agronomic practices and environmental factors affecting yield variations could further enhance overall okra production. These findings offer crucial insights for sustainable agriculture, emphasizing the importance of tailored variety selection. Consequently, based on these parameters, Punjab selection and F1 Glory emerge as promising varieties, recommended for further evaluation and utilization in vegetable cultivation and biomass production.

Ethics committee approval

The authors approve the ethics committee.

Data availability

Data will be made available on request.

References

- Abdelhamid, H. N., & Mathew, A. P. (2021). Cellulose-Based Materials for Water Remediation: Adsorption, Catalysis, and Antifouling. *Frontiers in Chemical Engineering*, 3(December), 1–23. <https://doi.org/10.3389/fceng.2021.790314>
- Akandé, O.M, Oluwatoyinbo I. F, Makinde A.E, Adepoju. S. A, & Adepoju. S.I. (2010). Response of Okra to Organic and Inorganic Fertilization. *Nature and Science*, January. https://www.researchgate.net/publication/266266881_Response_of_Okra_to_Organic_and_Inorganic_Fertilization
- Ampara, I. N., Of, D., & Lanka, S. R. I. (2021). EFFECTS OF DIFFERENT FERTILIZERS ON GROWTH AND YIELD OF OKRA (*Abelmoschus esculentus*) cv. HARITHA IN AMPARA DISTRICT OF SRI LANKA. *International Journal of Biology, Pharmacy and Allied Sciences*, 10(12 (SPECIAL ISSUE PART 2)). <https://doi.org/10.31032/ijbpas/2021/10.12.2050>
- Antar, M., Lyu, D., Nazari, M., Shah, A., Zhou, X., & Smith, D. L. (2021). Biomass for a sustainable bioeconomy: An overview of world biomass production and utilization. *Renewable and Sustainable Energy Reviews*, 139(January), 110691. <https://doi.org/10.1016/j.rser.2020.110691>
- Attarde, S., Narkhede, S., Patil, R., & Ingle, S. (2012). Effect of organic and inorganic fertilizers on the growth and nutrient content of *Abelmoschus esculentus* (okra crop). *International Journal of Current Research*, 4(10), 137–140.
- Brocks, S., & Bareth, G. (2018). Estimating barley biomass with crop surface models from oblique RGB

Comment [Ma23]: Which one?

Comment [Ma24]: Write it in a better way. And should incorporate the intellectual reasoning. You should add the varieties of the comparing studies

Comment [Ma25]: Rephrase it

Comment [Ma26]: Merge this information

Comment [Ma27]: This is the poor way of data presentation. Rephrase it. Reviewer is unable to understate this information

Comment [Ma28]: Please set all the references according to the journal demand.

- imagery. *Remote Sensing*, 10(2). <https://doi.org/10.3390/rs10020268>
- Bulo, A., Umrao, R., & Rina, L. (2019). Effect of different plant spacing on growth and yield of okra (*Abelmoschus esculentus*) under subabul (*Leucaena leucocephala*) based alley cropping system. *International Journal of Chemical Studies*, 7(5), 1345–1349.
- Chai Siah, L. (2017). Extraction of bio-flocculant from okra using hydrothermal and microwave extraction methods combined with a techno-economic assessment. June, 180.
- Freidenreich, A., Chanda, S., Dattamudi, S., & Jayachandran, K. (2022). Effect of Glyphosate and Carbaryl Applications on Okra (*Abelmoschus esculentus*) Biomass and Arbuscular Mycorrhizal Fungi (AMF) Root Colonization in Organic Soil. *Horticulturae*, 8(5). <https://doi.org/10.3390/horticulturae8050415>
- Gomez, K. A., & Gomez, A. A. (1984). *Statistical procedures for agricultural research*. John Wiley & Sons, Ltd.
- Gupta, P. K., Patra, S., & Samanta, K. K. (2021). Potential of Okra for Application in Textiles: A Review. *Journal of Natural Fibers*, 18(11), 1788–1800. <https://doi.org/10.1080/15440478.2019.1697997>
- Imoloame, E. O., & Usman, M. (2018). Weed biomass and productivity of okra (*Abelmoschus esculentus* (L) Moench) as influenced by spacing and pendimethalin-based weed management. *Journal of Agriculture Sciences*, 63(4), 379–398. <https://doi.org/https://doi.org/10.2298/JAS1804379I>
- Jonah, P. M., & Kwaga, Y. M. (2019). Genetic interrelationship among quantitative traits and path analysis of some West African okra (*Abelmoschus caillei*) genotypes. *Agriculture Science and Technology*, 11(1), 3–7. <https://doi.org/10.15547/ast.2019.01.001>
- Lee, M., Lin, Y. L., Chiueh, P. Te, & Den, W. (2020). Environmental and energy assessment of biomass residues to biochar as fuel: A brief review with recommendations for future bioenergy systems. *Journal of Cleaner Production*, 251. <https://doi.org/10.1016/j.jclepro.2019.119714>
- Maclaren, C., Swanepoel, P., Bennett, J., Wright, J., & Dehnen-Schmutz, K. (2019). Cover crop biomass production is more important than diversity for weed suppression. *Crop Science*, 59(2), 733–748. <https://doi.org/10.2135/cropsci2018.05.0329>
- Moosavi, S. A., Aghaalkhani, M., Ghobadian, B., & Fayyazi, E. (2018). Okra: A potential future bioenergy crop in Iran. *Renewable and Sustainable Energy Reviews*, 93(April), 517–524. <https://doi.org/10.1016/j.rser.2018.04.057>
- Moosavi, S. A., Shokuhfar, A., Lak, S., Mojaddam, M., & Alavifazel, M. (2020). Integrated application of biochar and bio-fertilizer improves yield and yield components of cowpea under water-deficient stress. *Italian Journal of Agronomy*, 15(2), 94–101. <https://doi.org/10.4081/ija.2020.1581>
- Nath, H., Saikia, A., Goutam, P. J., Saikia, B. K., & Saikia, N. (2021). Removal of methylene blue from water using okra (*Abelmoschus esculentus* L.) mucilage modified biochar. *Bioresource Technology Reports*, 14(March), 100689. <https://doi.org/10.1016/j.biteb.2021.100689>
- Sharma, V., & Singh, P. K. (2023). Performance of AquaCrop model for predicting yield and biomass of okra (*Abelmoschus esculentus*) crop. *The Indian Journal of Agricultural Sciences*, 93(8), 899–905. <https://doi.org/10.56093/ijas.v93i8.133319>
- Singha, A. S., & Guleria, A. (2015). Utility of chemically modified agricultural waste okra biomass for removal of toxic heavy metal ions from aqueous solution. *Engineering in Agriculture, Environment and Food*, 8(1), 52–60. <https://doi.org/10.1016/j.eaef.2014.08.001>
- Sousa, G. G. de, Mendonça, A. de M., Sales, J. R. da S., Junior, F. B. da S., Moraes, J. G. L., & Sousa, J. T. M. de. (2020). Morphophysiological characteristics of okra plants submitted to saline stress in soil with organic fertilizer. *Comunicata Scientiae*, 11, 1–8. <https://doi.org/10.14295/cs.v11i0.3241>
- Vadenbo, C., Tonini, D., & Astrup, T. F. (2017). Environmental Multiobjective Optimization of the Use of Biomass Resources for Energy. *Environmental Science and Technology*, 51(6), 3575–3583. <https://doi.org/10.1021/acs.est.6b06480>
- Verma, M., Loha, C., Sinha, A. N., & Chatterjee, P. K. (2017). Drying of biomass for utilising in co-firing with coal and its impact on environment – A review. *Renewable and Sustainable Energy Reviews*, 71(February 2015), 732–741. <https://doi.org/10.1016/j.rser.2016.12.101>
- Yang, J., Ching, Y. C., & Chuah, C. H. (2019). Applications of lignocellulosic fibers and lignin in bioplastics: A review. *Polymers*, 11(5), 1–26. <https://doi.org/10.3390/polym11050751>