

Original Research Article

Growth Performance, Haematological and Blood Biochemical Characteristics of Broiler Chickens fed varied Levels of Dietary Shea Caterpillar (*CirinaButyrospermi*) Larvae Meal

ABSTRACT

Aims: The increasing feed cost of poultry production makes a search for suitable alternatives essential. A five-week study was conducted to determine the outcome of dietary shea caterpillarlarvaemeal (SCM)on growth performance, haematological and blood biochemical characteristics of broiler chickens.

Study design:A Completely Randomized Design (CRD) was used for the experiment.

Place and Duration of Study:Department of Animal Science Education (Poultry Section of the Animal farm) of Akenten Appiah-Menka University of Skills Training and Entrepreneurship Development, Asante Mampong, Ghana. The study lasted for five weeks.

Methodology:

Ninety 4-week-old sexed Cobb 500 broiler chickens were assigned to five dietary treatments with three replicates. Each treatment had 18 birds with 6 birds per replicate in a male to female ratio of 4:2. Five dietary treatments were formulated to replace anchovy at 0%, 25%, 50%, 75% and 100% with shea caterpillar larvae meal and designated as SCM^{0%}(The Control), SCM^{25%}, SCM^{50%}, SCM^{75%}, and SCM^{100%} respectively.Data collected included proximate and amino acid analysis of shea caterpillar meal, growth performance, blood profile and cost benefit analysis. Data collected were subjected to analysis of variance with General Linear Model (GLM) procedure of Statistical Analysis System (SAS for windows, version 7).

Results:

The proximate and amino acids results showed that shea caterpillar meal contains 60.42 % crude protein, a crude fibre of 7.90 %, and 3306.76 kcal/kg of metabolizable energy as well as fifteen amino. There was a significant difference (P=0.05) among all the treatment means for daily feed and water intake, final body weight and total weight gain. For mean daily feed intake, SCM^{0%} was similar to SCM^{25%}, but statistically different from SCM^{50%}, SCM^{75%} and SCM^{100%}. Birds fed dietary SCM^{25%} (5134g/bird) consumed more feed while the least was recorded from birds fed SCM^{75%} (4615g/bird).Similar trend was observed for total water intake.

Mean Final weight and total weight gain, birds fed SCM^{0%}, SCM^{25%} and SCM^{50%} were similar, but statistically better than SCM^{75%} and SCM^{100%}.It was noticed that final weight and total weight gain decreases as the inclusion levels of shea caterpillar meal increases.

The haematological and biochemical attributes were not significant except for red cell distribution width coefficient of variation and globulin respectively.

The red cell distribution width coefficient of variation values of birds on all the treatments was similar and more than twice the value obtained for the control treatment SCM^{0%}. The globulin values of the birds fed the control, were similar to SCM^{25%}and SCM^{50%}but were significantly different from birds fed SCM^{75%}and SCM^{100%}.

Conclusion:It wasconcluded that the substitution of SCM for fish meal at a level not exceeding 50% had no detrimental effect but improved growth performance.

Keywords: Shea caterpillar (Cirinabutyrospermi), proximate composition, growth performance, haematological, biochemical, broilers.

1. INTRODUCTION

The present human population is projected at 7.6 billion people and it is estimated that by the year 2050, the world population will be centred around 9.2 billion individuals which will probably put pressure on global food security [1]. To ameliorate this challenge, poultry production primarily has the potential to efficiently accomplish the present and future food security in the world due to their short maturity period. Broiler production sub-sector serves as a source of employment, and income and offers not only superior protein food but also significant vitamins and minerals to reduce malnutrition among the vulnerable in developing countries and across the world at large [2].

Nevertheless, the poultry sector in Ghana is confronted with numerous challenges such as diseases, imported frozen chicken, high cost and scarcity of conventional sources of feed ingredients such as protein leading to adulteration in some cases [3]. More than 70% of the total cost of production of poultry in Ghana is due to feed costs alone [4].

A search for a suitable non-conventional feedstuff especially protein resources as alternative to fish meal is necessary and timely for saving the poultry industry from total collapse. One possible alternative source of animal protein ingredient is edible insects such as the shea caterpillar (*Cirinabutyrospermi*) larvae. The shea caterpillar is an edible insect from the family Saturniidae and order Lepidoptera [5] and is locally called 'Taantuni' in Dagaare/Wale. The shea caterpillars arise seasonally on shea-nut trees around June each year [6] and feed exclusively on the leaves of the host tree. Shea caterpillar larva meal is rich in protein, iron, fat, minerals, and amino acids [7, 8] and serves as a major source of food in the human and fish diets [5, 9].

In spite of the numerous benefits of shea caterpillars, there is limited evidence on its utilization as a source of animal protein ingredient for poultry in Ghana where the shea plant is found. This study was therefore conducted to determine the feeding value of dietary shea caterpillar (*Cirinabutyrospermi*) larva meal on growth performance, haematological and blood biochemical characteristics and cost savings of broiler chickens.

2.0 MATERIAL AND METHODS

2.1 Location of experiment and duration

The study was carried out at the Poultry Section of the Animal farm of the Department of Animal Science Education, of Akenten Appiah-Menka University of Skills Training and Entrepreneurial Development (AAMUSTED), Mampong, Ghana and lasted for five weeks. Asante Mampong is a town positioned in the North-Eastern part of the Ashanti region within longitudes 0° 05"W and 1° 30"W and latitudes 6°55" N and 7°30"N. The Municipality lies within the wet semi-equatorial forest zone lying between the Guinea Savannah to the North and Rain Forest to the South. It has a bimodal rainfall pattern [10].

2.2 Source and preparation of shea caterpillar larva meal

The already-dried shea caterpillar larvae were purchased from the Wa market in the Upper West Region of Ghana. The shea caterpillars were further sun-dried for three consecutive days and milled at the Poultry unit of the Animal Farm using a hammer mill with a 2mm sieve. The resulting product was further milled by a grinding miller into powder and kept in sacks at a clean, dried area until it was incorporated into the experimental diets. The texture and particle sizes of the milled shea caterpillars were similar to fish meal. The appearance (Plates 1) was pale yellow and the scent was like dried ground okra.



A- Shea caterpillar (SC) B- Hammer milled SC C- Milled SC in corn mill

Plate 1: Dried shea caterpillar and milled particle sizes

2.3 Proximate and amino acid analysis of shea caterpillar larva meal

Samples of shea caterpillar larva meal and fish meal were subjected to proximate analysis according to procedure of Association of Official Analytical Chemists [11]. The amino acid composition of the shea caterpillar meal was analysed at Noguchi Memorial Institute for Medical Research, Accra, to determine both indispensable and dispensable amino acids using High-Performance Liquid Chromatography (HPLC) as described by [12].

2.4 Determination of coliform load in milled shea caterpillar

A sample of shea caterpillar was subjected to microbial analysis according to the procedure in [13] in the Microbiology Laboratory at AAMUSTED, Asante Mampong. The total coliforms of the feed ingredient (Shea Caterpillar meal) were tested using Merk's Chromo cult coliform Agar. Ten (10) fold serial dilution was made with phosphate-buffered solution (PBS) up to 10^{-4} . One millilitre (1ml) of each dilution was inoculated by pour plate method triplicate. The plates were incubated in an inverted position at 37°C for 24 hours.

2.4.1 Total plate count

Dark-blue to violet colonies were counted as *Escherichia coli* (*E. coli*) and the salmon to red as other coliforms. Indole test was conducted on salmon to red colonies on a plate to differentiate *E. coli* from other coliforms and the colouration turned into chilly red indicating that the observed colonies were *E. coli* [39].

2.5 Management of pre-experimental birds

Two hundred (200) one-day-old Cobb 500 broiler chicks were obtained from Darko farms located in Kumasi, Ghana, and brooded for 4 weeks in a deep litter system. The birds were fed a starter diet and water *ad libitum* during the brooding period and the recommended medication and routine activities were followed according to the guidelines of Veterinary Service Directorate [14]

2.6 Experimental Birds and Design

Ninety (90) 4-week-old Cobb 500 birds were selected out of the initial stock based on weight and body condition. The selected birds were balanced by weight and allocated to five treatments and replicated three times in a completely randomized design. Each replicate had 4 males and 2 females. The birds were housed in 15 pens with dimensions of 1.8 m x 0.88m with a floor space of 0.264 m^2 per bird.

2.7 Experimental Diets

The composition of the experimental diets is presented in Table 1. Shea caterpillar larva meal was used to partially and completely replace fish meal in five dietary treatments. Five dietary treatments were formulated to replace anchovy at 0%, 25%, 50%, 75% and 100% with shea caterpillar larvae meal and designated as SCM^{0%}, SCM^{25%}, SCM^{50%}, SCM^{75%}, and SCM^{100%} respectively.

Table 1: Composition of experimental diets

| Feed ingredient | SCM ^{0%} | SCM ^{25%} | SCM ^{50%} | SCM ^{75%} | SCM ^{100%} |
|----------------------------|-------------------|--------------------|--------------------|--------------------|---------------------|
| Maize grain | 57.7 | 57.7 | 57.7 | 57.7 | 57.7 |
| anchovy | 16 | 12 | 8 | 4 | 0 |
| Shea Caterpillar | 0 | 4 | 8 | 12 | 16 |
| Wheat bran | 14 | 14 | 14 | 14 | 14 |
| Soya bean meal | 9 | 9 | 9 | 9 | 9 |
| Oyster shell | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 |
| Premix | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 |
| Dicalcium | 1 | 1 | 1 | 1 | 1 |
| Salt | 0.37 | 0.37 | 0.37 | 0.37 | 0.37 |
| Toxin binder | 0.03 | 0.03 | 0.03 | 0.03 | 0.03 |
| Total(kg) | 100 | 100 | 100 | 100 | 100 |
| Calculated analysis | | | | | |
| Crude protein(%) | 20.47 | 20.47 | 20.47 | 20.47 | 20.47 |
| Ether Extract (%) | 3.45 | 3.66 | 3.88 | 4.09 | 4.31 |
| Crude Fibre (%) | 3.42 | 3.7 | 3.97 | 4.25 | 4.52 |
| Lysine (%) | 1.58 | 1.56 | 1.52 | 1.49 | 1.46 |
| Methionine (%) | 0.50 | 0.46 | 0.43 | 0.40 | 0.37 |
| Calcium (%) | 0.65 | 0.51 | 0.37 | 0.23 | 0.09 |
| Phosphorus (%) | 0.75 | 0.67 | 0.59 | 0.51 | 0.43 |
| M E (kcal/kg) | 2762.39 | 2791.43 | 2820.47 | 2849.51 | 2878.55 |

M E- MetabolizableEnergy; SCM- Shea Caterpillar Meal ,%-percentage

2.8 Parameters measured

Data on proximate, amino acid, growth performance, haematological and biochemical characteristics and cost benefit on feed were recorded.

2.8.1 Proximate and amino acid analysis of shea caterpillar larva meal

Samples of shea caterpillar larva meal subjected to proximate analysis according to AOAC [11]. The amino acid composition of the shea caterpillar larvae meal and fish meal were analysed to determine the quantity of both essential and non-essential amino acids using High-Performance Liquid Chromatography (HPLC) as described by [12].

2.8.2 Growth performance

Growth parameters determined were daily feed intake, total feed intake, initial weight, and final body weight. Average daily feed intake was determined by dividing daily feed intake by number of birds per replicate, total feed intake per bird was determined by summation of daily feed intake over the study period. Daily water intake was found by dividing amount of daily water intake by number of birds in a replicate and the values were added together to obtain total water intake. Electronic Kitchen scale (Zhongshan Jinli Electronic Weighing Equipment Co.,Ltd,China) was used to weigh experimental birds to determine initial live weight, weekly weight and final live weight. Total weight gain was calculated by the difference between initial live weights from weekly live weights and feed conversion ratio was calculated by dividing feed intake by weight gain and average feed conversion was then determined.

2.8.3 Haematological parameters

Blood samples were collected from 15 birds selected at random, (one male bird per replicate) after the area was disinfected with cotton wool dampened with methylated spirit. A disposable sterilised 5ml syringe and needle were inserted into the right-wing vein (armpit) of each bird and 2mm of blood were drawn and discharged into sterilised vacutainer test tubes containing Ethylene Diamine Tetra Acetic Acid (EDTA) to prevent coagulation. A Rayto Automatic Haematology Analyser (Oscar Medicare Pvt. Ltd, India) was used to determine the haematological parameters such as Red Blood Cells, White Blood Cells, Platelets, Haemoglobin, Lymphocytes, Mean Corpuscular Haemoglobin, Mean Cell Volume, Mean Corpuscular Haemoglobin Concentration, Red Cell Distribution width standard deviation, and Red Cell distribution width coefficient of variation, as well as Platelets Distribution Width.

2.8.4 Biochemical attributes

Blood samples were collected from 15 birds selected at random from the five treatments involving a male bird per replicate. The blood samples were collected carefully from the right-wing vein of each bird after the area was disinfected with cotton wool moistened with methylated spirit using 5ml disposable sterilized syringes and dispensed gently into plain vacutainer tubes without EDTA and kept in a vacuum flask with ice cubes and analysed for the biochemical attributes using Mindray BA 88A semi-auto Chemistry Analyser (Shenzhen Mindray Bio-Medical Electronic Co.,Ltd,China). Total protein was assessed using the Biuret method whereas albumin was found using the Bromo Cresol Green method. Total cholesterol was determined using the CHOP-PAP method and globulin value was estimated by deducting albumin from total protein. Triglycerides, high-density lipoprotein, and low-density lipoprotein were also determined according to the procedure of [15]

2.8.5 Cost-benefit effectiveness

The cost of feed per bird was calculated by multiplying the per kilogram cost of feed by the total feed consumed per bird. The profit on feed alone was calculated by subtracting the cost of feed from the selling price of a bird [40].

2.9 Statistical analysis

The data collected were subjected to analysis of variance using General Linear Model (GLM) procedure of Statistical Analysis System (SAS for windows, version 7(2008)). Differences between means were separated using the Tukeys test at a 5% level. The statistical model used was defined as:

$$Y_{ijklm} = \mu + \beta_1 X_{1i} + \beta_2 X_{2j} + \beta_3 X_{3k} + \beta_4 X_{4l} + \beta_5 X_{5m} + \epsilon_{ijklm}$$

Where:

Y_{ijklm} represents the response variable being measured (e.g., proximates, feed intake, body weight gain, feed conversion ratio, haematology and biochemical profile).

μ is the overall mean or the intercept term, representing the baseline or average response when all the explanatory variables are zero.

$\beta_1, \beta_2, \beta_3, \beta_4,$ and β_5 the coefficients associated with the explanatory variables $X_1, X_2, X_3, X_4,$ and $X_5,$ respectively.

X_{1i} represents the control treatment, Shea caterpillar larvae meal at 0% (SCM^{0%}).

3. RESULTS AND DISCUSSION

3.1 Proximate analysis of shea caterpillar larva meal

The proximate analysis of the shea caterpillar larvae meal is presented in Table 2. The outcome showed that the caterpillar meal has a high crude protein content of 60.42% which fell within the normal range (20 - 80%) of protein content for edible insects of the order Lepidoptera [16]. The Crude protein of 60.42 % reported in this study is higher than the 55% CP [7] but less than 62%[17] and 63% [12]. Also, Amao *et al.*,[18] reported a superior value of 66.74% for Westwood (*Cirinaforda*) larvae meal. Similarly, [19] reported 50.30% CP content for silkworm caterpillar meals which also disagreed with current findings. The disparity in crude protein content observed is attributed to the quality of shea leaves as feed, and the mode of processing the shea caterpillars. According to [7], the differences in the proximate composition of insects may perhaps be attributed to the host plant since the amount of protein differs within similar species as well as different species of insects.

The ash content of 3.9% reported in this study was lower than the 4.8% [7] and 5.1% [17]. It was observed that some insect larvae meals are rich in ash content than shea caterpillar larvae meals. Ash content indicates the mineral potential in a feed[7]. The difference in ash content could be attributed to the quality of shea leaves consumed and the developmental stages of the caterpillar larvae. The high ash content could also be due to the hard body of the insect owing to the high chitin content[20]. The crude fibre content of 7.9% in this study was in agreement with the 7.5% reported for Westwood(*Cirinaforda*) larvae meal [18], but differed from 2.8%, 9.1% and 10.9% reported by [7], [21] and[19] respectively. Fibre in insects is chitin which cannot be absorbed or degraded into the blood and therefore affect the digestibility of proteins [18, 12]. The fibre content in this study was within the suggested limit of 7 % to improve the efficiency of digestion[22]. The difference in fibre content may perhaps be attributed to the extra fibrous feed of shea caterpillar larvae and the presence of bristles and setae on the caterpillar [12]. The fat value of 10.4% was lower than the 12% [12] and 14.5% indicated by [17]. Also, the nitrogen-free extract content of 6.3% was more than the 3% obtained by [7] and less than the value of 12.6%[17].

Table 2. Proximate composition of shea caterpillar larva meal

| Parameter | Shea caterpillar meal composition |
|--------------------------------|-----------------------------------|
| Moisture content (%) | 11.08 |
| Ash content (%) | 3.9 |
| Crude protein (%) | 60.42 |
| Crude fat (%) | 10.4 |
| Crude fibre (%) | 7.9 |
| Nitrogen Free Extract (%) | 6.3 |
| Metabolizable energy (kcal/kg) | 3306.76 |

3.2 Amino acid composition of shea caterpillar larva meal and fish meal

Table 3 shows the essential and non-essential amino acids composition of shea caterpillar larva meal. The shea caterpillar has fifteen different amino acids with a total amino acid content of 64.52mg/100g which disagreed with 59.52g/100g and 70.39g/100g amino acids reported by [7]and [17] respectively. The differences observed could be attributed to the stage of larvae development, the quality of the shea leaves consumed, the geographical location of shea trees, and the efficiency of the analysis procedures [12].

Similarly, in this study, fish meal used consisted of 15 amino acids with a total content of 37.071mg/100g, which differs from 42.27mg/100g reported by [9]. The disparities observed could be attributed to different types of fish used to prepare fish meal, method of processing, and drying.

The amino acid content of SCM was higher than the fish meal and therefore could be implied that the SCM was better in terms of the relatively high amino acid content of the shea caterpillar larva meal. However, the accurate feeding value is determined by the animal.

Table 3. Amino acid composition of shea caterpillar larva meal and fish meal

| Amino acid | Shea caterpillar (mg/100g) | Fish meal (mg/100g) |
|------------------|----------------------------|---------------------|
| Leucine* | 2.51 | 1.95 |
| Lysine* | 3.78 | 1.39 |
| Phenylalanine* | 2.59 | 5.83 |
| Histidine* | 4.03 | |
| Isoleucine* | 3.69 | 1.95 |
| Threonine* | 3.09 | 1.58 |
| Tryptophan* | 5.49 | 6.72 |
| Valine* | 3.07 | 1.73 |
| Methionine* | 1.18 | 1.75 |
| Total EAA | 29.34mg/100g | 22.89mg/100g |
| Alanine** | 7.35 | 1.57 |
| Aspartic acid** | 2.92 | 1.58 |
| Glutamic acid** | 8.67 | |
| Glycine** | 2.85 | |
| Serine** | 5.95 | 1.59 |
| Proline** | 7.35 | 1.68 |
| Asparagine** | | 1.56 |
| Glutamine** | | 1.39 |
| Tyrosine** | | 4.80 |
| Total NEAA | 35.09mg/100g | 14.17mg/100g |
| Total EAA+NEAA | 64.52mg/100g | 37.01mg/100g |

*Essential Amino acid (EAA), ** Non- Essential Amino Acid (NEAA)

3.3 Microbial analysis of shea caterpillar meal

The microbial analysis showed that shea caterpillar meal contains coliform up to 200,000 CFU/g (expressed as 2.0×10^5) which was higher than estimated limits of 103 CFU/g, but it was safe for consumption. The coliform count in foodstuffs that need supplementary cooking and salty meat is 103 CFU/g Lim *et al.* (2012) as cited in [23].

The coliform levels which were high indicates that the shea caterpillar meal might have been contaminated and this could be attributed probably to unhygienic methods of processing, mode of drying, exposed ways of advertising for buyers at marketplaces, and poor storage facilities. According to Cornell University[24]coliforms have been used as an indicator of unhygienic conditions and the quality of food and water products. Therefore, decent sanitation and personal hygiene, particularly during home-based food processing and storage should be adhered to by all actors' in the caterpillar value chain since the caterpillars are used as food or feed ingredient.

3.4 Effect of shea caterpillar larvae meal on growth performance

Mean daily feed intake, daily water intake, mean final weight and mean total weight gain differed ($P=.05$) among the dietary treatments except total feed intake, and feed conversion ratio (Table 4).

Mean daily feed intake of birds fed SCM^{0%} was similar to SCM^{25%}, but statistically different from SCM^{50%}, SCM^{75%} and SCM^{100%}. Treatments SCM^{50%} and SCM^{100%} were similar in daily feed intake but differed from SCM^{75%}.

Also, SCM^{25%} recorded superior mean daily water intake compared with the other treatment means. However, SCM^{50%} and SCM^{75%} were similar but lower than SCM^{0%} and superior to SCM^{100%}.

Mean final weight and total weight gain of SCM^{0%}, SCM^{25%} and SCM^{50%} were similar ($P=.05$), but statistically better than birds fed SCM^{75%} and SCM^{100%} which were also alike (Table 4).

It was observed that the increasing level of the shea caterpillar larva meal resulted in decreased feed intake of the birds. The reduced feed intake was particularly observed for broilers on the dietary shea caterpillar meal above 50% replacement for anchovy and this was due to reduced palatability and fibre content of the SCM. Decreased feed intake has been reported to be associated with high fibre content of feed [25], energy level, feed shape, health, and management practices affect the consumption of feed by table birds [26], the quality of the feed as well as how efficiently the birds utilize the feed [27]

The lower final body weight and total weight gain of broiler growers fed 75% and 100% replacement level of SCM-based diet is attributed to decreased feed intake. Additionally, chitin component of the shea caterpillar larvae could influence nutrient availability due to stronger bond with some nutrients and thus might have compromised nutrient utilisation for body growth. Campus [28] reported that grasshopper meal inclusion at 100% showed higher feed intake and weight gain which was contrary to the present study. In this study, at 100% inclusion of SCM, the feed intake was lowest and the growth rate was relatively slow.

Table 4: Effect of shea caterpillar larvae meal on growth performance of broiler chickens

| Parameters | Treatments | | | | | SEM | P-v |
|----------------------------------|---------------------|---------------------|---------------------|---------------------|---------------------|-------|-------|
| | SCM ^{0%} | SCM ^{25%} | SCM ^{50%} | SCM ^{75%} | SCM ^{100%} | | |
| Initial weight, g/bird | 720.0 | 726.7 | 733.3 | 720.0 | 726.7 | 7.30 | 0.682 |
| Mean daily feed intake, g/bird | 142.14 ^a | 146.68 ^a | 135.23 ^b | 138.71 ^b | 131.86 ^c | 1.493 | <.001 |
| Total feed intake, g/bird | 4975 | 5134 | 4733 | 4855 | 4615 | 176.9 | 0.329 |
| Mean daily water intake, ml/bird | 329.37 ^b | 335.29 ^a | 314.91 ^c | 316.43 ^c | 306.34 ^d | 0.742 | <.001 |
| Total water intake, ml/bird | 11,528 | 11,735 | 11,022 | 11,075 | 10,722 | 386.1 | 0.400 |
| Final body weight, g/bird | 2794 ^a | 2746 ^a | 2778 ^a | 2393 ^b | 2304 ^b | 86.5 | 0.005 |
| Mean total weight gain, g/bird | 2074 ^a | 2019 ^a | 2045 ^a | 1673 ^b | 1577 ^b | 92.1 | 0.008 |
| Feed Conversion Ratio | 2.50 | 2.67 | 2.47 | 2.94 | 3.00 | 0.145 | 0.080 |

^{abcd} means values bearing different superscripts in the same row are significantly ($P=.05$) different
g= Gram, P= Probability, SEM=Standard Errors of Means, V=Value

3.5 Effect of dietary shea caterpillar larvae meal on haematological parameters of broilers

Haematological parameters of birds fed different inclusion rates of shea caterpillar meal have been presented in Table 5. There was no significant difference ($P=.05$) among all the haematological indices except red cell distribution width coefficient of variation (RDW-CV). The red cell distribution width coefficient of variation of birds of all the treatments was similar and higher than SC^{0%}, the control. The RDW is used with parameters such as MCH, MCHC and MCV to describe a population of RBC. High RDW has been reported to be associated with nutritional deficiency (Particularly vitamin12 and iron) [29,30] and could indicate that the SCM may be deficient in these nutrients, thus requiring supplementation in the diet for better erythropoietin [7].

However, values obtained for WBC were above the reference range reported by [31]. The white blood cells help to fight infections, defend the body against micro-organisms and produce antibodies in immune response [32] and show enhanced well-being of the birds [33] as a result of producing high WBC against potential infection threat or by stimulation of T-cells. The haematological result of this study agrees with the work done by [33] who reported that Black soldier fly meal in broiler diets did not show any significant effect on haematological characteristics of the blood except WBC, MCV and platelets. The result is also similar to the report of [34] and [35] in a study using silkworm caterpillar meal.

Table 5: Effect of shea caterpillar meal on haematological parameters of broiler chickens

| Parameters | Treatments | | | | | SEM | P-value |
|---------------------------|-------------------|--------------------|--------------------|--------------------|---------------------|------|---------|
| | SCM ^{0%} | SCM ^{25%} | SCM ^{50%} | SCM ^{75%} | SCM ^{100%} | | |
| WBC (10 ⁶ /l) | 64.70 | 62.90 | 58.70 | 58.5 | 67.9 | 6.15 | 0.787 |
| LYM (10 ⁶ /l) | 58.00 | 56.90 | 54.60 | 54.4 | 62.1 | 5.18 | 0.824 |
| RBC (10 ¹² /l) | 1.780 | 1.97 | 1.91 | 1.79 | 1.957 | 0.13 | 0.741 |
| HGB(g/dl) | 7.20 | 7.97 | 7.90 | 7.07 | 7.30 | 0.38 | 0.374 |
| MCHC(g/dl) | 31.13 | 31.37 | 31.17 | 30.20 | 30.13 | 0.48 | 0.284 |
| MCH (pg.) | 40.60 | 40.53 | 40.53 | 39.50 | 37.73 | 1.22 | 0.441 |
| MCV(fl) | 130.20 | 129.2 | 130.10 | 131.0 | 124.9 | 2.66 | 0.539 |
| RDW-CV | 4.22 ^b | 11.87 ^a | 11.87 ^a | 11.83 ^a | 12.73 ^a | 1.87 | 0.046 |
| RDW-SD (fl) | 67.30 | 64.40 | 64.70 | 65.40 | 66.1 | 3.01 | 0.961 |
| HCT% | 23.17 | 25.37 | 25.93 | 23.47 | 24.20 | 1.44 | 0.614 |
| PLT (10 ⁹ /l) | 19.00 | 27.30 | 20.00 | 18.70 | 32.0 | 4.51 | 0.214 |
| MPV (fl) | 5.97 | 6.40 | 5.90 | 5.70 | 6.07 | 0.29 | 0.570 |
| PDW | 0.10 | 0.12 | 0.12 | 0.12 | 0.11 | 0.01 | 0.384 |

^{ab} means values bearing different superscripts in the same row are significantly ($P=0.05$) different
SEM=Standard Error of Means=Probability, White Blood Cell (WBC), Lymphocytes(LYM), Red Blood Cell (RBC), Haemoglobin (Hb), Mean Corpuscular Haemoglobin Concentration (MCHC), Mean Corpuscular Haemoglobin (MCH), Mean Corpuscular Volume (MCV), Red Cell Distribution Width Standard Deviation (RDW-SD), Red Cell Distribution Width Coefficient of Variation (RDW-CV), Haematocrit (HCT), Platelets (PLT), Mean Platelets Volume (MPV), platelets Distribution Width (PDW), Mean Platelets Volume (MPV), Percentage (%), L-litre

3.6 Effect of dietary shea caterpillar meal on biochemical parameters of broiler chickens

Table 6 shows the biochemical parameters of birds fed different inclusion rates of shea caterpillar meal. The results showed no significant differences ($P=0.05$) among all the treatment means except globulin. The globulin values of birds fed SCM^{0%} were similar to SCM^{25%} and SCM^{50%} but were significantly different from SCM^{75%} and SCM^{100%}. However, the values for SCM^{25%}, SCM^{50%}, SCM^{75%} and SCM^{100%} were also similar. Globulin plays an important role in liver function, blood clotting, and fighting infection.

Birds fed the control diet (SCM^{0%}) had a higher total protein value (33.3 g/l) than all the SCM treatments. Total protein content decreases as shea caterpillar meal inclusion levels increase. This indicates that the control diet may be of good quality than the SCM probably due to stronger bonding with minerals. The values obtained were all higher than the normal reference range of 3.0g/l - 4.9g/l [36]. This showed that probably the various diets were rich in protein and that the shea caterpillar did not adversely affect the total protein content. This finding was in line with [37] who indicated that the inclusion of maggot meal in the diet of broiler chickens did not show a significant effect for all the biochemical parameters.

Table 6: Effect of shea caterpillar meal on blood biochemical indices of broiler chickens

| Parameters | Treatments | | | | | SEM | P-v |
|----------------------|--------------------|---------------------|---------------------|--------------------|---------------------|--------|-------|
| | SCM ^{0%} | SCM ^{25%} | SCM ^{50%} | SCM ^{75%} | SCM ^{100%} | | |
| Total Protein (g/l) | 33.3 | 27.3 | 27.0 | 25.0 | 22.0 | 2.49 | 0.085 |
| Albumin (g/l) | 11.67 | 9.67 | 10.67 | 10.67 | 8.33 | 0.955 | 0.217 |
| Globulin (g/l) | 21.67 ^a | 17.67 ^{ab} | 16.33 ^{ab} | 14.33 ^b | 13.67 ^b | 1.706 | 0.050 |
| Cholesterol (mmol/L) | 2.70 | 2.50 | 2.97 | 2.67 | 2.67 | 0.2082 | 0.637 |
| Triglyceride(mmol/L) | 0.53 | 0.70 | 1.23 | 0.97 | 0.80 | 0.1125 | 0.255 |
| HDL (mmol/L) | 1.17 | 1.07 | 1.50 | 1.07 | 1.10 | 0.1125 | 0.096 |
| LDL (mmol/L) | 1.30 | 1.10 | 1.17 | 1.17 | 1.20 | 0.1164 | 0.239 |

^{ab} means values bearing different superscripts in the same row are significantly ($P=0.05$) different
SEM=Standard Error of Means, P=Probability, LDL=Low Density Lipoprotein, HDL=High Density Lipoprotein,

3.7 Cost-benefit of incorporating shea caterpillar meal in broiler chicken diets

The shea caterpillar meal at varied inclusion level showed significant differences (Table 7). There was a significant difference among the treatment means in terms of cost of feed per bird, selling price of bird and profit on feed alone. Birds fed SCM^{0%} and SCM^{25%} had higher feed cost while SCM^{100%} incurred the least, but similar to SCM^{50%} and SCM^{75%}. Birds fed the SCM^{0%}, SCM^{25%} and SCM^{50%} had similar selling price which was significantly higher than those fed SCM^{75%} and SCM^{100%}. The profit on feed of birds fed SCM^{0%}, SCM^{25%} and SCM^{50%} was similar and significantly higher than birds fed SCM^{75%} and SCM^{100%}. But profit on the birds fed SCM^{25%} and SCM^{75%} were not different.

The birds fed at 50% SCM, recorded a profit per bird of GH¢38.4 followed by GH¢37.3 for the control (SCM^{0%}) and GH¢35.6 for 25% SCM and GH¢26 for 100% SCM which had the least profit recorded. The inclusion levels at 75% SCM or more reduced the profit margins making such inclusion levels less efficient in commercial poultry production. Hence the best inclusion rate to maximize profit by a farmer should not exceed 50% of the shea caterpillar meal at the finisher phase. The reduction in the cost of the feed per bird on the shea caterpillar meal-based diets could be attributed to the price difference between a fish meal and a shea caterpillar meal. Since the prevailing price of fish meal was expensive compared to the SCM. Oyegokeet *al.*[38] reported that decreasing levels of Westwood (*Cirinaforda*) incorporation in the diets of broiler chickens has led to a reduction of costs of production and therefore, economically suitable to use in diets of broiler chickens. This agreed with the findings of this present study. Similarly,[19]also reported that the replacement of silkworm caterpillar meal with fish meal indicates that cost per kilogram gain reduced as silkworm caterpillar meal inclusion declined at the starter phase suggesting more profit.

Table 7. Cost-benefit effectiveness of using shea caterpillar larvae meal in broiler diets

| Description | Treatments | | | | | SEM | P-v |
|-----------------------|---------------------|--------------------|---------------------|---------------------|---------------------|--------|------|
| | SCM ^{0%} | SCM ^{25%} | SCM ^{50%} | SCM ^{75%} | SCM ^{100%} | | |
| Per kg feed cost | 4.19 | 4.08 | 3.98 | 3.87 | 3.77 | 0.000 | |
| Total feed intake | 4.975 | 5.134 | 4.733 | 4.855 | 4.615 | 0.1769 | 0.33 |
| kg/bird | | | | | | | |
| Cost of feed per bird | 20.76 ^{ab} | 20.97 ^a | 18.83 ^{bc} | 18.80 ^{bc} | 17.34 ^c | 0.650 | 0.03 |

(GH¢)

| | | | | | | | |
|------------------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------|------|
| Total Wt. Gain kg/bird | 2.074 ^a | 2.019 ^a | 2.045 ^a | 1.673 ^b | 1.577 ^b | 0.0921 | 0.01 |
| Feed Cost per kg | 10.07 | 10.44 | 9.36 | 11.26 | 11.03 | 0.610 | 0.17 |
| body Wt. Gain | | | | | | | |
| Selling price of birdGH¢) | 58.1 ^a | 56.5 ^a | 57.2 ^a | 46.8 ^b | 44.2 ^b | 2.35 | 0.01 |
| Profit on feed alone | 37.3 ^a | 35.5 ^{ab} | 38.4 ^a | 28.0 ^{bc} | 26.2 ^c | 2.50 | 0.01 |

^{abc} means values bearing different superscript in the same row are significantly ($P=0.05$) different.

SEM-standard errors of means, Wt. =weight, Kg=kilogram. P=Probability, V=value. One (1) US Dollar=GH¢11.98

4. Conclusion

This study concludes that dietary shea caterpillar larvae meal has a considerable amount of nutrients especially crude protein value and amino acids which can be utilized as an organic feed resource for broiler chickens. Shea caterpillar larvae meal had no detrimental effect on the growth performance and blood profile of broilers. The mean cost of feed per kilogram and mean cost of feed per bird declined gradually with increasing shea caterpillar larvae meal and at a rate of 50% or lower improves profit. SCM at a 50% inclusion rate or below is recommended in broilers diet while further studies involving digestibility of SCM is suggested.

CONSENT

All authors declared that 'written informed' consent was obtained from the approved parties for publication of this article and accompanying images.

ETHICAL APPROVAL (WHEREEVER APPLICABLE)

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee"

REFERENCES

1. George, S. (2018) Feeding the World in 2050 and Beyond. Michigan State University Extension retrieved from <https://www.canr.msu.edu/news/feeding-the-world-in-2050-and-beyond-part-1> on July 20, 2021
2. Ravindran, V. (2013). Poultry feed availability and nutrition in developing countries. *Poultry development review*, 2, 60-63.
3. Hagan, M. A. S. (2020). Challenges of poultry farmers at Ejisu municipality, Ghana. *Asian Journal of Agricultural Extension, Economics and Sociology*, 63-77.
4. Banson, K., Muthusamy, G., & Kondo, E. (2015). The import substituted poultry industry; evidence from Ghana.
5. Anankware, J. P., Osekre, E. A., Obeng-Ofori, D., & Khamala, C. (2016). Identification and classification of common edible insects in Ghana. *International Journal of Entomology Research*, 1(5), 33-39.
6. Bama, H. B., Dabire, R. A., Ouattara, D., Niassy, S., Ba, M. N., & Dakouo, D. (2018). Diapause disruption in *Cirina butyrospermi* Vuillet (Lepidoptera, Attacidae), the shea caterpillar, in Burkina Faso. *Journal of Insects as Food and Feed*, 4(4), 239-245.
7. Yapo, M. L., Amara, R. F., & Tuo, Y. (2017). Nutritional value of shea caterpillar (*Cirina butyrospermi* Vuillet) sold at the market of Korhogo (Côte d'Ivoire). *International Journal of Agronomy and Agricultural Research*, 10(5), 35-44.

8. Payne, C., Badolo, A., Sagnon, B., Cox, S., Pearson, S., Sanon, A., Bationo, F., & Balmford, A. (2020a). Effects of defoliation by the edible caterpillar “chitoumou” (*Cirinabutyrospermi*) on harvests of shea (*Vitellaria paradoxa*) and growth of maize (*Zea mays*). *Agroforestry Systems*, 94(1), 231–240.
9. Anvo, M. P. M., Aboua, B. R. D., Compaoré, I., Sissao, R., Zoungrana-Kaboré, C. Y., Kouamelan, E. P., & Toguyéni, A. (2017). Fish meal replacement by *Cirinabutyrospermi* caterpillar's meal in practical diets for *Clarias gariepinus* fingerlings. *Aquaculture Research*, 48(10), 5243-5250.
10. Ghana Statistical Service (2014). Population and Housing Census District Analytical Report Mampong Municipal.
11. AOAC. (1995). Official methods of Analysis of AOAC International, 16th Ed, (Vol.1). AOAC International Arlington, USA.
12. Anankware, J. P. (2017). Neglected and underutilised insects of Ghana: Identification, spatial distribution and utilisation for food, feed and nutrition [PhD Thesis]. Kumasi: Kwame Nkrumah University of Science and Technology.
13. Merck Microbiology Manual 12th edition retrieved on 12th July, 2021 on https://www.laboquimia.es/pdf_catalogo/MERCK_Manual_de_microbiologia_12a_edicion.pdf.
14. Veterinary Services Directorate (VAD). (2018) National Harmonised Poultry Vaccination Schedule. VSD/GPP/USDA Publication.
15. Keller A (1984) Total Serum Protein Determination. Clinical Chemistry, Theory, Analysis and Correlation. St. Louis, Mosby Company, USA.
16. Hlongwane, Z. T., Slotow, R., & Munyai, T. C. (2020). Nutritional composition of edible insects consumed in Africa: A systematic review. *Nutrients*, 12(9), 2786.
17. Anvo, M. P. M., Toguyéni, A., Otchoumou, A. K., Zoungrana-Kaboré, C. Y., & Kouamelan, E. P. (2016a). Nutritional qualities of edible caterpillars *Cirinabutyrospermi* in southwestern of Burkina Faso. *International Journal of Innovation and Applied Studies*, 18(2), 639-645.
18. Amao, O. A., Oladunjoye, I. O., Togun, V. A., Olubajo, K., & Oyaniyi, O. (2010). Effect of Westwood (*Cirinaforda*) larva meal on the laying performance and egg characteristics of laying hen in a tropical environment. *International Journal of Poultry Science*, 9(5), 450–454.
19. Ijaiya, A. T., & Eko, E. O. (2009a). Effect of replacing dietary fish meal with silkworm (*Anapheinfrecta*) caterpillar meal on growth, digestibility and economics of production of starter broiler chickens. *Pakistan Journal of Nutrition*, 8(6), 845-849.
20. Raksakantong, P., Meeso, N., Kubola, J., & Siriamornpun, S. (2010). Fatty acids and proximate composition of eight Thai edible terri-colous insects. *Food Research International*, 43(1), 350–355.
21. Adeniji, A. A. (2007). Effect of replacing groundnut cake with maggot meal in the diet of broilers. *International Journal of Poultry Science*, 6(11), 822-825.
22. Koney E.B.M, (2004). *Poultry Health and Production*. 4th Edition, Advent Press. Accra, Ghana.
23. Ramashia, S. E., Tangulani, T., Mashau, M. E., & Nethathe, B. (2020). Microbiological quality of different dried insects sold at Thohoyandou open market, South Africa. *Food Res*, 4, 2247–2255.
24. Cornell University. (2007). Coliform Bacteria. Indicators in food and water. Dairy Foods Science Notes, 1-2.
25. Onu, P. N., Madubuike, F. N., Onu, D. O., & Ekenyem, B. U. (2011). Performance and economic analysis of broiler starter chicks fed enzyme supplemented sheep manure-based diets. *ARPJ Agric Biol Sci*, 6, 14-1.
26. Ferket, P. R., & Gernat, A. G. (2006). Factors that affect feed intake of meat birds: A review. *Int. J. Poult. Sci*, 5(10), 905-911.
27. Egbunike, G. N., Agiang, E. A., Owosibo, A. O., & Fatufe, A. A. (2009). Effect of protein on performance and haematology of broilers fed cassava peel-based diets. *Archivos de zootecnia*, 58(224), 655-662.

28. Campus, L. (2009). The Effect of Replacing Graded Levels of Fish meal with Grasshopper Meal in Broiler Starter Diet. * Hassan, AA** Sani, I,* Maiangwa, MW and*** Rahman, SA. *PAT*, 5(1), 30-38.
29. Cabo del Riego, J. M., Núñez-Iglesias, M. J., Paz Carreira, J., Blanco Hortas, A., Álvarez Fernández, T., NovíoMallón, S., ... & Freire-Garabal Núñez, M. (2022). Red Cell Distribution Width as a Predictive Factor of Celiac Disease in Middle and Late Adulthood and Its Potential Utility as Celiac Disease Screening Criterion. *International Journal of Environmental Research and Public Health*, 20(1), 66.
30. Fici, F., Celik, T., Balta, S., Iyisoy, A., Unlu, M., Demitkol, S., & Grassi, G. (2013). Comparative effects of nebivolol and metoprolol on red cell distribution width and neutrophil/lymphocyte ratio in patients with newly diagnosed essential hypertension. *Journal of cardiovascular pharmacology*, 62(4), 388-393.
31. Aengwanich, W., Simaraks, S., & Chinrasri, O. (2004). Hematological, electrolyte and serum biochemical values of the Thai indigenous chickens (*Gallus domesticus*) in northeastern, Thailand. *Songklanakarin Journal of Science and Technology*, 26(3), 425-430.
32. Etim, N. N., Williams, M. E., Akpabio, U., & Offiong, E. E. (2014). Haematological parameters and factors affecting their values. *Agricultural Science*, 2(1), 37-47.
33. Opoku, O., Atuahene, C. C., Adjei, M. B., Quaye, B., & Adu, M. A. (2018). Black soldier fly (*hermetia illucens*) larvae meal (BSFLM) as alternative dietary protein source for broiler chickens; effects on growth performance, carcass and haematological parameters. *Ghanaian Journal of Animal Science*, 9(1), 25-31.
34. Biasato, I., Gasco, L., De Marco, M., Renna, M., Rotolo, L., Dabbou, S. & Schiavone, A. (2017). Effects of yellow mealworm larvae (*Tenebrio molitor*) inclusion in diets for female broiler chickens: implications for animal health and gut histology. *Animal Feed Science and Technology*, 234, 253-263.
35. Ijaiya, A. T., & Eko, E. O. (2009b). Effect of replacing dietary fish meal with silkworm (*Anaphe infracta*) caterpillar meal on performance, carcass characteristics and haematological parameters of finishing broiler chicken. *Pakistan Journal of Nutrition*, 8(6), 850-855.
36. Cynthia, K. (2011). The Merck Veterinary Manual 10th (tenth) edition. *Merck Sharp & Dohme Corp.*
37. Akpodiete, O. J., & Okagbare, G. O. (2002). Haematological and biochemical indices of meat-and egg-type chickens fed maggot meal. *Nigerian Journal of Animal Science*, 5(1).
38. Oyegoke, O. O., Ayandiran, T. A., & Akintola, A. J. (2013). Cost effectiveness of replacing fish meal with *Cirinaforda* (westwood) larva in the diets of broilers. *Agr Biol JN Am*, 3(4), 55-59.
39. Naratama, M. R., & Santoso, I. (2020). Non-fecal and fecal coliform tests of ready-to-eat food and drinks using fluorogenic and chromogenic media. In *Journal of Physics: Conference Series* (Vol. 1442, No. 1, p. 012064). IOP Publishing.
40. Kwafo, S., Kagya-Agyemeng, J.K., Annor, S.Y. and Bonsu, F.R.K. (2019). Growth performance, blood profile and economic efficiency of layer chicks fed diets containing graded levels of moringa leaf meal at pre-laying phase. *International Journal of Innovative Research and Development* 8(9):132-138. DOI No.:10.2494/ijird/2019/v8/i9/SEP19024