

# **Evaluation of okra varieties for morphological and phenological traits cultivated under Sub-Humid plains of South-East Rajasthan**

## **ABSTRACT**

Vegetable crop okra is well known for its Indian vegetable export and household consumption contribution and thus should be explored for cultivation under various agro climatic zones. Higher production of this crop is possible by the cultivation of varieties which shows remarkable enhanced returns, compared to other cultivars grown at same climatic conditions and inputs applied. Hadoti region of South-East Rajasthan, can serve to okra production in better way if the productive varieties can be screened for this region, which is lacking till date. Hence an attempt is made to recommend a variety, which performs better under South-East climatic conditions of Rajasthan. The present study was conducted in the summer season during 2022 and 2023 at the Instructional Farm, Department of Horticulture, School of Agricultural Sciences, Career Point University, Kota, Rajasthan. 12 varieties of okra viz. Parbhani Kranti, Parbhani Bhindi, Arka Anamika, Shahiba, Shan, No. 55, Harita, Jhilmil, No. 64, Sonal, Shakti, and Punjab Selection were evaluated in randomized block design, replicated thrice. Analysis of variance revealed that all varieties differed significantly for all the traits. There was a wide range of variation for earliness, growth, yield and quality parameters as well. These results showed presence of significant varietal effect. However, the present study demonstrated the superiority of Arka Anamika over other varieties in terms of maximum mean plant height (89.07 cm), number of leaves per plant (25.64), number of branches per plant (13.88), leaf length (19.03 cm), leaf width (25.32 cm) and leaf area (250.04 cm<sup>2</sup>) while minimum mean days taken to flowering opening to fruit picking (5.51 DAS) and days taken to first picking (47.16 DAS) was recorded in variety Punjab Selection under soil and climatic condition of Hadoti region of Rajasthan.

**KEYWORDS:** *Evaluation, Flowering, Growth, Varieties*

## **INTRODUCTION**

Okra [*Abelmoschus esculentus* (L.) Moench] has occupied a prominent position among

vegetables. Okra is known by many local names in different parts of the world. It is called lady's finger in England, Gumbo in U.S.A. and Bhindi in India. In India, okra is one of the most important vegetable crop grown for its tender green fruits during summer and rainy seasons. It is a short duration crop propagated through seeds, cherished for its tender and scrumptious green fruits used in curries, soups or in canned, dehydrated or frozen forms for off-season consumption (Neeraja *et al.*, 2004). Okra is more remunerative than the leafy vegetables, while crop has not adapted in India as leafy vegetable as in for East countries. The roots and stems are useful for clearing cane juice from which gur or jaggery is prepared (Chauhan, 1972). Its ripe seeds are roasted, ground and used as a substitute for coffee in Turkey (Mehta, 1959).

The foremost challenges faced by okra crop is that it is ravaged by many species of insect-pests and diseases throughout its growth period. Among these, jassid, fruit and shoot borer, powdery mildew and yellow vein mosaic are quite serious and major restraining factors in okra cultivation. The challenge to the existence of mankind has always been to produce adequate quantity of food from the available acreage to meet the requirements of ever expanding world population. The rate of yield gain in crop improvement programme must match the rate of population growth so, as to avoid malnutrition and hunger. A lot of okra hybrids/varieties are being grown by the farmers, but best performing hybrids/varieties of okra having desirable quantitative and qualitative characters such as adaptability to adverse environments and resistance to biotic and abiotic stresses resulting into better monetary return to the vegetable growers. Keeping in view, it is essential to work out on the appropriate quantitative and qualitative characters of okra crop so that maximum yield and high-quality produce could be obtained. This is common fact that the varieties showing better performance under one locality may not be suitable for another locality or region. So that the main aim of experiment is to find out the better performance of a particular okra variety in Hadoti region of Rajasthan and screen out to which variety well adapted and produced maximum yield to another variety under South-East region of Rajasthan.

## **Materials and Methods**

### ***Location***

Kota district is located at 25.18° N to 75.83° E Latitude in South Eastern Rajasthan. It covers an area of 221.36 km<sup>2</sup>. Agro-climatically, the district falls in Zone V, known as Humid South Eastern Plain. The average rainfall in the region is 660.6. mm. Maximum temperature range in the summer is 40 to 48°C and minimum 1.0- 2.6°C during winter. Major *kharif* crops of the district are soybean, maize and pulses. While in *rabi*, wheat, mustard, coriander and garlic are main crops.

### ***Experimental Details***

The present studies were carried out at the Department of Horticulture, School of Agricultural Sciences, Career Point University, Kota (Rajasthan) during the year 2022 and 2023. 12 varieties of okra viz. Parbhani Kranti, Parbhani Bhindi, Arka Anamika, Shahiba, Shan, No. 55, Harita, Jhilmil, No. 64, Sonal, Shakti, and Punjab Selection were evaluated in randomized block design, replicated thrice. Raised bed planting system at 60 x 30 cm spacing was adapted to grow the crop. The FYM 20 tonnes and RDF 150:100:100 NPK applied per hectare respectively, as basal dose.

### ***Measurement of the morphological parameters***

The plant height was taken from the tagged plant in each replication with the help of a meter scale, total number of leaves were counted from tagged plants in each replication after completion of harvesting period and expressed as average number of leaves per plant, number of branches of selected plants was counted and average was worked out, length of leaf and width was measured from randomly selected five leaves from every genotype with the help of scale and then average was recorded, leaf area of random five plants from each treatment was recorded by using electronic leaf area meter. The all plant morphological parameters was recorded at 30,60 and 90 days after sowing the crop.

### ***Measurement of the phenological parameters***

Days taken to 50% flowering was recorded after sowing the crop when 5-6 plants in each replication started to flower and average number of days from sowing date was calculated to make the observation. The number of days taken to first picking was recorded from sowing date was calculated to make the observation. Number of days required from the date of sowing to first picking of fruits was recorded.

## RESULTS

### *Morphological parameters*

Data observed on morphological parameters showed that there was considerable variation among the all varieties during all the stage of plant growth. Among the different varieties the maximum mean plant height (89.07 cm), number of leaves per plant (25.64), number of branches per plant (13.88), leaf length (19.03 cm), leaf width (25.32 cm) and leaf area (250.04 cm<sup>2</sup>) was found superior in variety Arka Anamika. However, lowest plant height (64.39cm), number of leaves per plant (18.24) and number of branches per plant (10.22) was recorded in Shan variety. The leaf length (13.97 cm), leaf width (19.54 cm) and leaf area (204.89 cm<sup>2</sup>) was found lowest in Punjab Selection variety 90 days after sowing (Table 1 to 6).

### *Phenological parameters*

Data observed on phenological parameters showed that there was considerable variation in different varieties. The minimum mean days taken to 50% flowering was recorded in Shakti (43.15 DAS) whereas the maximum days taken to 50% flowering was found in Shan (47.28 DAS). Similarly, minimum mean days taken to flowering opening to fruit picking was recorded in Punjab Selection (5.51 days) whereas the maximum days taken to flowering opening to fruit picking was found in Shan (9.51 days). However, minimum mean days taken to first picking was recorded in Punjab Selection (47.76 days) whereas the maximum days (54.02 days) taken to first picking was found in variety Shan (Table 7).

## DISCUSSION

It is clear from the experimental results that significant variations were recorded amongst the different varieties of okra for the studied on growth parameters viz. plant height, number of leaves per plant, number of branches per plant, leaf length, leaf width and leaf area.

The maximum mean plant height was recorded in “Arka Anamika” (89.07cm) while the lowest plant height was found in “Shan” (64.39cm). The significant differences amongst the varieties for plant height could be attributed to their genetic constitution and inheritance of the character as well as differences in apical dominance controlled by the endogenous auxin level of the genotypes. The obtained results are in accordance with earlier finding of Sharma *et al.* (2016), Singh *et al.* (2017) for plant height; Kandasamy *et al.* (2015), Darshan *et al.* (2016) and Deepanshu and Shamad (2017).

The maximum mean numbers of branches per plant(13.88) and leaves per plant(25.64) were found in 'Arka Anamika' while the minimum numbers of branches per plant (10.22) and leaves per plant (18.24) were found in 'Shan'. The differences amongst the varieties and hybrids for number of branches per plant could be attributed to their differential endogenous auxin and cytokinin contents which regulate the lateral growth and bud development, as low auxin and higher cytokinin levels promote lateral growth and bud development or vice-versa. The variation in number of leaves per plant may be due to varying branching in plants. The findings are in agreement with the results of Deepanshu and Shamad (2017), Simon *et al.* (2013), Muluken *et al.* (2016) and Kumar *et al.* (2019) for number of branches per plant and Kumar *et al.* (2016) for number of leaves per plant.

The maximum mean leaf length(19.03 cm) leaf width(25.32 cm) were found in 'Arka Anamika' while the minimum mean leaf length (10.22) and leaf width (19.54) were found in 'Punjab Selection'. The differences amongst the different varieties for leaf length and leaf width could be attributed to their differential endogenous auxin and cytokinin contents which regulate the lateral growth and bud development, as low auxin and higher cytokinin levels promote lateral growth and bud development or vice-versa. The variation in length and leaf width may be due to varying branching in plants. The findings are in agreement with the results of Deepanshu and Shamad (2017) and Kumar *et al.* (2019) and Das *et al.* (2012) for number of branches per plant and Kumar *et al.* (2016) and Singhet *et al.* (2017) for number of leaves per plant.

The largest leaves(250.04 cm<sup>2</sup> area) were recorded in 'Arka Anamika' while the smallest leaves (204.89 cm<sup>2</sup> area) in 'Punjab Selection'. The variation in leaf area amongst the different varieties of okra might be due to their genetic makeup governing the nutrient uptake behaviour, photosynthetic efficiency and their environmental interactions. The observed results are in accordance to Chandramouli *et al.* (2016) and Bendale *et al.* (2003).

The flowering parameters *viz.* days taken to 50% flowering, days taken to flowering opening to fruit picking and days taken to first picking were significantly influenced by the different varieties of okra.

The minimum mean days taken to 50% flowering were recorded in 'Shakti'(43.15 DAS), while the most late 50% flowering were recorded in 'Shan'

(47.28 DAS). Similarly minimum mean days taken to flowering opening to fruit picking was recorded in “Punjab Selection” (5.51 days) whereas the maximum mean days taken to flowering opening to fruit picking was found in “Shan” (9.51 days). On the other hand minimum mean days taken to first picking was recorded in “Punjab Selection” (47.76 days), whereas the maximum days taken to first picking was found in “Shan” (54.02 days). The early or late production of flowers by different varieties and hybrids of okra might have been genetically controlled and better adaptability to particular environmental conditions as supported by the earlier findings of Kandasamy *et al.* (2015), Kumar *et al.* (2017), Duggi *et al.* (2013), Chandra *et al.* (2014), Kumar *et al.* (2013) and Bagwale *et al.* (2016) for day taken to 50 per cent flowering and Adeoluwa and Kehinde (2011) for early flowering.

## CONCLUSION

Based on the overall effect the variety Arka Anamika was highest in terms of all growth characteristics but Punjab Selection was found earliness in flowering and fruiting. Hence, variety Punjab Selection may be considered worth for better harvest of the crop under Hadoti region or South-East Rajasthan.

## REFERENCES

- Adeoluwa, O.O. and Kehinde, O.B. (2011). Genetic variability studies in West Africanokra (*Abelmoschus caillei*). *Agriculture and Biological Journal of North America*, **2** (10): 1326-1335.
- Bagwale, S.B., Jawale, L.N., Deosarkar, D.B. and Jadhav, R.A. (2016). Genetic variability studies for yield, yield contributing and quality traits inokra [*Abelmoschus esculentus* (L.) Moench.]. *Indian Journal of Agricultural Research*, **50** (6): 614-618.
- Bendale, V.W., Kadam, S.R., Bhave, S.G., Mehta, J.L. and Pethe, U.B. (2003). Genetic variability and correlation studies inokra [*Abelmoschus esculentus* (L.) Moench]. *O rissa Journal of Horticulture*, **31**(2): 1-4.
- Chandra, S., Bhardwaj, M., Kumar, R., Kumar, D., Kumar, S., Gautam, N., Dogra, B. and Sharma, S. (2014). Estimation of parameters of variability for

different quantitative traits in okra [*Abelmoschus esculentus* (L.) Moench]. *International Journal of Farm Sciences*, **4**(3): 33-41.

Chandramouli, B., Shrihari, D., Rao, A. V. D. D. and Rao, M. P.

(2016). Studies on genetic variability, heritability and genetic advance in okra [*Abelmoschus esculentus* (L.) Moench] genotypes. *Plant Archives*, **16**(2): 679-682.

Chauhan, D. V. S. (1972). *Vegetable Crops*. Naya Prakash, Calcutta-6. pp. 711.

Darshan S. Nikitha J. and Arya K. (2016). Genetic variability studies for yield and yield parameters in okra [*Abelmoschus esculentus* (L.) Moench]. *Advances in life sciences*, **5**(17): 6539-6541.

Das, S., Chattopadhyay, A., Chattopadhyay, S. B., Dutta, S. and Hazra, P. (2012). Genetic parameters and path analysis of yield and its components in okra [*Abelmoschus esculentus* (L.) Moench] at different sowing dates in the Gangetic plains of eastern India. *African Journal of Biotechnology*, **11**(95): 16132-16141.

Deepanshu and Shamad, A. (2017). Genetic variability, heritability and correlation coefficient in okra [*Abelmoschus esculentus* (L.) Moench] in Allahabad agro climatic conditions. *Plant Archives*, **17**(2): 1597-1602.

Duggi, S., Magadum, S. K., Srinivasraghavan, A., Kishor, D. S. and Oommen, S. K. (2013). Genetic analysis of yield and yield-attributing characters in okra [*Abelmoschus esculentus* (L.) Moench]. *International Journal of Agriculture Environment and Biotechnology*, **6**(1): 45-50.

Kandasamy, R.

(2015). Genetic divergence in okra [*Abelmoschus esculentus* (L.) Moench]. *International Journal of Agricultural Sciences*, **11**(1): 158-160.

Kumar, A., Kumar, S. and Maji, S. (2016). Genetic variability, heritability and genetic advance studies in okra [*Abelmoschus esculentus* (L.) Moench]. *International Journal of Agriculture Sciences*, **8**(57): 3122-3124.

Kumar, A., Kumar, M., Sharma, V. R., Singh, M. K., Singh, B. and Chand, P. (2019). Genetic variability, heritability and genetic advance studies in genotypes of okra

[(*Abelmoschus esculentus* (L.) Moench]. *Journal of Pharmacognosy and Phytochemistry*, **8**(1): 1285-1290.

Kumar, K.V.V., Venkatesha, K.T., Asif, M., Gangappa E. and Pitchaimuthu M. (2013). Genetic variability studies in okra [*Abelmoschus esculentus* (L.) Moench]. *International Journal of Plant Sciences*, **8**(1): 187-192.

Mehta, Y. R. (1959). *Vegetable Crops*. NayaPrakash, Calcutta-6. pp.711.

Muluken, D., Wassu, M. and Endale, G. (2016). Variability, heritability and genetic advance in ethiopian okra [*Abelmoschus esculentus* (L.) Moench] collections for tender fruit yield and other agro-morphological traits. *Journal of Applied Life Sciences International* **4**(1): 1-12.

Neeraja, G.Vijaya, M.Chiranjeevi, C.Gautham, B. (2004) Screening of okra hybrids against pest and diseases. *Indian Journal of Plant Protection*; 2004. **32**(1): 129-131.

Sharma, P.K., Mishra, D.P. and Pandey, A. (2016). Genetic variability studies for yield and its contributing traits in okra [*Abelmoschus esculentus* (L.) Moench]. *Journal of Applied and Natural Science*, **8**(3): 1634-1637.

Simon, S.Y., Gashua I.B., and Musa, I. (2013). Genetic variability and trait correlation studies in okra [*Abelmoschus esculentus* (L.) Moench]. *Agriculture and Biology Journal of North America*, **4**(5): 532-538.

Singh, A.P., Kumar, P.P., Kumar, B.A. and Bahadur, V. (2017). Studies on genetic variability, heritability and character association in okra [*Abelmoschus esculentus* (L.) Moench]. *International Journal of Bio-resource and Stress Management*, **8**(3): 457-462.

Singh, N., Singh, D.K., Sati, U.C., Rawat, M. and Pandey, P. (2017). Genetic analysis studies in Okra [*Abelmoschus esculentus* (L.) Moench]. *Indian Journal of Pure and Applied Bioscience*, **5**(4): 361-367.

**Table .1 Performance of different varieties of okra with respect to plant height**

| Treatment       | Varieties                     | Plant height (cm) 30 |             |             | Plant height (cm) 60 |             |             | Plant height (cm) 90 |             |             |
|-----------------|-------------------------------|----------------------|-------------|-------------|----------------------|-------------|-------------|----------------------|-------------|-------------|
|                 |                               | DAS                  |             |             | DAS                  |             |             | DAS                  |             |             |
|                 |                               | 2022                 | 2023        | Pooled      | 2022                 | 2023        | Pooled      | 2022                 | 2023        | Pooled      |
| T <sub>1</sub>  | Parbhani Kranti               | 16.68                | 18.36       | 17.52       | 29.97                | 32.11       | 31.04       | 71.97                | 73.48       | 72.72       |
| T <sub>2</sub>  | Parbhani Bhindi               | 16.35                | 19.03       | 17.69       | 28.88                | 31.02       | 29.95       | 74.53                | 76.04       | 75.28       |
| T <sub>3</sub>  | Arka Anamika                  | 19.43                | 22.11       | 20.77       | 38.17                | 40.31       | 39.24       | 88.32                | 89.83       | 89.07       |
| T <sub>4</sub>  | Shahiba                       | 17.43                | 20.11       | 18.77       | 30.63                | 32.77       | 31.70       | 76.55                | 78.06       | 77.3        |
| T <sub>5</sub>  | Shan                          | 16.03                | 18.71       | 17.37       | 27.03                | 29.17       | 28.10       | 63.64                | 65.15       | 64.39       |
| T <sub>6</sub>  | No. 55                        | 16.46                | 19.21       | 17.83       | 28.41                | 30.55       | 29.48       | 68.30                | 69.81       | 69.05       |
| T <sub>7</sub>  | Harita                        | 16.99                | 19.67       | 18.33       | 30.05                | 32.19       | 31.12       | 69.18                | 70.69       | 69.93       |
| T <sub>8</sub>  | Jhilmil                       | 16.16                | 18.84       | 17.50       | 29.47                | 31.61       | 30.54       | 70.45                | 71.96       | 71.2        |
| T <sub>9</sub>  | No. 64                        | 17.52                | 19.20       | 18.36       | 32.68                | 34.82       | 33.75       | 79.33                | 80.84       | 80.08       |
| T <sub>10</sub> | Sonal                         | 18.03                | 20.71       | 19.37       | 34.9                 | 37.04       | 35.97       | 82.93                | 84.44       | 83.68       |
| T <sub>11</sub> | Shakti                        | 17.19                | 19.87       | 18.53       | 30.69                | 32.83       | 31.76       | 67.99                | 69.50       | 68.74       |
| T <sub>12</sub> | Punjab Selection              | 15.90                | 18.58       | 17.24       | 29.03                | 31.17       | 30.10       | 71.01                | 72.52       | 71.76       |
|                 | <b>SEm±</b>                   | <b>1.63</b>          | <b>1.71</b> | <b>0.82</b> | <b>1.66</b>          | <b>1.68</b> | <b>1.65</b> | <b>2.24</b>          | <b>2.02</b> | <b>1.78</b> |
|                 | <b>C.D.(p<sub>0.05</sub>)</b> | <b>2.31</b>          | <b>5.02</b> | <b>2.40</b> | <b>4.88</b>          | <b>4.94</b> | <b>4.86</b> | <b>6.59</b>          | <b>5.92</b> | <b>5.22</b> |

**Table 2 Performance of different varieties of okra with respect tonumber of leaves per plant**

| Treatment       | Varieties           | Number of Leaves 30 |             |             | Number of Leaves 60 |             |             | Number of Leaves 90 |             |             |
|-----------------|---------------------|---------------------|-------------|-------------|---------------------|-------------|-------------|---------------------|-------------|-------------|
|                 |                     | DAS                 |             |             | DAS                 |             |             | DAS                 |             |             |
|                 |                     | 2022                | 2023        | Pooled      | 2022                | 2023        | Pooled      | 2022                | 2023        | Pooled      |
| T <sub>1</sub>  | Parbhani Kranti     | 6.78                | 8.01        | 7.39        | 14.02               | 15.38       | 14.70       | 21.14               | 22.53       | 21.84       |
| T <sub>2</sub>  | Parbhani Bhindi     | 7.31                | 8.54        | 7.92        | 13.27               | 14.63       | 13.95       | 20.42               | 21.79       | 21.10       |
| T <sub>3</sub>  | Arka Anamika        | 8.84                | 10.07       | 9.45        | 17.38               | 16.06       | 18.06       | 24.94               | 26.33       | 25.64       |
| T <sub>4</sub>  | Shahiba             | 7.41                | 8.64        | 8.02        | 14.62               | 15.98       | 15.30       | 22.74               | 24.13       | 23.44       |
| T <sub>5</sub>  | Shan                | 6.84                | 8.07        | 7.45        | 12.40               | 13.76       | 13.08       | 17.54               | 18.93       | 18.24       |
| T <sub>6</sub>  | No. 55              | 7.18                | 8.41        | 7.79        | 13.39               | 14.75       | 14.07       | 21.10               | 22.49       | 21.80       |
| T <sub>7</sub>  | Harita              | 6.44                | 7.67        | 7.05        | 14.14               | 15.5        | 14.82       | 22.54               | 23.93       | 23.24       |
| T <sub>8</sub>  | Jhilmil             | 7.04                | 8.27        | 7.65        | 14.00               | 15.36       | 14.68       | 20.67               | 22.06       | 21.37       |
| T <sub>9</sub>  | No. 64              | 7.71                | 8.94        | 8.32        | 14.70               | 18.74       | 15.38       | 23.25               | 24.64       | 23.95       |
| T <sub>10</sub> | Sonal               | 8.41                | 9.64        | 8.69        | 15.70               | 17.06       | 16.38       | 23.44               | 24.83       | 24.14       |
| T <sub>11</sub> | Shakti              | 7.32                | 8.55        | 7.93        | 14.14               | 15.5        | 14.82       | 22.48               | 23.87       | 23.18       |
| T <sub>12</sub> | Punjab Selection    | 6.58                | 7.81        | 7.19        | 14.92               | 16.28       | 15.60       | 20.3                | 21.69       | 21.00       |
|                 | <b>SEm±</b>         | <b>0.43</b>         | <b>0.47</b> | <b>0.38</b> | <b>0.79</b>         | <b>0.66</b> | <b>0.69</b> | <b>1.00</b>         | <b>1.01</b> | <b>1.06</b> |
|                 | <b>C.D.(p=0.05)</b> | <b>1.28</b>         | <b>1.38</b> | <b>1.13</b> | <b>2.33</b>         | <b>1.95</b> | <b>2.04</b> | <b>2.93</b>         | <b>2.96</b> | <b>3.12</b> |

**Table 3 Performance of different varieties of okra with respect tonumber of branches per plant**

| Treatment       | Varieties           | Number of Branches |             |             | Number of Branches 60 |             |             | Number of Branches 90 |             |             |
|-----------------|---------------------|--------------------|-------------|-------------|-----------------------|-------------|-------------|-----------------------|-------------|-------------|
|                 |                     | 30 DAS             |             |             | DAS                   |             |             | DAS                   |             |             |
|                 |                     | 2022               | 2023        | Pooled      | 2022                  | 2023        | Pooled      | 2022                  | 2023        | Pooled      |
| T <sub>1</sub>  | Parbhani Kranti     | 3.12               | 3.44        | 3.28        | 8.18                  | 8.41        | 8.30        | 10.47                 | 10.66       | 10.57       |
| T <sub>2</sub>  | Parbhani Bhindi     | 2.85               | 3.17        | 3.01        | 7.78                  | 8.01        | 7.90        | 10.83                 | 11.02       | 10.93       |
| T <sub>3</sub>  | Arka Anamika        | 3.85               | 4.17        | 4.01        | 8.68                  | 8.91        | 8.80        | 13.78                 | 13.97       | 13.88       |
| T <sub>4</sub>  | Shahiba             | 2.92               | 3.24        | 3.08        | 8.15                  | 8.38        | 8.27        | 11.05                 | 11.24       | 11.15       |
| T <sub>5</sub>  | Shan                | 3.25               | 3.57        | 3.41        | 6.98                  | 7.21        | 7.10        | 10.12                 | 10.31       | 10.22       |
| T <sub>6</sub>  | No. 55              | 3.38               | 3.70        | 3.54        | 7.84                  | 8.07        | 7.96        | 10.88                 | 11.07       | 10.98       |
| T <sub>7</sub>  | Harita              | 3.18               | 3.50        | 3.34        | 8.12                  | 8.35        | 8.24        | 12.72                 | 12.91       | 12.82       |
| T <sub>8</sub>  | Jhilmil             | 3.12               | 3.44        | 3.28        | 8.11                  | 8.34        | 8.23        | 10.66                 | 10.85       | 10.76       |
| T <sub>9</sub>  | No. 64              | 3.38               | 3.70        | 3.54        | 9.38                  | 9.61        | 9.50        | 11.75                 | 11.94       | 11.85       |
| T <sub>10</sub> | Sonal               | 3.45               | 3.77        | 3.61        | 8.28                  | 8.51        | 8.40        | 12.44                 | 12.63       | 12.54       |
| T <sub>11</sub> | Shakti              | 3.25               | 3.57        | 3.41        | 8.52                  | 8.75        | 8.64        | 13.49                 | 13.68       | 13.59       |
| T <sub>12</sub> | Punjab Selection    | 3.32               | 3.64        | 3.48        | 8.65                  | 8.88        | 8.77        | 13.12                 | 13.31       | 13.22       |
|                 | <b>SEm±</b>         | <b>0.16</b>        | <b>0.24</b> | <b>0.22</b> | <b>0.35</b>           | <b>0.33</b> | <b>0.45</b> | <b>0.83</b>           | <b>0.35</b> | <b>0.46</b> |
|                 | <b>C.D.(p=0.05)</b> | <b>0.49</b>        | <b>0.72</b> | <b>0.65</b> | <b>1.05</b>           | <b>0.97</b> | <b>1.34</b> | <b>2.44</b>           | <b>1.05</b> | <b>1.35</b> |

**Table 4 Performance of different varieties of okra with respect to leaf length (cm) per plant**

| Treatment       | Varieties           | leaf length (cm) 30 DAS |             |             | leaf length (cm) 60 DAS |             |             | leaf length (cm) 90 DAS |             |             |
|-----------------|---------------------|-------------------------|-------------|-------------|-------------------------|-------------|-------------|-------------------------|-------------|-------------|
|                 |                     | 2022                    | 2023        | Pooled      | 2022                    | 2023        | Pooled      | 2022                    | 2023        | Pooled      |
| T <sub>1</sub>  | Parbhani Kranti     | 8.13                    | 8.45        | 8.29        | 14.80                   | 15.05       | 14.92       | 16.19                   | 16.49       | 16.34       |
| T <sub>2</sub>  | Parbhani Bhindi     | 8.33                    | 8.65        | 8.49        | 14.00                   | 14.25       | 14.12       | 16.30                   | 16.60       | 16.45       |
| T <sub>3</sub>  | Arka Anamika        | 10.09                   | 10.41       | 10.25       | 17.09                   | 17.34       | 17.21       | 18.87                   | 19.17       | 19.03       |
| T <sub>4</sub>  | Shahiba             | 9.14                    | 9.46        | 9.30        | 14.94                   | 15.19       | 15.06       | 16.61                   | 16.91       | 16.76       |
| T <sub>5</sub>  | Shan                | 7.53                    | 7.85        | 7.69        | 12.53                   | 12.78       | 12.65       | 14.99                   | 15.29       | 15.14       |
| T <sub>6</sub>  | No. 55              | 8.16                    | 8.48        | 8.32        | 14.12                   | 14.37       | 14.24       | 15.53                   | 15.83       | 15.68       |
| T <sub>7</sub>  | Harita              | 8.99                    | 9.31        | 9.15        | 13.80                   | 14.05       | 13.92       | 16.28                   | 16.58       | 16.44       |
| T <sub>8</sub>  | Jhilmil             | 8.82                    | 9.14        | 8.98        | 12.94                   | 13.19       | 13.06       | 15.83                   | 16.13       | 15.98       |
| T <sub>9</sub>  | No. 64              | 9.40                    | 9.72        | 9.56        | 15.26                   | 15.51       | 15.38       | 16.91                   | 17.21       | 17.06       |
| T <sub>10</sub> | Sonal               | 9.81                    | 10.13       | 9.97        | 16.08                   | 16.33       | 16.20       | 17.47                   | 17.77       | 17.62       |
| T <sub>11</sub> | Shakti              | 8.48                    | 8.80        | 8.64        | 12.70                   | 12.95       | 12.82       | 15.02                   | 15.32       | 15.17       |
| T <sub>12</sub> | Punjab Selection    | 7.88                    | 8.20        | 8.04        | 11.68                   | 11.93       | 11.80       | 13.82                   | 14.12       | 13.97       |
|                 | <b>SEm±</b>         | <b>0.50</b>             | <b>0.44</b> | <b>0.50</b> | <b>0.71</b>             | <b>0.47</b> | <b>0.59</b> | <b>0.83</b>             | <b>0.45</b> | <b>0.58</b> |
|                 | <b>C.D.(p=0.05)</b> | <b>1.46</b>             | <b>1.31</b> | <b>1.44</b> | <b>2.10</b>             | <b>1.39</b> | <b>1.72</b> | <b>2.44</b>             | <b>1.32</b> | <b>1.69</b> |

**Table .5 Performance of different varieties of okra with respect to leaf width (cm) per plant**

| Treatment       | Varieties           | leaf width (cm) 30 DAS |             |             | leaf width (cm) 60 DAS |             |             | leaf width (cm) 90 DAS |             |             |
|-----------------|---------------------|------------------------|-------------|-------------|------------------------|-------------|-------------|------------------------|-------------|-------------|
|                 |                     | 2022                   | 2023        | Pooled      | 2022                   | 2023        | Pooled      | 2022                   | 2023        | Pooled      |
| T <sub>1</sub>  | Parbhani Kranti     | 9.64                   | 9.96        | 9.80        | 19.67                  | 19.92       | 19.79       | 21.51                  | 21.66       | 21.51       |
| T <sub>2</sub>  | Parbhani Bhindi     | 9.45                   | 9.77        | 9.61        | 18.65                  | 18.90       | 18.77       | 20.97                  | 21.12       | 20.97       |
| T <sub>3</sub>  | Arka Anamika        | 12.10                  | 12.42       | 12.26       | 22.59                  | 22.84       | 22.71       | 25.32                  | 25.47       | 25.32       |
| T <sub>4</sub>  | Shahiba             | 10.08                  | 10.40       | 10.24       | 20.31                  | 20.56       | 20.43       | 22.27                  | 22.42       | 22.27       |
| T <sub>5</sub>  | Shan                | 8.51                   | 8.83        | 8.67        | 16.97                  | 17.22       | 17.09       | 18.75                  | 18.9        | 18.75       |
| T <sub>6</sub>  | No. 55              | 9.01                   | 9.33        | 9.17        | 18.52                  | 18.77       | 18.64       | 20.14                  | 20.29       | 20.14       |
| T <sub>7</sub>  | Harita              | 9.74                   | 10.06       | 9.90        | 18.87                  | 19.12       | 18.99       | 21.38                  | 21.53       | 21.38       |
| T <sub>8</sub>  | Jhilmil             | 9.53                   | 9.85        | 9.69        | 17.76                  | 18.01       | 17.88       | 21.81                  | 21.96       | 21.81       |
| T <sub>9</sub>  | No. 64              | 10.34                  | 10.66       | 10.50       | 20.84                  | 21.09       | 20.96       | 23.41                  | 23.56       | 23.41       |
| T <sub>10</sub> | Sonal               | 10.93                  | 11.25       | 11.09       | 21.53                  | 21.78       | 21.65       | 23.98                  | 24.13       | 23.98       |
| T <sub>11</sub> | Shakti              | 9.04                   | 9.36        | 9.20        | 19.28                  | 19.53       | 19.40       | 22.04                  | 22.19       | 22.04       |
| T <sub>12</sub> | Punjab Selection    | 8.11                   | 8.43        | 8.27        | 16.84                  | 17.09       | 16.96       | 19.54                  | 19.69       | 19.54       |
|                 | <b>SEm±</b>         | <b>0.50</b>            | <b>0.51</b> | <b>0.68</b> | <b>1.04</b>            | <b>0.57</b> | <b>0.62</b> | <b>0.67</b>            | <b>0.83</b> | <b>0.70</b> |
|                 | <b>C.D.(p=0.05)</b> | <b>1.46</b>            | <b>1.50</b> | <b>1.98</b> | <b>3.06</b>            | <b>1.68</b> | <b>1.83</b> | <b>1.95</b>            | <b>2.44</b> | <b>2.00</b> |

**Table .6 Performance of different varieties of okra with respect to Leaf Area (cm<sup>2</sup>) per plant**

| Treatment       | Varieties           | Leaf Area (cm <sup>2</sup> ) 30 DAS |             |             | Leaf Area (cm <sup>2</sup> ) 60 DAS |             |             | Leaf Area (cm <sup>2</sup> ) 90 DAS |             |             |
|-----------------|---------------------|-------------------------------------|-------------|-------------|-------------------------------------|-------------|-------------|-------------------------------------|-------------|-------------|
|                 |                     | 2022                                | 2023        | Pooled      | 2022                                | 2023        | Pooled      | 2022                                | 2023        | Pooled      |
| T <sub>1</sub>  | Parbhani Kranti     | 149.14                              | 150.46      | 149.80      | 184.37                              | 186.29      | 185.33      | 220.96                              | 223.92      | 222.44      |
| T <sub>2</sub>  | Parbhani Bhindi     | 156.27                              | 157.59      | 156.93      | 191.39                              | 193.31      | 192.35      | 226.05                              | 229.01      | 227.53      |
| T <sub>3</sub>  | Arka Anamika        | 172.06                              | 173.38      | 172.72      | 223.53                              | 225.45      | 224.49      | 248.56                              | 251.52      | 250.04      |
| T <sub>4</sub>  | Shahiba             | 161.96                              | 163.28      | 162.62      | 199.28                              | 201.20      | 200.24      | 230.63                              | 233.59      | 232.11      |
| T <sub>5</sub>  | Shan                | 118.32                              | 119.64      | 118.98      | 167.54                              | 169.46      | 168.50      | 198.49                              | 201.45      | 199.97      |
| T <sub>6</sub>  | No. 55              | 123.39                              | 124.71      | 124.05      | 173.14                              | 175.06      | 174.10      | 205.73                              | 208.69      | 207.21      |
| T <sub>7</sub>  | Harita              | 134.60                              | 135.92      | 135.26      | 177.46                              | 179.38      | 178.42      | 211.61                              | 214.57      | 213.09      |
| T <sub>8</sub>  | Jhilmil             | 144.10                              | 145.42      | 144.76      | 179.65                              | 181.57      | 180.61      | 216.07                              | 219.03      | 217.55      |
| T <sub>9</sub>  | No. 64              | 164.57                              | 165.89      | 165.23      | 208.69                              | 210.61      | 209.65      | 235.07                              | 238.03      | 236.55      |
| T <sub>10</sub> | Sonal               | 168.72                              | 170.04      | 169.38      | 217.32                              | 219.24      | 218.28      | 240.18                              | 243.14      | 241.66      |
| T <sub>11</sub> | Shakti              | 137.38                              | 138.70      | 138.04      | 191.26                              | 193.18      | 192.22      | 204.56                              | 207.52      | 206.04      |
| T <sub>12</sub> | Punjab Selection    | 127.13                              | 128.62      | 127.96      | 182.08                              | 184.00      | 183.04      | 203.41                              | 206.37      | 204.89      |
|                 | <b>SEm±</b>         | <b>2.21</b>                         | <b>2.24</b> | <b>2.55</b> | <b>2.41</b>                         | <b>2.47</b> | <b>2.55</b> | <b>1.66</b>                         | <b>2.41</b> | <b>1.90</b> |
|                 | <b>C.D.(p=0.05)</b> | <b>6.51</b>                         | <b>6.60</b> | <b>7.49</b> | <b>7.06</b>                         | <b>7.21</b> | <b>7.50</b> | <b>4.88</b>                         | <b>7.06</b> | <b>5.50</b> |

**Table 7 Performance of different varieties of okra with respect to phonological characteristics**

| Treatment       | Varieties           | Days Taken to 50% Flowering |             |             | Days taken to flower opening to fruit picking |             |             | Days Taken to First Picking |             |             |
|-----------------|---------------------|-----------------------------|-------------|-------------|---|-------------|-------------|-----------------------------|-------------|-------------|
|                 |                     | 2022                        | 2023        | Pooled      | 2022  | 2023        | Pooled      | 2022                        | 2023        | Pooled      |
| T <sub>1</sub>  | Parbhani Kranti     | 44.39                       | 43.07       | 43.73       | 7.66  | 7.36        | 7.51        | 51.47                       | 49.55       | 50.51       |
| T <sub>2</sub>  | Parbhani Bhindi     | 46.66                       | 45.34       | 46.00       | 8.66  | 8.36        | 8.51        | 52.58                       | 50.66       | 51.62       |
| T <sub>3</sub>  | Arka Anamika        | 43.99                       | 42.67       | 43.33       | 6.66  | 6.36        | 6.51        | 50.22                       | 48.30       | 49.26       |
| T <sub>4</sub>  | Shahiba             | 44.79                       | 43.47       | 44.13       | 7.04  | 6.74        | 6.89        | 50.87                       | 48.95       | 49.91       |
| T <sub>5</sub>  | Shan                | 47.94                       | 46.62       | 47.28       | 9.66  | 9.36        | 9.51        | 54.98                       | 53.06       | 54.02       |
| T <sub>6</sub>  | No. 55              | 46.36                       | 45.04       | 45.70       | 9.31  | 9.01        | 9.17        | 53.26                       | 51.34       | 52.30       |
| T <sub>7</sub>  | Harita              | 44.97                       | 43.65       | 44.31       | 7.93  | 7.63        | 7.78        | 51.45                       | 49.53       | 50.49       |
| T <sub>8</sub>  | Jhilmil             | 45.42                       | 44.10       | 44.76       | 7.33  | 7.03        | 7.19        | 51.52                       | 49.60       | 50.56       |
| T <sub>9</sub>  | No. 64              | 45.83                       | 44.51       | 45.17       | 6.87  | 6.57        | 6.72        | 52.08                       | 50.16       | 51.12       |
| T <sub>10</sub> | Sonal               | 43.99                       | 42.67       | 43.34       | 6.64  | 6.34        | 6.49        | 51.37                       | 49.45       | 50.41       |
| T <sub>11</sub> | Shakti              | 44.15                       | 42.49       | 43.15       | 5.92  | 5.62        | 5.77        | 49.49                       | 47.57       | 48.53       |
| T <sub>12</sub> | Punjab Selection    | 42.93                       | 41.61       | 42.27       | 5.66  | 5.36        | 5.51        | 48.72                       | 46.80       | 47.76       |
|                 | <b>SEm±</b>         | <b>0.80</b>                 | <b>0.83</b> | <b>0.67</b> | <b>0.61</b>                                   | <b>0.33</b> | <b>0.62</b> | <b>1.09</b>                 | <b>1.67</b> | <b>0.62</b> |
|                 | <b>C.D.(p=0.05)</b> | <b>2.35</b>                 | <b>2.44</b> | <b>1.95</b> | <b>1.80</b>                                   | <b>0.98</b> | <b>1.83</b> | <b>3.20</b>                 | <b>4.89</b> | <b>0.89</b> |