

Johnson Grass (*Sorghum halepense*) Interference, its Effect on Crops Yield and Soil attributes under Different Cropping Systems and Management Practices

ABSTRACT

Johnson grass (*Sorghum halepense*) is one of the most detrimental and toxic weeds among the weed species. Its environment is expanding unceasingly due to deeply expanding of its rhizomes in the soil, self-pollinating reproduction strategy, accelerated growth and poor management which results in production losses of major agronomic crops. Mungbean and sorghum are important crops grown on significant acreage around the world, and a major constraint for their production losses is weeds interference. Thus, field studies were conducted to examine Johnson grass interference, control, and recovery under different management practices and cropping system and its effects on crops production and soil health. Our results indicated that the Johnson grass density was minimized by the application of plastic sheet mulch treatment under wheat-mungbean cropping system. Johnson grass competition had a significant impact on growth and grain yield of both mungbean and sorghum. The crop yield parameters; 100 grain weight and grain yield were lowest in the non-treated treatment (control), while the greatest values of these parameters were recorded in the plastic sheet mulch and wheat straw mulch treatments with wheat-mungbean cropping system. Similarly, soil building attributes; available nitrogen, available phosphorus, available potassium and organic matter content were positively affected by the interaction of plastic sheet mulch + wheat-mungbean cropping system. Overall, we have concluded that wheat-mungbean and wheat-sorghum are most resilient cropping system with the implementation of plastic sheet mulch and wheat straw mulch to suppress the spread of Johnson grass.

Keywords: Johnson grass, crop yield, soil health, cropping systems, legumes, cereals

Introduction

Johnson grass (*Sorghum halepense*) is the world's most precarious and persistent weed that is C4 perennial graminoid plant species from the Poaceae family (Holm *et al.* 1997). Agricultural land and natural biodiversity have been severely deprived by the spread of Johnson grass across all over the globe; Asia, Africa, America, and Europe, covering a one third of the world's field area (Chirita *et al.* 2007). It is ranked sixth among the worst weeds in whole world that has infested 30 different crops in 53 countries and invading millions of hectares (Valverde & Gressel 2006). It was introduced as a perennial fodder crop, but its invasive and tenacious nature has made it a nuisance to agricultural productivity (Hoffman & Buhler, 2002; Binimelis *et al.* 2009). Because of its high core competencies and allelopathic potential, *Sorghum halepense* is eminent for having a detrimental impact on neighboring plant growth and development (Novak *et al.* 2009; Huang *et al.* 2015). In cultivated regions, it has resulted in significant yield losses in economically important crops such as wheat, soybean, maize, cotton, vegetables, and fruits (Uludag *et al.* 2007; Uremis *et al.* 2009). When the cyanide content of weed is high, grazing on *S. halepense* causes harm to cattle, sheep, and horses during winter (Henderson, 2001). Because of its multiple ways of propagation, fast-growing behavior and resilience to extreme climatic fluctuations, *S. halepense* may thrive in a range of locations and ecological niches (Mihovsky & Pachev 2012; Vila-Aiub *et al.* 2013). These biological characteristics of *S. halepense* have

contributed to its reputation as a difficult-to-control weed, as well as impacting the effectiveness of intercultural operations to control *S. halepense* in diverse crops (Dalley & Richard 2008; Heap 2014).

Cropping systems are described as the pattern of crops that are grown throughout the large area, as well as the methods used to cultivate crops (Blanco-Canqui & Lal, 2010). The sequence of crops that are grown on larger areas to get maximum benefits are referring to cropping system and for successful completion of crop production is improved by management practices. Different managerial measures, including as tillage, crop residue management, cropping sequence, nutrients, irrigation, and erosion control, are necessary for the effective and efficient development of crops in a given cropping system (Blanco-Canqui & Lal, 2010). The agricultural system management has a positive influence on soil and water conservation. Soil fertility is improved, soil erosion is reduced, and soil properties are improved with a well-balanced farming approach. Cropping methods with poor management, on the other hand, lead to decreased soil fertility and increased erosion. Mono cropping has a number of negative implications, including poor soil qualities, increased fertilizer and pesticide usage, weed and insect infestations, and decreased crop yields (Blanco-Canqui & Lal, 2010).

Mulches have been used for vegetable cultivation since ancient times, and it got its origin from word ‘molsch’, a German word which has meaning "easy to decay" (Lightfoot 1994). This practice of covering soil to intercept the germination of weeds and loss of moisture which optimizes crop production by spreading of different covering material is termed as mulches (Nalayini 2007; Kader *et al.* 2019). They not only improve the yield production but also nourish the soil, ameliorate the soil penetration, diminish the runoff, reduce the rate of evapotranspiration and restrict the weeds emergence to large extent (Rathore *et al.* 1998). They have variety of important environmental benefits including soil and plant root temperature regulation, decreased nutrient losses, reduced soil erosion and compaction, and improved physical soil conditions (Ngouajio & McGiffen 2004; Lamont 2005).

Mulching works as a barrier to light penetration under the surface, preventing weeds from completing photosynthetic activities. The efficient way of getting rid of annual weeds is to use the mulch (Ahmad *et al.* 2015; Ahmad *et al.* 2020). The emergence and growth of weeds is inhibited by use of mulches and other cultural practices that have been used to overcome weeds are minimized due to their effectiveness. The most effectual mulch that is applied to control the spread of weeds in cotton is plastic sheet. The weed density can be significantly decreased by cumulative quantity of mulch (Ahmad *et al.* 2015; Ahmad *et al.* 2020)

Materials and Methods

Experimental Location and Treatments

The experiment was conducted in field on University Research Farm, Koont, PMAS Arid Agriculture University, Rawalpindi in growing year 2021–2022. The climate of URF, Koont, Rawalpindi is arid to semi-arid. The metrological data gathered throughout the experiment is depicted in Figure 1.

Three cropping systems (CS) were used in this experiment: wheat-sorghum (CS₁), wheat-mungbean (CS₂) and wheat-fallow (CS₃). Four weed control techniques were factorial paired

with the CS: control (C), deep ploughing (DP), wheat straw mulch (WSM), and plastic sheet mulch (PSM). The experiment was set up in a split-plot arrangement. Cropping systems were arranged in main plots while sub plots were consisted of management practices. This experiment had three replications and total plot size was 30m x 20m. During the wheat growing season, weeds were physically eradicated upon their emergence while in summer season Johnson grass was controlled by applying different management strategies. In control treatment, nothing was applied which allowed the weed to emerge and grow for the comparison with other treatments while in DP treatment, deep ploughing was done before sowing in all the main plots including fallow where nothing was grown in both years. Wheat straw mulch (WSM) was spread manually on the soil after land preparation and before sowing of the mungbean and sorghum. Plastic sheet mulch (PSM) was applied on soil between lines 15 days after sowing mungbean and sorghum while it was applied immediately after land preparation in fallow plot.

Crop Husbandry

The experimental area received 10 cm of pre-soaking irrigation during both seasons. For the preparation of seedbed, soil had to achieve field capacity. Table 1 shows the crops that were grown according to the suggested production technology for the area. With a hand drill, all the crops were manually seeded in lines. The irrigation was completely dependent on precipitation. Urea and di-ammonium phosphate were used as fertilizers (DAP). All the phosphorus was added to soil at time of sowing and 1/3rd portion of nitrogen was applied at sowing time. The rest of the nitrogen was applied during the first and second irrigations, respectively. In cropping season, diseases, insects, and pests were managed by implementing the necessary agronomic and crop protection methods. The grain yield from each plot of each crop was estimated when moisture level was about 12%.

Table 1 Crop husbandry of different crops encompassed in various cropping systems (2021–2022).

Crops	Sowing Time	Varieties	Seed Rate (kg/ha)	Fertilizer NPK (kg/ha)	P-P (cm)	R-R (cm)	Harvesting Time	Harvest Method
Wheat	20 Nov	Barani-2017	125	60-75-0	-	25	09 April	Manual
Mungbean	13 April	NM-2011	75	20-60-0	10	30	08 Sep	Manual
Sorghum	13 April	DS-2003	20	100-60-0	15	60	13 Sep	Manual

P–P = Plant spacing; R–R = Row spacing

Weed Parameters of Johnson grass

Plant density (plants per m²), plant height (cm), plant fresh weight (g) and plant dry weight (g) were the attributes to determine for Johnson grass. The whole data was calculated from plots of mungbean, sorghum and fallow. Weed density was calculated from 1m² sample during the booting stage of Johnson grass and plant height was estimated from 10 randomly selected plants. To find out the fresh weights, we had averaged the fresh weights of 10 selected plants for each plot and then dried these samples in oven at 120 Celsius for 24 hours to determine the dry weights with analytical balance.

Agronomic Traits of Mungbean and Sorghum

The agronomic and yield parameters of mungbean and sorghum were taken from each experimental unit. It includes plant height (cm), 100 grain weight (g), straw yield (kg/ha) and grain yield (kg/ha) for mungbean. Similarly, plant height (cm), 100 grain weight (g), straw yield (kg/ha) and grain yield (kg/ha) was the traits for sorghum, Plant height was taken from 10 random plants from each plot. For calculating 100 grain weight, we had averaged the 10 samples from each experimental unit containing 100 grain weight. Straw yield was determined by harvesting plants from area of 1m², dried these samples in oven at 120 Celsius for 24 hours and then weighted. Similarly, plants from 1m² were harvested and then threshed to calculate grain yield of wheat.

Soil Analysis

Composite soil samples (0–15 cm depth) were obtained from each experimental unit for postharvest soil analysis of Available N, P, K and soil organic content (SOM). To find out the value of soil organic matter content we will multiply organic carbon (OC) by 1.73 while Walkley and Black method was followed to determine OC titrimetrically. Kjeldahl technique was used to calculate available nitrogen. To find the available phosphorus content, soil was agitated with a 0.03 M NH₄F— 0.025 M HCl solution at a pH of less than 7.0. By employing the ammonium bicarbonate-DTPA method (AB-DTPA), the value of available K was evaluated.

Statistical Analysis

Utilizing Statistical Software, IBM SPSS and R, data were statistically examined in order to identify substantial variability among various parameters. The least significant difference test was used to compare the means.

Results and Discussion

Soil Properties

During the season of study, different management practices (MP) and cropping systems (CS) had a substantial influence on available nitrogen. PSM had the greatest nitrogen value among management methods, whereas C had the lowest. In cropping systems (CS), CS₂ had the highest value of plant available nitrogen while cropping system CS₃ had the lowest value.





Figure 1a, b: Effect of numerous management practices (a) and cropping systems (b) on available nitrogen (mg/kg) C = Control, DP = Deep Ploughing, WSM = Wheat straw mulch, CS₁ = Wheat-Sorghum, CS₂ = Wheat-Mungbean, CS₃= Wheat-Fallow. Any two means sharing different letters are statistically different ($p \leq 0.05$) from each other.

During the two years of the study, MP and CS considerably changed the amount of P that was accessible by the plant in soil. Among the various management practices included in the study, it varied from 7.63 to 19.40 mg/kg. The PSM and WSM produced the greatest values of available P, whereas C has the minimum value of available P. Similar to this, among the several CS included in the study, available P varied from 5.90 to 19.4 mg/kg. The minimum and maximum values for available P were obtained for C₂ and C₃ cropping systems, respectively (Figure 2a, b).





Figure 2a, b: Effect of numerous management practices (a) and cropping systems (b) on available phosphorus (mg/kg) C = Control, DP = Deep Ploughing, WSM = Wheat straw mulch, CS₁ = Wheat-Sorghum, CS₂ = Wheat-Mungbean, CS₃= Wheat-Fallow. Any two means sharing different letters are statistically different ($p \leq 0.05$) from each other

Different management practices (MP) and cropping systems (CS) had a substantial influence on plant available potassium. PSM had the greatest potassium level among management treatments, whereas C had the lowest. Cropping system CS₂ had greatest value of potassium; however the CS₃ cropping system had the lowermost value (Figure 4). Similarly, available K ranged among the several CS included in the study from 121.43 to 185.81 mg/kg. The C₂ and C₃ cropping systems had estimated with maximum and minimum values for available K through the time of the experiment, respectively (Figure 3a, b)





Figure 3a, b: Effect of numerous management practices (a) and cropping systems (b) on available potassium (mg/kg) C = Control, DP = Deep Ploughing, WSM = Wheat straw mulch, CS₁ = Wheat-Sorghum, CS₂ = Wheat-Mungbean, CS₃= Wheat-Fallow. Any two means sharing different letters are statistically different ($p \leq 0.05$) from each other

Various management techniques (MP) and cropping systems (CS) both had a substantial influence on soil organic matter. In terms of management techniques, PSM recorded the greatest value for soil organic matter, whereas C recorded the lowest. The CS₂ cropping system had the highest value of soil organic matter, whereas the CS₃ cropping system had the lowest value.




Figure 4a, b: Effect of numerous management practices (a) and cropping systems (b) on soil organic matter (%) C = Control, DP = Deep Ploughing, WSM = Wheat straw mulch, CS₁ = Wheat-Sorghum, CS₂ = Wheat-Mungbean, CS₃= Wheat-Fallow. Any two means sharing different letters are statistically different ($p \leq 0.05$) from each other

Weed Parameters of Johnson grass

Plant Density (plants per m²)

Various management treatments and cropping system had significant effect on plant density of Johnson grass. However, lowest plant density was recorded in PSM which was followed by WSM while highest plant density was noted in C treatment followed by DP (Table 2). Among different cropping systems, maximum number of plants per m² was recorded in CS₃ while least possible number of plants in m² was experiential in CS₁ throughout the growing season (Table 2).

The interactive effect of cropping systems and management treatments in both experimental years drastically influenced plant density of observed Johnson grass. Plant density ranged from 43 to 7 plants per m² (Table 2). During the year of the research, all CS had the lowest plant density of Johnson grass with PSM, followed by WSM (Table 6). All CS had the maximum plant density of Johnson grass with C treatment, whereas CS₁ had the lowest (Table

2). CS₃ was the most afflicted cropping system overall, however CS₁ was slightly affected cropping system in this experiment.

Table 2: Effect of different cropping systems and management practices on plant density (plants m⁻²) of Johnson grass

2022				
	CS ₁	CS ₂	CS ₃	Mean
C	22.66d	30.33bc	43.00a	32.00a
DP	13.67ef	22.00d	32.67b	22.78b
WSM	9.33fg	14.00ef	26.67cd	16.67c
PSM	7.00g	12.67ef	16.00e	11.89d
Mean	13.16c	19.75b	29.58a	

The lowercase letters denote how respective cropping system differed among various management practices. C = Control, DP = Deep Ploughing, WSM = Wheat straw mulch, PSM = Plastic Sheet Mulch, CS₁=wheat-sorghum, CS₂=wheat-mungbean, CS₃= wheat-fallow, CS = cropping-systems and T= management treatments

Plant Height (cm)

The interactive effect of cropping systems and management treatments in the experimental year considerably influenced plant height of observed Johnson grass. The plant height varied from 223.76 to 73.17 cm (Table 3). All CS measured the least plant height of Johnson grass using PSM, which was then followed by WSM (Table 7). All CS recorded the optimal plant height of Johnson grass with C treatment, but the CS₁ recorded the minimum value (Table 3). Consequently, CS₃ had the highest plant height across all cropping systems, whereas CS₁ had the lowest plant height throughout all cropping systems.

Table 3: Effect of different cropping systems and management practices on plant height (cm) of Johnson grass

2022				
	CS ₁	CS ₂	CS ₃	Mean
C	194.42bc	217.64ab	223.74a	211.93a
DP	167.15d	185.08cd	200.06abc	184.10b
WSM	96.62fg	139.15e	168.91d	134.9c
PSM	74.60g	105.13f	73.17g	84.30d
Mean	133.20b	161.75a	166.47a	

The lowercase letters denote how respective cropping system differed among various management practices. C = Control, DP = Deep Ploughing, WSM = Wheat straw mulch, PSM = Plastic Sheet Mulch, CS₁=wheat-sorghum, CS₂=wheat-mungbean, CS₃= wheat-fallow, CS = cropping-systems and T= management treatments

Plant Fresh Weight (kg/ha)

Data regarding plant fresh weight is presented in Table 4. Plant fresh weight was not found statistically different for different CS but found highly significant for different treatment.

In management treatments (T), minimum plant fresh weight was recorded for PSM while maximum plant fresh weight was recorded for C followed by DP. In cropping systems (CS), lowest plant fresh weight was estimated in CS₁ while maximum fresh weight was observed in CS₃ which was statistically similar to CS₂ (Table 4).

The interactive effect of cropping systems (CS) and management treatments (T) profoundly influenced plant fresh weight of observed Johnson grass. The fresh weight of the plant varied from 441.73 to 132.44 g (Table 4). All CS measured the minimum plant height of Johnson grass using PSM, which was then followed by WSM (Table 8). All CS recorded the optimal plant height of Johnson grass with C treatment, but the CS₁ recorded the lowest values (Table 4). Furthermore, CS₃ had the highest plant height among all cropping systems, while CS₁ had the lowest plant height across all cropping systems.

Table 4: Effect of different cropping systems and management practices on plant fresh weight (g) of Johnson grass

2022				
	CS₁	CS₂	CS₃	Mean
C	403.90ab	422.88ab	441.73a	422.84a
DP	342.04c	355.54c	389.08bc	362.22b
WSM	168.46ef	258.29d	258.00d	228.25c
PSM	132.44f	199.29e	194.89e	175.54d
Mean	261.71b	309.00a	320.92a	

The lowercase letters denote how respective cropping system differed among various management practices. C = Control, DP = Deep Ploughing, WSM = Wheat straw mulchs, PSM = Plastic Sheet Mulch, CS₁=wheat-sorghum, CS₂=wheat-mungbean, CS₃= wheat-fallow, CS = cropping-systems and T= management treatments

Plant Dry Weight (cm)

Data regarding plant dry weight is presented in Table 5. Plant dry weight was found statistically different for both different CS and for different management treatments. In management treatments (T), minimum plant dry weight was recorded for PSM while maximum plant dry weight was recorded for (C) followed by deep ploughing (DP). In cropping systems (CS), lowest plant dry weight was estimated in CS₁ while maximum fresh weight was observed in CS₃ (Table 5).

Plant dry weight was changed drastically during both study years due to the interaction between cropping systems (CS) and management treatments (T). The plant dry weight varied from 93.49 to 11.53 g, respectively (Table 5). During this period of the study, PSM was used to record the minimum plant height of Johnson grass, followed by WSM (Table 5). CS₁ was reported with lowest value of plant dry weight, although all CS recorded the optimal plant height of Johnson grass with C treatment (Table 5). In general, CS₃ had the highest average plant dry weight across all cropping systems, whereas CS₁ had the lowest average plant dry weight across all cropping systems.

Table 5: Effect of different cropping systems and management practices on plant dry weight (g) of Johnson grass

2022				
	CS ₁	CS ₂	CS ₃	Mean
C	67.11c	80.96ab	93.49a	80.52a
DP	62.12c	71.45bc	85.19a	72.92a
WSM	26.48ef	36.71de	41.10d	34.76b
PSM	11.53g	14.89fg	17.55fg	14.65c
Mean	41.81c	51.0b	59.33a	

The lowercase letters denote how respective cropping system differed among various management practices. C = Control, DP = Deep Ploughing, WSM = Wheat straw mulchs, PSM = Plastic Sheet Mulch, CS₁=wheat-sorghum, CS₂=wheat-mungbean, CS₃= wheat-fallow, CS = cropping-systems and T= management treatments.

Agronomic Parameters of Mungbean

Plant height (cm)

Data on the subject of plant height of mungbean as prejudiced by management practices is presented in Fig. 5. Analysis of variance (ANOVA) of the data revealed that management practices significantly affected the plant height of mungbean. Plots sown where plastic sheet mulch treatment was applied produced plants with maximum plant height of 58.46 cm, while minimum plant height (37.17 cm) was observed in control where nothing was applied as management practices.




Figure 5: Plant Height (cm) of mungbean as affected by the various management treatments. C = Control, DP= Deep Ploughing, WSM= Wheat straw mulchs, PSM= Plastic Sheet Mulch. Error bars indicate the LSD 5% values.

100 grain weight (g)

Management techniques considerably changed the 100 grain weight. Different management practices considered in the study varied the 100 grain weight from 5.02 to 3.26 g in the growing year. The greatest values of 100 grain weight were obtained from the PSM (5.02 g)

and WSM (4.08 g), whilst the lowest values were obtained from the control (C) (3.26 g) (Figure 6).




Figure 6: 100 grain weight (g) of mungbean as affected by the various management treatments. C = Control, DP= Deep Ploughing, WSM= Wheat straw mulchs, PSM= Plastic Sheet Mulch. Error bars indicate the LSD5% values.

Grain yield (kg/ha)

In both research years, grain yield varied dramatically depending on management strategies. Among the various management approaches, grain yield varied between 1610.7 and 810.27 kg/ha. The maximum grain yield was produced by the PSM (1610.17 g) and the WSM (1150.52 kg/ha), whereas the C (810.27 kg/ha) produced the lowest average grain yield (Figure 7).




Figure 7: Grain yield (kg/ha) of mungbean as affected by the various management treatments. C = Control, DP= Deep Ploughing, WSM= Wheat straw mulchs, PSM= Plastic Sheet Mulch. Error bars indicate the LSD 5% values

Agronomic Traits of Sorghum

Plant height (cm)

Data regarding plant height of mungbean as influenced by management practices are presented in Fig. 8. Analysis of variance (ANOVA) of the data revealed that management practices significantly affected the plant height of mungbean. Plots sown where WSM treatment was applied produced plants with maximum plant height of 229.96 cm while minimum plant height 208.46 cm was found in case of control where nothing was applied as management practices.




Figure 8: Plant Height (cm) of sorghum as affected by the various management treatments. C = Control, DP= Deep Ploughing, WSM= Wheat straw mulchs, PSM= Plastic Sheet Mulch. Error bars indicate the LSD 5% values.

100 grain weight (g)

Analysis of variance (ANOVA) regarding 100 grain weight showed that management treatments were significant. Maximum 100 grain weight was 3.81 g recorded in PSM during. It was followed by PSM having 100 grain weight 3.55 g. Furthermore, the 100-grain weight of 2.70 g were found in control (C) during the period of experiment (Fig.9)




Figure 9: 100 grain weight (g) of sorghum as affected by the various management treatments. C = Control, DP= Deep Ploughing, WSM= Wheat straw mulchs, PSM= Plastic Sheet Mulch. Error bars indicate the LSD5% values.

Grain yield (kg/ha)

Grain yield varied dramatically depending on management strategies. Among the various management approaches examined in the research, it varied between 2214.6 kg/ha between the 4061 and 2555.18 kg/ha during the growing season. The PSM and WSM produced the greatest grain yields, (3595.99 and 4061 kg/ha), whereas control (C) (2555.18 kg/ha) produced the lowest grain yield (Figure 10).




Figure 10: Grain yield (kg/ha) of sorghum as affected by the various management treatments. C = Control, DP= Deep Ploughing, WSM= Wheat straw mulchs, PSM= Plastic Sheet Mulch. Error bars indicate the LSD 5% values.

Discussion

Shahzad *et al.*, (2021) recorded that plant height of major weeds like Johnson grass *Araganthus* spp. and *Partuala* spp. has been suppressed when emerged in sorghum fields. Farooq *et al.* (2020) showed that phenolic compounds present in sorghum played pivot role in suppression of plant height of these noxious weeds. Our outcomes are in proximity with the results of Azadbakht *et al.*, (2017) and Asif *et al.*, (2020) who found that weed density can be reduced by using plastic sheet mulch on soil surface during sowing. Naeem *et al.* (2021) recorded from study that sorghum-based cropping system with different weed management strategies diminished the plant density of Johnson grass. They also found that allelopathic effect of sorghum resulted in minimum. Our results are in agreement with those of Awal *et al.*, (2016). They found that growth of Johnson grass and other seasonal weeds can be abridged by using plastic sheet mulch in field during sowing of crops. However, David & Chandler, (1987) reported that good to excellent control of Johnson grass was observed for applying herbicide at 8 cm and 16 cm tall Johnson grass.

Azad *et al.*, (2015) found that application of sheet mulch diminishes the fresh weight of Johnson grass and other summer season weeds. While Mistaskas *et al.*, (2003) found that fresh weight of Johnson grass was diminished by the early application herbicides and its fresh weight increased with greater duration of time. Similarly, Czarnota *et al.* (2003) found that sorghum excreted allelopathic chemicals like hydrophobic compound (e.g. *sargoleone*), phenolic acids that inhibited the growth of weeds and diminished the fresh and dry weights of plants Patil *et al.* (2013) recorded that there was significant decrease in dry weight of weeds by the application of plastic sheet mulch as management practices. Czarnota *et al.* (2003) reported that sorghum excreted allelopathic chemicals like hydrophobic compound (e.g. *sargoleone*), phenolic acids that inhibited the growth of weeds and diminished the dry weight of plant. On the other hand, Ibrahim *et al.* (2013) also reported similar results that allelopathic extract put negative affect on weeds which suppressed the plant dry of weeds.

Yaqub & Shahzad (2009) indicated a gradual increase in plant height of mungbean when applied with plastic sheet due to solarization which results in reduction of attack of pests and emergence of weeds. Islam & Faruq (2009) also recorded the similar results of increased plant height in mungbean by the application of different plastic sheets. Agele *et al.* (2010) also observed that plastic mulches maximize the plant height of crops and Mohtisham *et al.* (2013) revealed that plastic sheet mulch improves the plant height to larger extent in rice crop. Similar findings were also reported by Kumar *et al.* (2008) that plant height significantly increased by the application of wheat straw mulch on sorghum. Similarly, Singh *et al.* (2014) also suggested when wheat straw mulch was applied to the crop resulted in the improvement of plant height due to early suppression of pests and weeds and matching results were also found by Khairnar *et al.* (2014) during the application of wheat straw mulch on crop under observation.

Burnside & Wicks (1965) had found that application of plastic sheet mulch also put substantial difference on the 100-grain weight of sorghum but El- Samnoudiet *al.* (2009) demonstrating a significant difference in 100 grain weight with the application of mulches during sowing the crop which was according to the results shown in Fig. 15. Additionally, Abdelrahman *et al.* (2016) also found that applying sheet mulch increases the 100-grain weight in almost all

crops. The results of Mohtisham *et al.* (2013) are in line with our research, showing the significant difference among the various treatments in 100 grain weight of crop. 100 grain weight was very much enhanced by using sheet mulch. Similar results were also demonstrated by Xu *et al.* (2007) where mulches improved the 100 grain weight of crop.

Unger & Jones (1981) reported an increase in grain yield and quality of sorghum with the application of sheet mulch as it enhanced the water use efficiency and nutrient uptake. El-Samnoudi *et al.* (2019) also showed that soil mulches increase the grain yield of sorghum to larger degree due to improved soil structure and proper addition of nutrients to soil.

Karim *et al.* (2006) reported that yellow polythene mulch increased the yield and yield contributing characters of tomato by suppressing the TPVV infection and these results were in line with Islam and Faruq (2009) which showed that application of polythene sheet as mulch increased the grain yield of mungbean by 11.73 %. Li *et al.* (2007) showed that plastic film mulch improves the yield in rice and application of polythene sheet as mulch increases the grain yield of rice as the density of weed has been diminished due to increase in temperature of soil (Mohtisham *et al.*, 2013). In sunflower grain yield (24 and 19 t/ha) has been recorded when plastic sheet mulch was applied (Agele *et al.*, 2010).

Conclusion

The results of this study indicated that Johnson grass competition had a strong impact on the growth and yield of mungbean and sorghum. The results showed that application of treatments PSM have suppressed the growth of Johnson grass and increased the yield of mungbean and sorghum. The weed density of Johnson grass in CS₁ was 7 plants/ m² and in CS₂ it was 12.60 plants/m² with the amendment of PSM. It was recorded maximum in non-treated control treatment in all the cropping systems. On the other hand, the average yield for mungbean for treatment PSM was 1610.17 kg/ha and sorghum yield was 4061 kg/ha during both study years. The plant density of Johnson grass in CS₁ + PSM was 69.10 % was lower than control while it was 58.25 % lower than control in CS₂ + PSM. The yield of mungbean was 49.67 % greater than control while sorghum yield has increased yield of 7.10 %.

Declaration of Interest

There is no conflict of interest among authors.

References

- Abdelrahman, N. A., Abdalla, E. A., Ibrahim, E. A., & El Naim, A. M. (2016). The effect of plastic mulch on growth and yield of rain-fed cowpea and watermelon in North Kordofan State of Sudan. *World*, 4(5), 139-142.
- Agele, S. O., Olaore, J. B., & Akinbode, F. A. (2010). Effect of some mulch materials on soil physical properties, growth and yield of sunflower (*Helianthus Annuus*, L). *Advances in Environmental Biology*, 368-376.

- Ahmad S, Raza M. A. S, Saleem M. F, Zahra S. S, Khan I. H, Ali M, Shahid A. M, Iqbal R, & Zaheer M. S (2015) Mulching strategies for weeds control and water conservation in cotton., *J. Agric. Biol. Sci* 8:299–306
- Ahmad S, Raza M. A. S, Saleem M. F, Zaheer M. S, Iqbal R, Haider I, Aslam M. U, Ali M, & Khan I. H (2020) Significance of partial root zone drying and mulches for water saving and weed suppression in wheat. *J. Anim. Plant. Sci.* 30:154–162
- Asif, M., Nadeem, M. A., Aziz, A., Safdar, M. E., Adnan, M., Ali, A., & Abbas, B. (2020). Mulching improves weeds management, soil carbon and productivity of spring planted maize (*Zeamays L.*). *International Journal of Botany Studies*, 5(2), 57-61.
- Awal, M. A., Dhar, P. C. & Sultan, M. S. (2016). Effect of mulching on micro climatic manipulation, weed suppression, and growth and yield of pea (*Pisum sativum L.*). *Journal of Agriculture and Ecology Research International*, 1-12.
- Azadbakht, A., Alebrahim, M. T. & Ghavidel, A. (2017). The effect of chemical and non-chemical control methods on weeds in potato (*Solanum tuberosum L.*) cultivation in Ardabil province, Iran. *Applied Ecology and Environmental Research*, 15(4), 1359-1372.
- Azad, B, Hassandokht, M. R. & Parvizi K. (2015): Effect of mulch on some characteristics of potato in Asadabad, Hamedan, Iran. *International Journal of Agronomy and Agricultural Research* 6(3): 139-147
- Binimelis R, Pengue W, & Monterroso I (2009) “Transgenic treadmill”: responses to the emergence and spread of glyphosate-resistant Johnsongrass in Argentina. *Geoforum* 40:623–633
- Blanco-Canqui, H.; & Lal, R. Cropping Systems. In Principles of Soil Conservation and Management; Springer: Dordrecht, The Netherlands, 2010; pp. 165–193.
- Burnside, O. C., & Wicks, G. A. (1965). Effect of Herbicides and Cultivation Treatments on Yield Components of Dryland Sorghum in Nebraska 1. *Agronomy Journal*, 57(1), 21-24
- Chirita R, Grozea I, Sarpe N, Lauer K. F (2007) Control of *Sorghum halepense* (L.) species in western part of Romania. *Commun Agric Appl Biol Sci*, 73:959–964
- Czarnota, M. A., Paul, R. N., Weston, L. A., & Duke, S. O. (2003). Anatomy of sorgoleone-secreting root hairs of Sorghum species. *International Journal of Plant Sciences*, 164(6), 861-866.
- Dalley, C. D., & Richard, E. P. (2008). Control of rhizome Johnson grass (*Sorghum halepense*) in sugarcane with trifloxysulfuron and asulam. *Weed Tech*, 22(3), 397-401.
- David, C. B and J. M. Chandler. 1987. Effect of herbicide and weed height on Johnsongrass (*Sorghum halepense L.*) and cotton (*Gossypium hirsutum*) yield. *Weed Tech.* (1): 207-211.
- El-Samnoudi, I. M., Ibrahim, A. E. A. M., Abd El Tawwab, A. R., & Abd El-Mageed, T. A. (2019). Combined effect of poultry manure and soil mulching on soil properties,

- physiological responses, yields and water-use efficiencies of sorghum plants under water stress. *Communications in Soil Science and Plant Analysis*, 50(20), 2626-2639
- Farooq, M., Khan, I., Nawaz, A., Cheema, M. A., & Siddique, K. H. (2020). Using sorghum to suppress weeds in autumn planted maize. *Crop protection*, 133, 105162.
- Heap I (2014) Herbicide resistant weeds. In: Pimentel D, Peshin R (eds) Integrated pest management. *Springer*, Dordrecht, pp 281–301
- Henderson L (2001) Alien weeds and invasive plants. Plant Protection Research Institute Handbook No. 12. Paarl Printers, Cape Town, South Africa
- Hoffman M. L & Buhler D. D. (2002) Utilizing sorghum as a functional model of crop–weed competition. I. Establishing a competitive hierarchy. *Weed Sci* 50:466–472
- Holm L. G, Plucknett D. L, Pancho J. V & Herberger J. P (1997) Sorghum halepense L. Pers. In: Holm (ed) The world’s worst weeds, distribution and biology. The University Press of Hawaii, Honolulu, pp 54–61
- Huang H, Ling T, Wei S, & Zhang C (2015) A new 4-oxazolidinone from *Sorghum halepense* (L.) Pers. *RecNat Prod* 9:247–250
- Islam, M. T., & Faruq, A. N. (2009). Effect of some cultural practices for the management of mungbean yellow mosaic disease. *Journal of Sher-e-Bangla Agricultural University*, 3(1).
- Kader M. A, Singha A, Begum M. A, Jewel A, Khan F. H & Khan N. I (2019). Mulching as water-saving technique in dry land agriculture. *Bulletin of the National Research Centre* 43:1–6
- Karim, Z., Akanda, A.M., Hossain, M.S., Islam, M. M. & Rahman, M.M.E. (2006). Effect of polythene mulch on the management of Tomato purple vein virus (TPVV) in the field. *Bangladesh Journal of Plant Pathology*. 22(1&2):79-83.
- Khairnar, C. B., Goud, V. V., & Sethi, H. N. (2014). Pre-and post-emergence herbicides for weed management in mungbean.
- Kumar, V., Tripathi, S. S., Sachan, H. K., & Singh, V. P. (2008). Effect of weed management techniques on weed dynamics and green fodder yield of sorghum. *Indian Journal of Weed Science*, 40(1&2), 72-74.
- Lamont W. J. (2005) Plastics: modifying the microclimate for the production of vegetable crops. *Hortic. Technol.* 15:477–481
- Lightfoot D. R. (1994) Morphology and ecology of lithic-mulch agriculture. *Geo. Rev.* 25:172–185
- Mihovsky T & Pachev I (2012) Reduced tillage practices. *Banat’s J Biotech* 3:49–58

- Mistaskas, M. B., C. E. Tsoilis., E. G. Ilis, & D. A. Christos. 2003, Interference between corn and Johnsongrass (*Sorghum halepense* L.) from seed or rhizomes. *Weed Sci.* (51): 540-545
- Mohtisham A, Ahmad R, Ahmad Z, & Aslam M. R. (2013) Effect of different mulches techniques on weed infestation in aerobic rice (*Oryza sativa* L.). *Ame-Eur. J. Agric. Environ. Sci.* 13: 153-157
- Naeem, M., Farooq, M., Farooq, S., Ul-Allah, S., Alfarraj, S., & Hussain, M. (2021). The impact of different crop sequences on weed infestation and productivity of barley (*Hordeum vulgare* L.) under different tillage systems. *Crop Protection*, 149, 105759.
- Nalayini P. (2007) Poly-mulching a case study to increase cotton productivity. Senior scientist, Central Institute for Cotton Research, Regional Station, Coimbatore
- Ngouajio M. & McGiffen M.E. (2004) Sustainable vegetable production: effects of cropping systems on weed and insect population dynamics. *Acta. Hort.* 638:77–83
- Novak R., Dancza I., Szentey L. & Karaman J. (2009) Arable weeds of Hungary. The 5th National Weed Survey (2007–2008). Ministry of Agriculture and Rural Development, Budapest, Hungary
- Rathore A. L, Pal A. R. & Sahu K. K. (1998). Tillage and mulching effects on water use, root growth, and yield of rain-fed mustard and chickpea grown after lowland rice. *J. Sci. Food. Agric.* 78:149–161
- Shahzad, M., Jabran, K., Hussain, M., Raza, M. A. S., Wijaya, L., El-Sheikh, M. A., & Alyemeni, M. N. (2021). The impact of different weed management strategies on weed flora of wheat-based cropping systems. *Plos One*, 16(2), e0247137.
- Singh, G., Aggarwal, N., & Ram, H. (2014). Efficacy of postemergence herbicide imazethapyr for weed management in different mungbean (*Vigna radiata*) cultivars. *Indian Journal of Agricultural Sciences*, 84(4), 540-543.
- Uludag A., Gozcu D., Rusen M., Guvercin R.S. & Demir A. (2007). The effect of Johnsongrass densities (*Sorghum halepense* L. Pers.) on cotton yield. *Pak J Biol. Sci.* 10:523–525
- Unger, P. W., & Jones, O. R. (1981). Effect of soil water content and a growing season straw mulch on grain sorghum. *Soil Science Society of America Journal*, 45(1), 129-134
- Valverde B. E., & Gressel J., (2006) Dealing with the evolution and spread of *Sorghum halepense* glyphosate resistance in Argentina. <http://www.sinavimo.gov.ar/files/senasareport2006.pdf>
- Vila-Aiub M. M., Gundel P. E., Yu. Q.&Powles S. B. (2013). Glyphosate resistance in *Sorghum halepense* and *Lolium rigidum* is reduced at suboptimal growing temperatures. *Pest Manag. Sci.*69:228–232

- Xu, G. W., Zhang, Z. C., Zhang, J. H., & Yang, J. C. (2007). Much improved water use efficiency of rice under non- flooded mulching cultivation. *Journal of Integrative Plant Biology*, 49(10), 1527-1534.
- Yaqub, F., & Shahzad, S. (2009). Effect of solar heating by polyethylene mulching on sclerotial viability and pathogenicity of *Sclerotium rolfsii* on mungbean and sunflower. *Pak. J. Bot*, 41(6), 3199-3205.