MATHEMATICAL MODELING OF BIRD HARVESTING IN INTENSIVE POULTRY SYSTEM

ABSTRACT:
This work, presents a formulation of mathematical model of bird harvesting in an intensive poultry system, under the assumption that under a favourable environmental atmosphere and good management system, the birds have logistic growth. The model is analysed using methods from dynamical system theory and theory of calculus. It was established that the system has two steady state, the two equilibrium state are both locally asymptotically stable. The first one is stable if there is a bound on the harvest rate of the birds, which is proportional to the growth rate of the birds. The second equilibrium state is locally asymptotically stable (LAS) if $K < \frac{r(C + \gamma)}{\rho}$ that is if the carrying capacity is less than the ratio of the sum of $Per\ unit\ cost\ of\ birds$ and $Per\ unit\ tax\ on\ the\ bird$ to that of $Per\ unit\ price\ of\ the\ birds$. Further analysis indicates that the limiting population of bird, that is the maximum population of birds that the available resources in the system can sustain and also ensures harvesting profitability is given as $\frac{K\left(r-\frac{K\rho\,(r-\mu)-r(C+\gamma)}{K\rho}-\mu\right)}{r}$

This occurs at the point

$$(\bar{t},\bar{B}) = \left(\left(\frac{1}{\left(r-\beta\left(\frac{K\rho\,(r-\mu)-r(C+\gamma)}{K\rho}\right)-\mu\right)}\mathrm{Ln}\left(\frac{\frac{K\left(r-\frac{K\rho\,(r-\mu)-r(C+\gamma)}{K\rho}-\mu\right)}{r}-B_0}{B_0}\right),\left(\frac{K\left(r-\frac{K\rho\,(r-\mu)-r(C+\gamma)}{K\rho}-\mu\right)}{2r}\right)\right)\right)$$

Keywords: Profitability, Harvesting, Stability, Poultry, Birds

1Introduction

Poultry refers to birds that are kept or raised, either for private or commercial purposes and generally includes birds such as Chicken, Turkey, Duck, Goose, Quail, Pheasant, Pigeon, Guinea Fowl, Pea fowl, Ostrich, e.t.c [1]

Poultry farming is one of the smallest livestock investment that an individual or household can easily undertake. Income generation is the primary goal of this poultry keeping. Eggs can provide a regular income albeit small, while the sales of birds provide a more flexible source of income as required. [2]

A study of income generation in poultry farming in East Kakimantan, Indonesea showed that family poultry accounted for about 53 percent of total income and was used for food, school fees and unexpected expenses such as medicine [3]. Poultry are kept under a wide range of conditions, which can be classified into one of four broad production system such as free-range extensive system, Backyard extensive system, Semi-intensive system and Intensive system. The intensive systems are used by medium or large-scale commercial enterprises and also at the household level. In this system, birds are fully confined either in houses or cages.

[4] presented a mathematical modelling of the mortality rate of broilers chicken using a standard S-I-R model. Analysis of the modelshowed that poultry industry is one of the most successful agricultural venture in any nation's economy. They established that for successful rearing of broiler chicken, good welfare conditions must be maintained to avoid extreme case of the effective mortality rate being large. Also good management programmes must be developed in broiler production to combat early mortality. It has been established by many researchers [10, 11 12,13,19]that effective control ,reduction and prevention of infection in poultry can enhance better production and profitability in poultry.

UNDER PEER REVIEW

[7] stated that harvesting models based on ordinary differential equations are commonly used in the fishery industry and wildlife management to model the evolution of a population depleted by harvest mortality. He presented a project consisting of a series of scenarios based on fishery harvesting models to teach the application of theoretical concepts learned in a differential equations course to scenarios encountered in real fisheries. In this projects a thorough understanding of simplifying assumptions inherent in various models was required, as well as a qualitative analysis of phase portraits, bifurcations, and stability of steady states. Parameters were estimated and equations were solved both analytically and numerically. Students learn to respond to a professional request from a fishery in the form of a scientific report, which requires organizing and communicating assumptions, models, solution methods, results, and a final recommendation with clarity and professionalism.

[5], presented a mathematical model of fish harvesting in a common access fishery. The model was formulated and analysed using methods from dynamical system theory. The analysis of the model showed that the system has two steady states. The first one is unstable while the second one is globally asymptotically stable if the carrying capacity of the water has a lower bound.

[9], developed mathematical model of bacteria-nutrient harvesting in a cultured environment. The model which assumes that the rate of harvesting of these bacteria is constant results in a system of first order differential equations. Analyzing the model, it was discovered that the product of the maximum nutrient uptake per cell and the number of cells produced per unit of nutrient uptake is constant ($\boldsymbol{VY}= \boldsymbol{ln2}+ \boldsymbol{h}$). It was also assumed that the rate of harvesting of these bacteria varies and a corresponding model was developed. Analyzing this model using methods from dynamical systems theory, it was seen that the system has two steady states. The first steady state is unstable while the second is globally asymptotically stable if the carrying capacity of the environment has a lower bound, which is a ratio of the harvesting coefficient of the bacteria, cost per unit effort per unit price of the bacteria.

In this paper, we present a mathematical model of bird harvesting in an intensive poultry system. We assume that under a favourable environmental atmosphere and good management system the birds have logistic growth. The model is formulated and analysed using methods from dynamical system theory and theory of calculus.

2 FORMULATION OF THE MATHEMATICAL MODEL

2.1 ASSUMPTIONS

1. In the absence of epidemic outbreak, the population of birds has a logistic growth.

2. The population of birds is reduced by the number of birds harvested at time t provided the net income is positive

3. The per carpita rate of birds reproduction decreases with increase in the population of birds.

4. in the presence of favourable environmental conditions and good management system, death due to natural occurrences does not cause any damage.

5, There is no migration to or out of the poultry except for harvesting purpose.

6. Work done increases with positive revenue and decrease otherwise.

2.2 Symbols/Parameters

$B(t) \ = Population\ of\ birds\ at\ time\ t > 0$

$W(t) \ = Workdone\ in\ the\ poultry\ \ at\ time\ t > 0$

UNDER PEER REVIEW

$r \ = Rate\ of\ reproduction$

$K \ = Carrying\ capacity\ of\ the\ poultry\ house$

$\beta \ = harvest\ rate\ of\ birds\ which\ depends\ on\ the\ revenue\ and\ maturity$

$\mu \ = Na\ tural\ death\ rate\ of\ birds$

$\rho = Per\ unit\ price\ of\ the\ birds$

$\mathrm{C} = \ Per\ unit\ cost\ of\ birds$

$\gamma \ = Per\ unit\ tax\ on\ the\ birds$

2.3 The Model Equation

$$\frac{dB}{dt} = rB\left(1-\frac{B}{K}\right) - \beta WB - \mu B \ldots\ldots\ldots\ldots\ldots\ldots..(2.1)$$

$$\frac{dW}{dt} = \alpha W(\rho B - c - \gamma) \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots..(2.2)$$

Where $R = W(\rho B - c - \gamma) \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots..(2.3)$

3 Analysis of the Model

Using methods from [14, 15,16,20,21,22] the system (2.1) – (2.3) is analysed as follows:

$$\frac{dB}{dt} = rB\left(1-\frac{B}{K}\right) - \beta WB - \mu B \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots (3.1)$$

$$\frac{dW}{dt} = \alpha W(\rho B - \mathrm{C} - \gamma) \quad \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots.(3.2)$$

At equilibrium

$$rB\left(1-\frac{B}{K}\right) - \beta WB - \mu B = 0 \quad \ldots\ldots\ldots\ldots..(3.3)$$

$$\alpha W\rho B - \mathrm{C}\alpha W - \gamma\alpha W = 0 \quad \ldots\ldots\ldots\ldots\ldots.(3.4)$$

From $(5.2) B^0 = \frac{\mathrm{C} + \gamma}{\rho}$

From $(5.1) W^0 = \frac{K\rho(r-\mu) - r(\mathrm{C}+\gamma)}{K\rho\beta}$

Using methods from [6.17,18] , Linearizing, we obtain the Jacobian matrix thus:

$$J = \begin{pmatrix} r\left(1-\frac{B^0}{K}\right) - \frac{B^0}{K} - \beta W^0 - \mu & -\beta B^0 \\ \alpha\rho W^0 & \alpha\rho B^0 - \mathrm{C}\alpha - \gamma\alpha \end{pmatrix}$$

3.1 FIRST EQUILIBRIUM STATE

At $(B^0, W^0) = (0 \ , \ 0)$

Then

UNDER PEER REVIEW

$$J_0 = \begin{pmatrix} r-\mu & 0 \\ 0 & -C\alpha-\gamma\alpha \end{pmatrix}$$

$$|J_0 - \lambda I| = \begin{vmatrix} r-\mu-\lambda & 0 \\ 0 & -\alpha(C+\gamma)-\lambda \end{vmatrix}$$

This implies that, $\lambda_1 = r-\mu$, $\lambda_2 = -\alpha(C+\gamma)$

The system is stable at $(B^0, W^0) = (0, 0)$ if $r-\mu < 0$, $-\alpha(C+\gamma) < 0$. This implies $C+\gamma > 0$. That is, there must be a bound on the harvest rate of the the birds which is proportional to the growth rate of the birds.

3.2SECOND EQUILIBRIUM STATE

$$(B^0, W^0) = \left(\frac{C+\gamma}{\rho}, \frac{K\rho(r-\mu)-r(C+\gamma)}{K\rho\beta}\right)$$

$$J = \begin{pmatrix} r\left(1-\frac{B^0}{K}\right)-\frac{B^0}{K}-\beta W^0-\mu & -\beta B^0 \\ \alpha\rho W^0 & \alpha\rho B^0 - C\alpha-\gamma\alpha \end{pmatrix}$$

Substituting the endemic equilibrium point into the Jacobian, we obtain the Jacobian for the endemic equilibrium point.

$$J_E = \begin{pmatrix} \frac{-(C+\gamma)}{\rho K} & \frac{-\beta(C+\gamma)}{\rho} \\ \frac{\alpha\rho K-\alpha r(C+\gamma)}{K\beta} & 0 \end{pmatrix}$$

$$|J_E - \lambda I| = \begin{vmatrix} \frac{-(C+\gamma)}{\rho K}-\lambda & \frac{-\beta(C+\gamma)}{\rho} \\ \frac{\alpha\rho K-\alpha r(C+\gamma)}{K\beta} & 0-\lambda \end{vmatrix}$$

This implies that, $\rho K\lambda^2 + (C+r)\lambda + \delta = 0, where\ \delta = (C+r)[\alpha\rho K - \alpha r(C+\gamma)$

Hence $\lambda_1 = \frac{-(C+\gamma)}{2\rho K} + \frac{\sqrt{(C+\gamma)^2-4K\rho\delta}}{2K\rho}$, $\lambda_2 = \frac{-(C+\gamma)}{2\rho K} - \frac{\sqrt{(C+\gamma)^2-4K\rho\delta}}{2K\rho}$

For real λ, then $(C+\gamma)^2 - 4K\rho\delta \geq 0$

This implies that $(C+\gamma)^2 \geq 4K\rho\delta \Rightarrow C+\gamma) \geq \sqrt{4K\rho\delta}$

$$C+\gamma) \geq \sqrt{4K\rho(C+r)[\alpha\rho K - \alpha r(C+\gamma)}$$

$$\Rightarrow K \geq \frac{1}{\alpha\rho}\alpha r(C+\gamma)$$

The steady state is stable only if $\lambda_1 < 0\ and\ \lambda_2 < 0$ This is possible only if

$$\frac{-(C+\gamma)}{2\rho K} \pm \frac{\sqrt{(C+\gamma)^2-4K\rho\delta}}{2K\rho} < 0$$

$$\Rightarrow \frac{-(C+\gamma)}{2\rho K} < \pm\frac{\sqrt{(C+\gamma)^2-4K\rho\delta}}{2K\rho}$$

$$\Rightarrow -(C+\gamma) < \pm\sqrt{(C+\gamma)^2-4K\rho\delta}$$

UNDER PEER REVIEW

Squaring both sides, we have

$$(C + \gamma)^2 = (C + \gamma)^2 - 4K\rho\delta$$

$$(C + \gamma)^2 = (C + \gamma)^2 - 4K\rho((C + r)[\alpha\rho K - \alpha r(C + \gamma))$$

$$\Rightarrow 4K\rho((C + r)[\alpha\rho K - \alpha r(C + \gamma)) < 0$$

$$\Rightarrow \alpha\rho K - \alpha r(C + \gamma)$$

$$\Rightarrow \alpha\rho K < \alpha r(C + \gamma)$$

$$\Rightarrow K < \frac{r(C + \gamma)}{\rho}$$

Hence, the endemic steady state $(B^0, W^0) = \left(\frac{C + \gamma}{\rho}, \frac{K\rho(r - \mu) - r(C + \gamma)}{K\rho\beta}\right)$

Is locally asymptotically stable (LAS) if $K < \frac{r(C + \gamma)}{\rho}$ that is if the carrying capacity is less than the ratio of the sum of $Per\ unit\ cost\ of\ birds$ and $Per\ unit\ tax\ on\ the\ bird$ to that of $Per\ unit\ price\ of\ the\ birds$

3.3 LIMITING CAPACITY AND POINT OF LIMITING CAPACITY OF THE SYSTEM

Assuming the initial population of the bird is given as $B(0) = B_0$. $W = \frac{K\rho(r - \mu) - r(C+\gamma)}{K\rho\beta}$, $Workdone\ in\ the\ poultry$. From the theory of initial value problem of a differential equation and the methods in [8,9,10, 18,23],we obtain the population of the bird at time t to be

$$B(t) = \frac{\frac{K\left(r - \frac{K\rho(r - \mu) - r(C+\gamma)}{K\rho} - \mu\right)}{r}}{1 + \left(\frac{K\left(r - \frac{K\rho(r - \mu) - r(C+\gamma)}{K\rho} - \mu\right)}{r} B_0 - 1\right) e^{-\left(r - \beta\left(\frac{K\rho(r - \mu) - r(C+\gamma)}{K\rho}\right) - \mu\right)t}}$$

By taking the limit of the bird's population as t approaches infinity, the limiting population of bird, that is the maximum population of birds that the available resources in the system can sustain and also ensures harvesting profitability is given as $L = \frac{K\left(r - \frac{K\rho(r - \mu) - r(C+\gamma)}{K\rho} - \mu\right)}{r}$

This occurs at the point

$$(\bar{t}, \bar{B}) = \left(\left(\frac{1}{\left(r - \beta\left(\frac{K\rho(r - \mu) - r(C+\gamma)}{K\rho}\right) - \mu\right)} \mathrm{Ln}\left(\frac{\frac{K\left(r - \frac{K\rho(r - \mu) - r(C+\gamma)}{K\rho} - \mu\right)}{r} - B_0}{B_0}\right), \left(\frac{K\left(r - \frac{K\rho(r - \mu) - r(C+\gamma)}{K\rho} - \mu\right)}{2r}\right)\right)\right)$$

4CONCLUSION

The work, presents a mathematical model of bird harvesting in an intensive poultry system, under the following assumptions:In the absence of epidemic outbreak, the population of birds has a logistic growth. The population of birds is reduced by the number of birds harvested at time t provided the net income is positive. The per carpita rate of bird's reproduction decreases with increase in the population of birds. In the presence of favourable environmental conditions

UNDER PEER REVIEW

and good management system, death due to natural occurrences does not cause any damage.There is no migration to or out of the poultry except for harvesting purpose. Work done increases with positive revenue and decrease otherwise. The model is analysed using methods from dynamical system theory and theory of calculus. It was established that the system has two steady state, the two equilibrium state are both locally asymptotically stable. The first one is stable if there is a bound on the harvest rate of the birds, which is proportional to the growth rate of the birds. The second equilibrium state is locally asymptotically stable (LAS) if $K < \frac{r(C + \gamma)}{\rho}$ that is if the carrying capacity is less than the ratio of the sum of $Per\ unit\ cost\ of\ birds$ and $Per\ unit\ tax\ on\ the\ bird$ to that of $Per\ unit\ price\ of\ the\ birds$. Further analysis indicates that the limiting population of bird, that is the maximum population of birds that the available resources in the system can sustain and also ensures harvesting profitability is given as $L = \frac{K\left(r - \frac{K\rho(r-\mu) - r(C+\gamma)}{K\rho} - \mu\right)}{r}$

This occurs at the point $(\bar{t}, \bar{B}) =$

$$\left(\left(\frac{1}{\left(r - \beta\left(\frac{K\rho\,(r-\mu) - r(C+\gamma)}{K\rho}\right) - \mu\right)} \mathrm{Ln}\left(\frac{\frac{K\left(r - \frac{K\rho\,(r-\mu) - r(C+\gamma)}{K\rho} - \mu\right)}{r} - B_0}{B_0}\right), \left(\frac{K\left(r - \frac{K\rho\,(r-\mu) - r(C+\gamma)}{K\rho} - \mu\right)}{2r}\right)\right)\right)$$

UNDER PEER REVIEW

References

1. G. D Butcher et al, Dairy and Poultry Science, Faculty of Veterinary Medicine, Florida University.
2. https://en.wikipedia.org/wiki/Poultry#:~:text=%22Poultry%22%20can%20be%20defined%20as,and%20squabs%20(young%20pigeons).
3. Singh M, Mollier RT, Paton RN, Pongener N, Yadav R, Singh V, Katiyar R, Kumar R, Sonia C, Bhatt M, Babu S, Rajkhowa DJ and Mishra VK (2022) Backyard poultry farming with improved germplasm: Sustainable food production and nutritional security in fragile ecosystem. *Front. Sustain. Food Syst.* 6:962268. doi: 10.3389/fsufs.2022.962268
4. Inyama S.C(2008) Mathematical model of fish harvesting in a common access fishery, journal of mathematical sciences Vol19, N0 1 pages 43-47
5. Joshua E. E et al (2008) Mathematical modelling of mortality rate in broiler chickens, journal of Nigerian Association of mathematical physics vol. 12 page 437 -442
6. KinfeHailemariamHntsa, ZebebeTekaMengesha (2016) 'Mathematical modeling of Fish Resources Harvesting with Predator at Maximum Sustainable Yield, Journal of Mathematical Theory and Modeling ISSN 2224-5804 (Paper) ISSN 2225-0522 (Online) Vol.6, No.6, 2016
7. Danilo R. Diedrichs(2019)Using Harvesting Models to Teach Modeling with Differential Equations, PRIMUS Journal **Volume 29, 2019 - Issue 7: Modeling Approach to Teaching Differential Equations(Part 2) https://doi.org/10.1080/10511970.2018.1484397**
8. IA Agwu, SC Inyama, RA Umana, A Omame, N Ukanwoke, A Ofomata, HI Mbachu, ES Udofia, JI Uwakwe (2018) Determining the impact of variation of Harvesting Effort on the Qualitative Behaviour of a Coexistence Steady State Solution and its Stability in Prey-Predator Fishery Model, Academic Journal of Applied Mathematical Sciences Vol. 4 Issue 10 Pages 119-128, 2018

9. Udofia Ekere Sunday , Amos AmosIdungafa (2018) Mathematical Model of Bacteria-Nutrient Harvesting in A Cultured Environment, Journal of the Nigerian Association of Mathematical Physics Volume 46 (May, 2018 Issue), pp115 –118, 2018 © J. of NAMP
10. Udofia, Ekere Sunday, Sampson, Marshal Imeh (2014) Mathematical Model for the Epidemiology of Fowl Pox Infection Transmission That Incorporates Discrete Delay, IOSR Journal of Mathematics (IOSR-JM) e-ISSN: 2278-5728, p-ISSN: 2319-765X. Volume 10, Issue 4 Ver. V (July - Aug. 2014), PP 08-16 www.iosrjournals.org
11. Udofia Ekere Sunday and Inyama Simeon Chioma (2011) Mathematical Model of the Impact of Vaccination on the Transmission Dynamics of Fowl Pox in Poultry, Journal of Modern Mathematics and Statistics 5 (5-6): 102-105, 2012 ISSN 1994-5388 © 2011 Medwell Journals
12. Udofia Ekere Sunda and Inyama Simeon Chioma (2011) MathematicalModeling of the Transmission Dynamics of Fowl Pox in Poultry, Journal of Modern Mathematics and Statistics 5 (5-6): 106-111, 2012 ISSN 1994-5388 © 2011 Medwell Journals, 2011
13. UDOFIA EKERE SUNDAY, UDOH KENNETH JUMBO AND INYAMA SIMEON CHIOMA (2016) THE IMPACT OF MEDIA COVERAGE ON THE TRANSMISSION DYNAMICS OF FOWL POX IN POULTRY, International J. of Math. Sci. &Engg. Appls. (IJMSEA) ISSN 0973-9424, Vol. 10 No. I (April, 2016), pp. 103-114
14. Udofia, Ekere S, and Etukudo Idorenyin A (2019) Optimal Allocation of Biscuit Ingredient in the Production Process - An Invariant Property Based Algorithm Approach, Mathematical Theory and Modeling www.iiste.org ISSN 2224-5804 (Paper) ISSN 2225-0522 (Online) DOI: 10.7176/MTM Vol.9, No.4, 2019
15. ONUOHA JOY LJEOMA, INYAMA SIMEON CHIOMA , UDOFIA EKERE SUNDAY AND OMAME ANDREW (2015) Mathematical Model of the Transmission Dynamics of Swine Flu with the Vaccination of Non Newborns, International J. of Math. Sci. &Engg. Appls. (IJMSEA) ISSN 0973-9424, Vol. 9 No. I (March, 2015), pp. 1-17 www.ascent-journals.com
16. Ekere S. Udofia, (2023), Mathematical Model of Male Circumcision in HIV/AIDS Preventions, International Journal of Innovative Science and Research Technology, Volume 8, Issue 8, August – 2023 ISSN No: -2456-2165 www.ijisrt.com
17. O. Diekmann and .J. A. P. Heesterbeek, Mathematical epidemiology of infectiousdiseases: Model building, analysis and interpretation, Journal of the RoyalSociety Interface 7 (2000), no. 47, 873–885.
18. P. Driessche and Z. Shua, Global stability of infectious disease models using lyapunov functions, SIAM J. APPL.MATH 73 (2013), no. 4, 1513–1532.
19. Udofia Ekere Sunday and Inyama Simeon Chioma (2012), Application of Optimal Control to the Epidemiology of Fowl Pox Transmission Dynamics in Poultry, Journal of Mathematics and Statistics 8 (2): 248-252
20. **Udofia, EkereSunday, Sampson, Marshal Imeh** (2014) ,Mathematical Model of the Effect of Complacency in HIV/AIDS Preventions, IOSR Journal of Mathematics (IOSR-JM) e-ISSN: 2278-5728, p-ISSN: 2319-765X. Volume 10, Issue 6 Ver. V (Nov - Dec. 2014), PP 30-33 www.iosrjournals.org
21. ONUOHA JOY LJEOMA, INYAMA SIMEON CHIOMA AND UDOFIA SUNDAY EKERE(2014), Mathematical Model of the Transmission Dynamics of Swine Flu with the Vaccination of Newborns, International J. of Math. Sci. &Engg. Appls. (IJMSEA) ISSN 0973-9424, Vol. 8 No. V (September, 2014), pp. 217-229
22. Udofia, Ekere Sunday and Inyama, Simeon Chioma Mathematical model of Structural Strategy (Delayed First Intercourse) in HIVAIDS Prevention, Journal of the Nigerian Association of Mathematical Physics Volume 24 (July, 2013) pp 257-260 © J. of NAMP
23. ***Udofia, Ekere Sunday and Umana, Reuben Andrew (2014),* Weak Solutions of Boundary Value Problems *Journal of the Nigerian Association of Mathematical Physics Volume28, No. 1,* (*November, 2014*), pp21 – 30© *J. of NAMP***

UNDER PEER REVIEW