

# Effect Of Preservative Methods on Organoleptic Characteristics of Broiler Chicken NeckMeat

## Abstract

*The effect of preservative methods on organoleptic characteristics of broiler chicken neck is investigated, forty pieces of broiler neck meat was purchased from ipata frozen meat shop, ten (10) pieces each of the meat part was fried and roasted and evaluation was carried out almost immediately after which ten (10) each was again fried and roasted and keep at room temperature for two (2) weeks and at the end of the two weeks, sensory evaluation was again carried out, after each taste the questionnaire was filled, the nine point edonic scale was used as questionnaire, the panelist were ten (10) each for both boys and girls with their ages ranging from 18-22 years old, the data collected was then analyzed using Randomized combined block design (RCBD) statistic tool and at the end the panelist preferred the roasted meat ( $P < 0.05$ ), this may be as a result of the heat that actually killed the microorganisms that can cause spoilage while the oil that was used in frying may be the cause of spoilage (rancidity) in fried meat*

**KEYWORDS:** Organoleptics characteristics, frozen, preservatives, edonic scale, Statistic tool.

## INTRODUCTION

The continuous rise in human population all over the world with annual average of 7 billion has brought an increase in the demand for animal protein. Estimated current level of consumption of protein in the world is 7g per caput per day (Lawal et al., 2013).

If we must continue to produce food at our present rate, then there will be food scarcity in Nigeria and in deed all over the world, there is that need to increase the rate of food production to match the continuous increase in food production to avoid food scarcity and have food security (Lawal et al., 2013).

In Nigeria, the average daily consumption of 54g protein with 6.5g from animal source falls below the recommended daily protein of 86g and 8.4g of protein from animal source (Abu, et al., 2008)

To alleviate this situation, it has been realized that broiler production is the fastest and easiest route (Lara, 2008; Demirbas, 2009; Nworgu et al., 2000) since they are prolific, possess a high feed conversion ratio and are accepted by all, irrespective of religion.

Continuous production will at a point lead to excess and there will that need for storage or Preservation to support food security, various method of food preservation are available for commercial scale but those practiced by back yard farmers are to be considered in this study to avoid wastage from spoilage caused by microorganisms and this includes frying and roasting.

## 1.2 Justification

- i-Continuous production of broiler birds lead to more of the meat
- ii-Easier way of preserving meat is the cooking methods, refrigerating might be expensive for most Nigerians

### **1.3 Objectives of the study**

- i-Purchase the needed birds part for experimental reason
- ii-Ten(10) broiler chicken neck each were then fried and roasted and store for 2 weeks
- iii- Freshly prepared broiler neck by roasting and frying were evaluated for organoleptic characteristics and this were compare to the taste given after two (2) weeks
- iv- Invite panelist for evaluation

### **Materials And Methods**

Site of the experiments: The experiment was carried out in the Agricultural garden of Kwara State Polytechnic, Ilorin,

Meat sample: The fifty (50) pieces of sizeable broiler chicken neck was purchased from cold room and thirty (30) were use for evaluation immediately, ten (10) each was boiled, fried and roasted while the remaining twenty (20) was used by frying ten (10) and then roast then and keep for evaluation after two (2weeks)

Panelist: Twenty panelist was invited with ten (10) males and ten (10) females with their ages ranging from 18-23years old. They were all briefed on how to fill the questionnaire before embarking on it, and immidiatly after each taste the panelist has to rinse their mouth before tasting another sample,

Statistical analysis: All data collected was then collated and analysed using Randomised complete bloc design

## **RESULTS AND DISCUSSIONS**

### **4.1 Results**

Table 1 below shows the effect of cooking methods on broiler chicken neck, significant difference occurred in the colour of the meat ( $P < 0.05$ ), panelist preferred the roasted meat most and the least is fried one, this is the same result that was obtained with the meat when considering texture, though in this case the boiled meat is accepted least, the flavour, tenderness and juiciness is same with the fried meat being accepted most. . It is good to state here that the boiled meat value is just used as the control.

Table 1: Organoleptic characteristics of Broiler chicken neck cooked with different methods

Cooking methods	Boil	Fry	Roast	SEM
Colour	12.00 <sup>b</sup>	21.36 <sup>c</sup>	20.1 <sup>a</sup>	1.4
Flavour	13.00 <sup>c</sup>	24.03 <sup>a</sup>	21.76 <sup>b</sup>	1.7
Tenderness	21.00 <sup>a</sup>	21.00 <sup>a</sup>	19.00 <sup>b</sup>	1.4
Juiciness	23.00 <sup>b</sup>	25.00 <sup>a</sup>	11.90 <sup>c</sup>	6.1
Texture	14.03 <sup>b</sup>	22.00 <sup>a</sup>	21.73 <sup>a</sup>	4.0

a, b, c and d within the same column with different superscripts are significantly different at (P < 0.05).

## 4.2 Discussion

The roasted meat is most preferred meat based on the panelist after preservation for two (2) weeks, this may be as a result to the fact that the roasted meat really kill the micro-organism that can cause damage, the oil used in frying may be a reason for the that instigated the damage in the fry meat that causes the panelist to preferred the roasted meat

## References

- Abu, A.O., Onifade, A. A., Abanikanda, O.T. F. and Obineye, R.I. (2008): Status and promotional Strategies for Rabbit Production in Nigeria. 9<sup>th</sup> World Rabbits Congress Verona, Italy.
- Bowker, B., Hawkins, S and Zhuang, H (2014). Measurement of water-holding capacity in raw and freeze-dried broiler breast meat with visible and near-infrared spectroscopy. *Poultry Science*, 93:1–8.
- Bramblett, V.D, Hostetler, R.L, Vail, G.E and Fraudt, H.N. (1959): Quality of beef as affected by cooking at very low temperature for long periods of time. *Journal of Food science*, 13:707-710
- Bratzler, L.J (1971): Palatability factors affecting meat. The science of meat and products by J.F. price and B.S. Schweicert (2<sup>nd</sup> Edition) W.H. freeman and company 328-363.
- Brittaine, R, Nebambi, O and Lualadio, R (2010) Jatropha: A small- holder bioenergy crop; The potential for pro- poor development. Integrated crop management n<sup>o</sup>8. IFAD/ FAO
- Cassens, R.G and Cooper, C.C (1971); Effect of gender on red and white muscles breast and thigh of turkey meat quality. *Advance in food research* 19:1-74

- Cavitt, L. C., Youm, G. W., Meullenet, J. F., Owens, C. M. and Xiong, R. (2004). Prediction of poultry meat tenderness using razor blade shear, Allo-Kramer shear, and sarcomere length. *Journal of Food Science* 69:SNQ11–SNQ15.
- Chan, J. T. Y., Omana, D. A., and Betti, M.. 2011. Effect of ultimate pH and freezing on the biochemical properties of proteins in turkey breast meat. *Food Chemistry*, 127:109–117.
- Cheng, F. Y., Huang, C. W., Wan, T. C., Liu, Y. T., Lin, L. C. and Lou Chyr, C. Y. (2008). Effects of free-range farming on carcass and meat qualities of black-feathered Taiwan native chicken. *Asian Journal of Animal Science*. 21:1201–1206.
- Cheng C.S (1976) Biochemical studies of post-mortem aging, calcium ion and heating of bovine skeletal muscle. PhD Thesis, Iowa State University, Ames, USA, pp. 250
- Chikpah, S and Demuyakor, B. (2012) : Effect of spontaneous solid state fermentation on proximate and antinutrients composition of *Jatropha curcas* kernel meal obtained from four different Agro climatic area of Ghana. *Int Journal of Science research* Vol1 issue 11
- Chinmoy, B. Ernest, K. Yanful and Maurice, A. (2009) Bergounou biodiesel production from *Jatropha curcas* oil using potassium carbonate as an unsupported catalyst, *International Journal of ChemReaction Eng.*, 7(72), 1-20
- Claus, H. Laccases (2004). Structure, reactions, distribution. *Micron*, 35, 93–96.
- Connor, W. E. (2000). Importance of n-3 fatty acids in health and disease. *The American Journal of Clinical Nutrition*, V. 71(1):171S-175S.
- Cox, A.C (1975); Crab meat in broiler diets. *Poultry Science*, 60; 1567 (Abstract).
- Demirbas, A(2009); Production of biodiesel from algae oil, Part A; Recovery, utilization and Environmental effect, energy sources, 31, No 2, 163-168
- Lawal W.S, Banjoko I.K, Ahmed S.A, Abdulkadri Jimoh, Adebayo Foshade and Zakariyau Mujidat  
2013 Effect of combined method of detoxification of *Jatropha curcas* on organoleptic characteristics of broiler meat . *Journal of Meat Science*, 76(4): 675-681
- Lara, L.J.C, Baião, N.C, Rocha, J.S,R, Lana, A.M.Q, Cançado, S.V, Fontes, D.O, (2008) . Influence of physical form of ration and line on the performance and yield of broiler cuts. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* ;60(4):970-978.
- Nworgu FC, Egbunike GN, Ogundola FI.(2000); Performance and nitrogen utilization of broilers fed full fat extruded soybean meal and full fat soybean. *Tropical Animal Production Investment*. 3:47–54

UNDER PEER REVIEW