

Original Research Article

Puerperal sepsis and its associated factors: Review of cases in a Tertiary Hospital in Jigawa, North-west Nigeria.

ABSTRACT

Background: Puerperal sepsis is an infection of the genital tract occurring anytime between the rupture of membranes and 42nd day postpartum. It is one of the leading causes of maternal mortality accounting for 11% maternal death.

Objective: The aim of the study was to determine the incidence of puerperal sepsis and associated complications. (Factors)

Material and method: It was a retrospective study carried out in the department of Obstetrics and Gynecology of Rasheed Shekoni Federal University Teaching Hospital. Data obtained was analyzed with IBM SPSS version 21.0. Measured variables were expressed in percentage. Test for association was done using chi-square between the dependent and independent variables, setting non-parametric test, setting P-value at <0.05 at 95% level of confidence.

Results: The incidence of puerperal sepsis was of 10.7%. The mean age of the patients was 25.85 ± 7.5. The mean parity was 3.89 ± 4.5. Majority (74.71%) of the women were unbooked and had home delivery (69.3%). Fever (90.7%) was the most frequent presenting complaints. The commonest (75.0%) organism cultured was staphylococcus aureus. Anemia was the commonest (63.6%) complication.

Conclusion: The incidence of Puerperal sepsis in this study is high. Low literacy level, unemployment, un-booked status and home delivery were found to be predisposing factors.

Keywords: Puerperal sepsis, anemia, home delivery, Jigawa.

Comment [mo1]: This section should capture readers attention pls.

Comment [mo2]: Did you measure incidence, if not please remove

Comment [mo3]: Cross-check with the title above please

Comment [mo4]: Which type of retrospective cohort study, pls, indicate

Comment [mo5]: Which variables, categorical or continuous variables, please indicate

Comment [mo6]: Not necessary please

Comment [mo7]: PLEASE REMOVE, No Test of association was conducted. Just stick to the descriptive statistics. Indicate sample size, and sampling technique, means of data collection etc

Comment [mo8]: Retrospective study don't measure incidence unless the type employed is justified. pls check and address it

Comment [mo9]: Please check above

Comment [mo10]: Please where are the supportive findings at the results section

Comment [mo11]: Here please arrange them in ascending order, eg. Anaemia, Home delivery etc

INTRODUCTION

Puerperal sepsis is one of the leading causes of maternal mortality accounting for approximately 11% of maternal death globally.⁽¹⁾ Puerperal sepsis is defined as infection of the genital tract occurring at any time between the onset of rupture of membranes or labour and 42nd day postpartum in which fever and one or more of the following are present: pelvic pain, abnormal vaginal discharge, abnormal smell/foul odour of discharge, delay in the rate of reduction of size of uterus (< 2cm/day during first 8 days).^(2,3) According to international statistical classification of disease and related problems (ICD-10), puerperal sepsis refers to complication related to the puerperium. It includes not only puerperal endometritis but also encompasses other puerperal infections such as: infection of obstetric surgical wounds, infection of other genital tract including cervicitis and vaginitis, urinary tract infection following delivery, pyrexia of unknown origin following delivery and extra-genitourinary complications related to the puerperium.⁽⁴⁾

Puerperal sepsis is the 3rd leading cause of direct maternal death in ~~the~~ developing countries.⁽¹⁾ The case fatality rate of puerperal sepsis was reported to be as high as 8%,⁽⁵⁾ and it complicates 1-8% of all deliveries.⁽⁶⁾ Furthermore, the risk factors for puerperal sepsis include: low socioeconomic status, low literacy level, un-booked status, prolonged rupture of membranes, prolonged labor, obstructed labour, repeated vaginal examination during labour, anemia in pregnancy, caesarean section, home delivery and poor hygiene.^(1,7,8)

Regarding causative agents, puerperal sepsis is usually poly-microbial comprising of aerobic and anaerobic organisms.⁽⁶⁾ The commonly isolated organisms in Nigeria are Escherichia coli, staphylococcus aureus and Klebsiella species; which are often indigenous to the lower genital tract.^(2, 8) However exogenous organisms, including Niesseriagonorrhoea and chlamydia trachomatis are also isolated.⁽²⁾

~~Similarly, P~~ Puerperal sepsis if untreated ~~may~~ could result in several complications. The early complications include progression of the infection resulting in pelvic abscess, septicaemia, septic shock, renal failure and multiple organ failure leading to death.⁽⁹⁾ Long term complications include

chronic pelvic pain, chronic Pelvic Inflammatory Disease (PID), ectopic pregnancy, secondary amenorrhoea and infertility.⁽⁹⁾ Therefore, the aim of this study was to determine the incidence of puerperal sepsis and associated complications at Rasheed Shekoni Teaching Hospital, (RSTH), Dutse., Jigawa State, Nigeria

MATERIAL AND METHODS

Background to study setting

This is a retrospective study carried out in the department of Obstetrics and Gynaecology of Rasheed Shekoni Teaching hospital, Dutse Jigawa State between 1st June 2021 and 31st May 2023. Rasheed Shekoni Teaching hospital is one of the tertiary health facilities in the state. It serves as a referral center for patients within Jigawa and neighbouring states.

The study participants were all the patients admitted into the postnatal ward and managed for puerperal sepsis during the period under review. The case files of the participants were retrieved from the record department. Relevant information obtained from the files included socio-demographic data, parity, booking status, mode of delivery, place of delivery, presenting complaints, complications and microbiology result.

The data obtained was checked for completeness and accuracy. Data analysis was carried out using IBM statistical package for social sciences (SPSS) version 21.0. Measured variables were expressed in percentage. Test for association was done using chi-square non-parametric test, setting P-value at <0.05.

The study was carried out after obtaining approval from the Health Research and Ethics committee of Rasheed Shekoni Teaching Hospital Jigawa, Nigeria (Approval number- RSSH/GEN/226/V.I)

RESULTS

There were 816 obstetric admissions during the period under review, of which 88 patients were managed for puerperal sepsis. This gave an incidence of 10.7%. However, only 75(85.2%) case files were retrieved for analysis. The mean age of the patients was 25.85 ± 7.5 , with a range of 17 — 40 years. Majority

Comment [mo12]: Please the information provided is good but renirch its contents. What makes it a problem to research, any gap identified. at least 800 wards would be enough. This is very scanty, top up please to reflect on the study phenomenon

Formatted: Font color: Red

Comment [mo13]: Which type pls

Comment [mo14]: I think authors did not follow journal format in writing the methodology section, pls strictly adhere to the standard of journal write up. Very, Very necessary here

Formatted: Font color: Red

Most of the patients were not educated (68%), not employed (88%) and resided in the rural areas (77.3%).

The socio-demographic characteristics are presented in Table 1.

Comment [mo15]: Talk on the age, level of education, occupation as well pls.

Table 1: Socio-demographic characteristics of the participants

Variables	Frequency (n)	Percentage (%)
I-Age		
Mean age= 25.8 ± 7.5		
<20	22	29.3
20-25	22	29.3
26-30	10	13.3
31-35	18	24.0
>35	3	4.0
II-Education level		
No formal education	51	68.0
Primary	8	10.6
Secondary	14	18.7
Tertiary	2	2.7
III-Occupation		
Unemployed	66	88.0
Student	2	2.7
Civil servant	5	6.6
Business	2	2.7
IV-Residence		
Rural	58	77.3
Urban	17	22.7

Comment [mo16]: This is not type of occupation, please change to others

The Obstetric characteristics of the participants are shown in Table 2. The mean parity was 3.89 ± 4.5 with a parity range of 1 – 10. The occurrence of puerperal sepsis was almost equally distributed among the parity groups. Majority (74.71%) of the women were un-booked and had home delivery (69.3%), supervised by unskilled attendants (60%). Most (84.0%) of the women had vaginal delivery.

Table 2: Obstetrics characteristics of the study participants

Variables	Frequency (n)	Percentage (%)
I-Parity		
Mean parity= 3.89 ± 4.5		
Primiparous	23	30.7
Multiparous	24	32.0
Grand-multiparous	28	37.3
II-Booking status		
Un-booked	56	74.7
Booked:	19	25.3
a-booked at study hospital	2	10.5
b-booked at referring hospital	5	26.3
c-booked elsewhere	12	63.2
III- Place of delivery		
Home	52	69.3
Hospital:	23	30.7
a-study hospital	2	8.7
b-other hospitals	21	91.3
IV-Duration of labour		
<24 hours	16	21.3
>24 hours	27	36.0
Not known	32	42.7
V-Mode of delivery		
Vaginal	63	84.0
Caesarean section	12	16.0
VI-Accoucheur		
Unskilled birth attendant	45	60.0
Midwife/Nurse/Chew	18	24.0
Doctor	12	16.0

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

CHEW_{hew} = Community _Hhealth _Eextension _Wworker

Fever (90.7%) was the most frequent presenting complaints. AlsoAgain, delay in health seeking is observed with more than half (58.7%) of the patients presenting to the facility more than a week following the onset of symptoms. Majority (89.3%) of the patient did not have vaginal or wound swab microscopy, culture and sensitivity. Among the eight (10.7%) patients who had swabs taken for culture, 2 (25.0%) yielded no growth. The most common (75.0%) organism cultured was staphylococcus aureus. The isolated organisms were most sensitive to (83.3%) to levofloxacin. The findings are detailed in Table 3.

Table 3: Presenting complaints and microbial culture

Presenting complaint	Frequency (n)	Percentage (%)
Fever	68	90.7
Abdominal pain	41	54.7
Abnormal vaginal discharge	22	29.3
Abdominal distension	17	22.7
Vaginal bleeding	7	9.3
Uterine sub-involution	4	6.7
Duration of symptoms at time of presentation		
1-7 days	31	41.3
>7 days	44	58.7
Culture and sensitivity		
Not done	67	89.3
Done	8	10.7
Culture result n=8		
Staphylococcus	6	75.0
Klebsiella spp	4	50.0
Streptococcus spp	4	50.0
Escherichia coli	2	25.0
No growth	2	25.0
Sensitivity pattern n=6		
Levofloxacin	5	83.3
Ofloxacin	4	66.7
Ciprofloxacin	3	50.0
Cefuxime	3	50.0
Erythromycin	2	33.3

Comment [mo17]: Check spelling please

Azithromycin	1	16.6
--------------	---	------

Comment [mo18]: You can present this section better if you use graph please.)line graph would be ideal

As shown in Table 4; more than half (58.7%) of the women had complications; of which anaemia was the commonest (63.6%). In the same vein 5 (6.7%) maternal death was recorded.

Comment [mo19]: Check spelling please (anaemia)

Table 4: Outcome of puerperal sepsis among participants

Variables	Frequency (n)	Percentage (%)
I-Outcome:	n=75	
Complications	44	58.7
No complication	31	41.3
II-Complications:	n=44	
Anaemia	28	63.6
Abdomino-pelvic abscess	10	22.7
Surgical site infection	8	18.2
Acute kidney injury	5	11.4
Septicaemia	2	4.5
Septic shock	1	2.3
Psychosis	1	2.3
III-Discharge summary		
Discharged	58	77.3
DAMA	12	16.0
Death	5	6.7

DAMA= Discharged against medical advice

DISCUSSION

The incidence of puerperal sepsis is 10.7% in this study. This is similar to 9.34% observed in Port Harcourt, south-south Nigeria.⁽¹⁰⁾ However, the incidence observed in this study is higher than the 0.9% reported in Sokoto, north-west Nigeria. On the contrary, the incidence is lower than the 16.7% in Jos, north-central Nigeria.⁽¹²⁾ This variation in incidence may reflect health seeking behavior and access to health facilities in these areas.

Comment [mo20]: Present general introduction t the study. indicate the significant findings and processed with your discussion please. Very necessary here

The mean age of the patient in this study was 25.8% ± 7.5. This is similar to the mean age recorded in Sokoto. ⁽¹¹⁾ Puerperal sepsis was highest among the age groups of <20 and 20 – 25. A significant proportion (68%) of the patient had no formal education. This is similar to the finding of **some studies**.^{(11,}

Comment [mo21]: Which study please, where was it conducted

¹³⁾ However, this is in contrast with the findings of Oriji et al where majority (60.6%) had secondary education.⁽¹⁴⁾ More so, most of the patients (88%) were unemployed as also reported in **some studies**.^{(11,}

Formatted: Font color: Red

¹³⁾ Low literacy level and low socio-economic status are significant risk factors for puerperal sepsis⁽¹¹⁾ and have been shown to have effect on health seeking behavior.⁽¹⁵⁾ Likewise, most (77.3%) of the patients reside in the rural areas. This is in line with the findings by Signh et al and Demisse **et al. et al.**^(16, 17) This is in contrast to a study from Bangladesh. ⁽¹⁸⁾ High incidence of puerperal sepsis among rural dwellers may be as a result of unclean home delivery, low literacy level and poor awareness of the need for antenatal care.

Majority (74.7%) of the patients did not book for **ANC**. This is similar to the findings of **some studies**.^(10, 19)

Comment [mo22]: Write in full please

Pregnant women during **ANC** are educated on danger signs of pregnancy and need to present to hospital when problem arise. The parity of these patients with puerperal sepsis in this study was almost evenly distributed. This is in contrast to the findings of some studies^(11, 16, 20) The current study found that 69.3% of the patients had home delivery, supervised by unskilled attendants. This is consistent with the findings of **other studies**.^(11, 14, 16, 17) Home deliveries increase the risk of puerperal sepsis due to ignored asepsis,

Formatted: Font color: Red

Comment [mo23]: As above

contact with contaminated materials and some harmful traditional practices. In the study sub-region, cultural values and husband's or his family's choice is also a determinant of place of delivery,⁽²¹⁾ which together with low literacy level and poverty contribute to high incidence of home delivery. In the current study only 23(30.7%) patients managed for puerperal sepsis had deliveries in the hospital; and among them only 2(8.7%) delivered in the study hospital. Others (91.3%) had their deliveries at other primary and secondary health facilities. Referred cases are more likely to develop puerperal sepsis when compared to those from the study hospital. This was similarly observed in Uganda and Pakistan.^(11, 20) A major proportion (84%) of the patients had vaginal delivery and this is similar to the reports of some studies.^(11, 16) More so, the patients who were in labour for more than 24hours (36%) had higher prevalence of the disease and this also share similarity with some studies in Ethiopia and Nepal^(17, 22)

Comment [mo24]: As above pls

Fever (90.7%) was the most frequent presenting complaints. Regarding laboratory confirmation of the diagnosis, most (89.3%) of the patients did not have swab taken for microscopy, culture and sensitivity (MCS). For those that had the swab MCS, the most common (75.0%) organism cultured was staphylococcus aureus. This is similar to findings in Maiduguri and Sudan. ^(19, 25) However this is in contrast to the findings of some studies in which Klebsiella was the most isolated organism. ^(10, 16, 23) Moreover, the isolated organisms were most sensitive to Levofloxacin (83.3%) which was in contrast to the findings in Sokoto (Ceftriaxone), ⁽¹¹⁾ Yenagoa (Amoxicillin – Clavulanic acid) ⁽¹⁴⁾ and India (Gentamycin).⁽¹⁶⁾

Similarly more than half (58.7%) of the patient managed developed or presented with complications; with anaemia being the commonest (63.6%). Also Again this study reported 5(6.7%) maternal death. Oriji et al in their study did not record any mortality. ⁽¹⁴⁾ However, Sulaiman et al had a higher (15.1%) mortality. ⁽¹¹⁾ The patients' status on referral was contributory to the mortality: Majority (97.3%) of these patients were referred from other health facilities. Additionally, more than half (58.7%) of these patients presented more than a week following the onset of symptoms with already developed complications. Additionally, some (16%) of the patients requested and were discharged against medical advice (DAMA). This was because of the medical bills in the study (tertiary) hospital and unusual feeding expenses for the accompanying relatives. These families had enjoyed free maternal health care services (including caesarean section) in all the government owned primary and secondary health care facilities in the sub-region.

Nevertheless, majority (77.3%) of these patients were discharged from the hospital after full recovery.

LIMITATION

This is a hospital based study as such does not reflect what is obtainable in other government owned and private hospitals where the socio-economic status and place of residence may differ.

CONCLUSION

The incidence of Puerperal sepsis in this study is high. Low literacy level, unemployment, unbooked status and home delivery were found to be predisposing factors. Making health care affordable through poverty alleviation support, in addition to increase awareness on need for ANC where pregnant women will be given effective health education on safe delivery and postpartum care could help in reducing the incidence of puerperal sepsis in the study community.

Comment [mo25]: In full pls

Conflict of interest

Funding for the study

Competing Interest

Contributions of authors

Formatted: Justified, Line spacing: Double

REFERENCES

1. Say L, Chou D, Gemmil A, et al. Global causes of material death: a WHO systematic analysis. Lancet Glob Health 2014; 2:323-33.
2. World Health Organization. WHO recommendations for prevention and treatment of material peripartum infections. World Health Organization 2015; 80.
3. Momoh MA, Ezurgwuwie OJ, Ezeigwei HO. Causes and management of puerperal sepsis: The health personnel view point. Advanced Bio Research 2010; 4:154-8.
4. International Statistical Classification of Diseases and Related Health Problems 10th Revision (ICD-10) – 2015-WHO Version, 2015. Complications predominantly related to the puerperium (085-092).
5. Greer O, Shah NM, Sriskandan S, et al. Sepsis: precision-based medicine for pregnancy and puerperium. Int J Mol Sci 2019; 20: 5388.
6. Tamboli SS, Tamboli SB, Shrikhande S. Puerperal Sepsis: predominant organisms and their antibiotic sensitivity pattern. Int J Reprod contracept obstetGynecol 2016; 5:762-5.

7. Kiponza R, Bulandya B, Majigo MV. Laboratory confirmed puerperal sepsis in a national referral hospital in Tanzania: etiological agents and their susceptibility to commonly prescribed antibiotics. *BMC Infect Dis* 2019; 19:690.
8. Melkie A, Dagnaw E. Burden of puerperal sepsis and its associated factors in Ethiopia: a systematic review and meta-analysis. *Arch Public Heal* 2021; 79:1-11.
9. Atlaw D, Seyoum K. Puerperal sepsis and its associated factors among mothers in University of Gondar referral hospital, Ethiopia. *Int J Preg Child Birth* 2019; 5:190-5.
10. Ononuju CN, Nyengidiki TK, Ugboma HA, Bassey G. Risk factors and antibiogram of organisms causing puerperal sepsis in a tertiary health facility in Nigeria. *Trop J ObstetGynaecol* 2015; 32:73-82.
11. Sulaiman B, Tunau KA, Nasir S, Hassan M, Ahmed Y. Puerperal sepsis at Usmanu Danfodiyo University Teaching Hospital, Sokoto: A ten-year review. *Eur J Pharm Med Res* 2018; 5:569-73.
12. Mutahir J, Utoo B. Postpartum maternal morbidity in Jos, North-central Nigeria. *Niger J Clin Pract* 2011; 14:38.
13. Maritim CV, Jackim N, Margaret K. Associated factors with puerperal sepsis among reproductive age women in Nandi country, Kenya. *J Midwifery Reprod Health* 2017; 5:1032-40.
14. Oriji PC, Allagoa DO, IKoroc, Oguche IO, Oriji CE, Unachukwu CE. A Five Year Review of Puerperal sepsis and its complications at the Federal Medical Centre, Yenagoa, south-south Nigeria. *J Adv Microbiol* 2021; 21:55-63.
15. Ahmed MI, Alsammani MA, Babiker R. Puerperal Sepsis in a rural hospital in Sudan. *Mater Socio Med* 2013; 25:19-22.
16. Singh P, Tirkey S, Trivedi K, Hansda R, Prakesh J. Study of cases of puerperal sepsis, its socio-demographic factors, bacterial isolates and antibiotics sensitivity pattern. *J Family Med Prim Care* 2022; 11: 5155-60.
17. Demisse GA, Sifer SD, Kedir B, Fekene DB, Bulto Ga. Determinants of puerperal sepsis among postpartum women at public hospitals in west zone oromia regional state, Ethiopia. *BMC Pregnancy Childbirth* 2019; 19:95.

18. Taskin T, Sultana M, Islam T, Khan NA, Chowdhury S.M. Socio-demographic factors and puerperal Sepsis: experiences from two tertiary level hospitals in Bangladesh Int J Community Fam Med 2016; 1:1-4.
19. Bako B, Audu BM, Lawan ZM, Umar JB. Risk factors are microbial isolates of puerperal sepsis at the University of Maiduguri Teaching Hospital Maiduguri, North-eastern Nigeria ArchGynecolObstet 2012; 285:913-17.
20. Ngozi J, Tornes YF, Mukasa Pk, Salango W, Kabakyenga J, Sezalio M et al. Puerperal sepsis, the leading cause of maternal deaths at a Teaching University teaching Hospital in Uganda. BMC Pregnancy Childbirth 2016; 16:207.
21. Ekele BA, Tunau KA. Place of delivery among women who had antenatal care in a teaching hospital. Acta ObstetGynecol Scand 2007; 86:627 – 30.
22. Bhandary S Puerperal Sepsis and its cause in Patan hospital Nepal J ObstetGynecol 2015; 1:33-5.
23. Vanukuru J, Bagga R, Muthyala T, Gautam V, Sethi S, Jain V et al. A clinical and microbiological study of puerperal sepsis in tertiary care hospital in India. J Matern Fetal Neonatal Med 2019; 32:1931-7.