

# Computing the minimum polynomial, the function and the Drazin inverse of a matrix with Matlab

## Abstract

In this note we will discuss the known (but not well known) problem of finding the minimum polynomial and the function of a matrix providing the simplest proofs for undergraduate students. We will try to explain with fairly simple arguments how to compute the minimum polynomial of a matrix giving also the `matlab` code for its symbolic computation. Next we will describe the (symbolic) computation of the matrix of a function via the Hermite interpolation method which seems to be the simplest method for undergraduate students. Finally we shall see how we can compute the Drazin inverse given the  $n$ th power of a matrix  $A$ .

**ACM Classification** Mathematics of computing-Solvers

**Keywords** Minimum polynomial of a matrix, function of a matrix, MATLAB codes

## 1 Introduction

A problem that arise at least at the study of Markov chains is to compute the  $n$ th power of a stochastic matrix or the exponential matrix  $e^{tA}$ . A well known approach is via the diagonalization of the matrix  $A$  but this approach need the matrix  $A$  to be diagonalizable. Below, we will describe the Hermite interpolation method which seems to be the simplest way for undergraduate students. For more about the matrix function computation one can refer to the bibliography given at the end of this note (see for example [5], [7], [11]).

Let  $A_{k \times k}$  a matrix of real or complex numbers. If we want to compute symbolically the matrix  $A^n$  we can do the following. Find a polynomial  $p(x) = x^q + c_{q-1}x^{q-1} + \dots + c_0$  which is such that

$$p(A) = A^q + c_{q-1}A^{q-1} + \dots + c_0\mathbb{I}_{k \times k} = 0_{k \times k}$$

Dividing theoretically the polynomial  $x^n$ , for  $n \geq q$ , by  $p(x)$  we have

$$x^n = \pi(x)p(x) + v(x), \quad \forall x \in \mathbb{R}$$

where  $v(x)$  is a polynomial with degree at most  $q - 1$ . That means that the polynomial  $\Delta(x) = x^n - \pi(x)p(x)v(x)$  has all its coefficients equal to zero and that means that

$$A^n = v(A)$$

In general,  $v(A) = A^n$  for  $n \in \mathbb{N}$  except in the case where the polynomial  $p(x)$  has the zero number as its root with multiplicity greater than one. In this case it holds  $v(A) = A^n$  at least for  $n \geq q$ .

The only thing that we have to do is to find the coefficients of  $v(x)$ . In this problem, the roots of the polynomial  $p(x)$  will play an important role. If  $p_1, \dots, p_l$  are the roots with multiplicity  $j_1, \dots, j_l$  (with  $j_1 + \dots + j_l = q$ ) then we construct the following system, setting  $f(x) = x^n$ ,

$$v^{(r)}(p_i) = f^{(r)}(p_i), \quad r = 0, \dots, j_i - 1, \quad i = 1, \dots, l \quad (1)$$

where  $f^{(r)}$  is the  $r$  derivative of  $f$  and  $f^{(0)} = f$ .

Let us look the homogeneous system 1. According to the fundamental theorem of algebra any polynomial of degree  $n$  has exactly  $n$  complex roots not distinct in general. The degree of the polynomial  $v(\cdot)$  equal to  $q - 1$  therefore has only  $q - 1$  roots. However, from the homogeneous system 1 we deduce that the polynomial  $v(\cdot)$  has  $q$  roots and that is true only if the polynomial  $v(\cdot)$  is identically zero. Therefore the above homogeneous system has only the trivial solution and that means that the system 1 has a unique solution.

One polynomial with the property  $p(A) = 0_{k \times k}$  is the characteristic polynomial of  $A$ . The system 1 suggests that the less the degree of the polynomial  $p(x)$  the better. That means that we should find, if it is possible, the minimum polynomial with the property  $p(A) = 0_{k \times k}$ .

## 2 The computation of the minimum polynomial

Let an  $k \times k$  matrix of numbers  $A$ . Below we give the `matlab` function in order to compute the minimum polynomial of this matrix. The coefficients of the polynomial are in the vector  $v$ , i.e. the minimum polynomial is

$$m(x) = v(1)x^r + v(2)x^{r-1} + \dots + v(r+1)$$

```
function v=minimpoly(A)
[l,n]=size(A);
if l~=n
    fprintf('This is not a square matrix')
    return
end

r=1;
Aj=sym(eye(1));
B=Aj(:);
BB(1,1)=sym(1);
while BB(r,r)==1
    Aj=Aj*A;
    B=[B Aj(:)];
    BB=rref(B);
    r=r+1;
end
v=[1 transpose(flip(-BB(1:r-1,r)))];
end
```

Where this code come from? We shall explain below the mathematics beyond the above code (see for example [1], [16]).

**Definition 1** By  $P_A$  we denote all the monic polynomials which are such that  $p(A) = 0_{k \times k}$ .

It is well known that there exists the minimum monic polynomial (which is unique) of the matrix  $A$  that belongs to  $P_A$  and the proof can be found at any linear algebra book (see for example [2], [9], [10], [15]). We denote the minimum polynomial of  $A$  by  $m_A(x)$  and its degree by  $q_{m_A}$ .

**Definition 2** By  $b_A(q)$  we denote the vector which contain the matrix  $A^q$  column by column. By  $b_A(0)$  we denote the vector which contain the identity matrix column by column. By  $B_A(q)$  we denote the matrix with columns  $b_A(0), \dots, b_A(q)$ .

**Definition 3** By  $C_A(q)$  we denote the following set, given  $q \in \mathbb{N}$ ,

$$C_A(q) = \{c \in \mathbb{R}^q : B_A(q-1) \cdot c = -b_A(q)\}$$

**Proposition 1** If  $p(x) = x^q + c_{q-1}x^{q-1} + \dots + c_0$  and  $c = \begin{pmatrix} c_0 \\ \vdots \\ c_{q-1} \end{pmatrix}$  then

$$p(x) \in P_A \iff c \in C_A(q)$$

**Proof.** Let  $p(x) \in P_A$ . That means that

$$p(A) = A^q + c_{q-1}A^{q-1} + \dots + c_0\mathbb{I}_{k \times k} = 0_{k \times k}$$

In other words

$$c_{q-1}A^{q-1} + \dots + c_0\mathbb{I}_{k \times k} = -A^q \tag{2}$$

The above set of equations can be written at the following form

$$B_A(q-1) \cdot c = -b_A(q) \tag{3}$$

where  $c = \begin{pmatrix} c_0 \\ \vdots \\ c_{q-1} \end{pmatrix}$ . That is the systems 2 and 3 are equivalent and therefore the desired conclusion is true. □

**Definition 4** We denote by  $Q_A$  the following set of integers

$$Q_A = \{q \in \mathbb{N} : C_A(q) \neq \emptyset\}$$

**Theorem 1** Let  $m(x)$  the minimum polynomial of  $A$  with degree  $q_{m_A}$ . Then

$$q_{m_A} = \min Q_A$$

**Proof.** The set  $Q_A$  is nonempty because for  $q = k$  the  $C_A(q)$  contain the coefficients of the characteristic polynomial. Therefore the  $q^* = \min Q_A$  is well defined and of course  $q^* \in Q_A$ . That means that there exists a monic polynomial  $p_{q^*}(x)$  of degree  $q^*$  which is such that  $p_{q^*}(A) = 0_{k \times k}$ .

We know that the minimum polynomial exists and is unique therefore  $q_{m_A} \in Q_A$  so  $q_{m_A} \geq q^*$ . This immediately mean that  $q_{m_A} = q^*$  using proposition 1.  $\square$

Therefore in order to compute the minimum polynomial we have to find the minimum  $q$  for which the system  $B_A(q-1) \cdot c = -b_A(q)$  has a solution.

We have compared the above matlab code with the integrated function `minpoly` and see that in many cases is much faster.

### 3 The function and the Drazin inverse of the matrix $A$

Below we shall give two equivalent definitions of the function of a matrix. Before that we define the matrix  $v(A)$  where  $v(\cdot)$  is a polynomial.

**Definition 5** *Let a polynomial  $v(x) = a_q x^q + \dots + a_0$ . The matrix  $v(A)$  is defined to be the*

$$v(A) = a_q A^q + \dots + a_0 \mathbb{I}$$

**Definition 6 (Hermite Interpolation)** *Let a matrix  $A_{k \times k}$  and a function  $f : U \subseteq \mathbb{R} \rightarrow \mathbb{R}$ . The matrix  $f(A)$  is defined to be the  $v(A)$  where  $v(\cdot)$  is the polynomial which satisfies the system 1, given a polynomial  $p(x)$  which is such that  $p(A) = 0_{k \times k}$  and  $p_1, \dots, p_l$  are the roots with multiplicities  $j_1, \dots, j_l$  of  $p(x)$ .*

**Definition 7 (Taylor Expansion)** *Let a matrix  $A_{k \times k}$  and a function  $f : U \subseteq \mathbb{R} \rightarrow \mathbb{R}$ . The matrix  $f(A)$  is defined to be the  $v(A)$  where the  $v(\cdot)$  is the Taylor expansion of  $f$  around zero.*

It is not always defined the matrix  $f(A)$  as we have discussed in [6].

**Definition 8** *Let a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a polynomial  $p(x)$  with  $p_1, \dots, p_l$  are the roots of multiplicities  $j_1, \dots, j_l$  of  $p(x)$ . We say that the function  $f$  is well defined at the roots of  $p(x)$  if  $f^{(k)}(p_i)$ , for  $i = 1, \dots, l$  and  $k = 0, \dots, j_i - 1$  are well defined.*

**Proposition 2** *Let a matrix  $A_{k \times k}$  and set  $\Delta = \max_{i,j} |A_{ij}|$ . If the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  can be expanded in a Taylor series around zero which is absolute convergent for any  $x \in [-M, M]$  with  $M > k\Delta$  then the matrix  $f(A)$  is well defined by definition 7. Let a polynomial  $p(x)$  which is such that  $p(A) = 0_{k \times k}$ . Then the matrix  $f(A)$  is well defined by the definition 6 if the function  $f$  is well defined at the roots of  $p(x)$ .*

**Proof.** The matrix  $f(A)$  via the Taylor expansion is defined as

$$f(A) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} A^n$$

This matrix is well defined because the infinite series converges. Indeed, define by  $\Delta = \max_{i,j} |A_{ij}|$  we have that

$$|(A^n)_{ij}| \leq k^{n-1} \Delta^n$$

We will prove it by induction. For  $n = 1$  is obvious that  $|(A^1)_{ij}| \leq k^0 \Delta$ . We assume that it holds for some  $n$ , that is  $|(A^n)_{ij}| \leq k^{n-1} \Delta^n$  and we will prove that it holds for  $n + 1$ . We have that

$$|(A^{n+1})_{ij}| = \left| \sum_{l=1}^k (A^n)_{il} A_{lj} \right| \leq \sum_{l=1}^k k^{n-1} \Delta^n A_{lj} \leq k^n \Delta^{n+1}$$

Therefore

$$\left| \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} (A^n)_{ij} \right| \leq 1 + \left| \sum_{n=1}^{\infty} \frac{f^{(n)}(0)}{n!} (A^n)_{ij} \right| \leq 1 + \sum_{n=1}^{\infty} \frac{|f^{(n)}(0)|}{n!} k^{n-1} \Delta^n$$

That means that  $\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} (A^n)_{ij}$  converges absolutely and therefore the matrix  $f(A)$  is well defined.

On the other hand it is easy to see that the matrix  $f(A)$  via definition 6 is well defined as soon as the right hand side of 1 is well defined.  $\square$

**Remark 1** *It is easy to see that the matrices  $f(A), g(A), h(A)$  are well defined for any matrix  $A_{k \times k}$  for  $f(x) = e^{tx}$ ,  $g(x) = \cos(tx)$  and  $h(x) = \sin(tx)$ .*  $\square$

**Theorem 2** *Let a function  $f$  and a matrix  $A_{k \times k}$ . Suppose that the matrix  $f(A)$  is well defined for both definitions 7 and 6. Then the two matrices are equal.*

**Proof.** Let  $q$  the degree of the polynomial  $p(x)$  which is such that  $p(A) = 0_{k \times k}$  and the roots  $p_1, \dots, p_l$  with multiplicities  $j_1 \dots, j_l$ . For any  $m \in \mathbb{N}$  with  $m \geq q$  there are  $a_0(m), \dots, a_{q-1}(m)$  such that

$$A^m = a_{q-1}(m)A^{q-1} + \dots + a_0(m)\mathbb{I}$$

Then it is easy to see that,

$$\begin{aligned} f^{Taylor}(A) &= A^{q-1} \underbrace{\left( \sum_{n=q}^{\infty} \frac{f^{(n)}(0)}{n!} a_{q-1}(n) + \frac{f^{(q-1)}(0)}{(q-1)!} \right)}_{b_{q-1}} \\ &+ \dots + \mathbb{I} \underbrace{\left( \sum_{n=q}^{\infty} \frac{f^{(n)}(0)}{n!} a_0(n) + \frac{f^{(0)}(0)}{0!} \right)}_{b_0} \end{aligned}$$

In order to get the above result we have rearranged the terms and this can be done since the series is absolutely convergent.

We shall see below that the same coefficients  $b_0, \dots, b_{q-1}$  will appear if we compute the matrix  $f(A)$  via the Hermite interpolation. Computing  $A^m$  for  $m \geq q$  the system 1 will contain the following equations, for  $i = 1, \dots, l$ ,

$$\begin{aligned} a_{q-1}(m)p_i^{q-1} + \dots + a_0(m) &= p_i^m \\ a_{q-1}(m)(q-1)p_i^{q-2} + \dots + a_1(m) &= mp_i^{m-1} \\ &\vdots \\ a_{q-1}(m)(q-1) \dots (q-j_i+1)p_i^{q-j_i} + \dots + a_{j_i-1}(m) &= \\ m \cdot (m-1) \dots (m-j_i+2)p_i^{m-j_i+1} & \end{aligned}$$

We multiply these equations by  $\frac{f^{(m)}(0)}{m!}$  and then we sum from  $m = q$  to  $\infty$ . Therefore the above system will contain the following equations, for  $i = 1, \dots, l$ ,

$$\begin{aligned}
 p_i^{q-1} \sum_{m=q}^{\infty} a_{q-1}(m) \frac{f^{(m)}(0)}{m!} + \dots + \sum_{m=q}^{\infty} a_0(m) \frac{f^{(m)}(0)}{m!} &= \sum_{m=q}^{\infty} p_i^m \frac{f^{(m)}(0)}{m!} \\
 (q-1)p_i^{q-1} \sum_{m=q}^{\infty} a_{q-1}(m) \frac{f^{(m)}(0)}{m!} + \dots + \sum_{m=q}^{\infty} a_1(m) \frac{f^{(m)}(0)}{m!} &= \sum_{m=q}^{\infty} m p_i^{m-1} \frac{f^{(m)}(0)}{m!} \\
 &\vdots \\
 (q-1) \dots (q-j_i+1) p_i^{q-j_i} \sum_{m=q}^{\infty} a_{q-1}(m) \frac{f^{(m)}(0)}{m!} + \dots + \\
 &\quad + \sum_{m=q}^{\infty} a_{j_i-1}(m) \frac{f^{(m)}(0)}{m!} = \\
 &\quad \sum_{m=q}^{\infty} m \cdot (m-1) \dots (m-j_i+2) p_i^{m-j_i+1} \frac{f^{(m)}(0)}{m!}
 \end{aligned}$$

We add the corresponding terms in each equation so the right hand side equal to  $f(p_i)$ ,  $f'(p_i)$ ,  $\dots$ ,  $f^{(j_i-1)}(p_i)$ . Then the unknown coefficients of the above system are the  $b_0, \dots, b_{q-1}$  and therefore we obtain the desired result.  $\square$

The `matlab` function for this computation is `matfun(A,f)` and can be found at the following link <https://www.mathworks.com/matlabcentral/fileexchange/132518-computing-the-minimum-polynomial-of-a-matrix>. At this link one can find several auxiliary functions and examples.

The Drazin inverse of a matrix can be used at the context of differential equations, Markov chains and others (see for example [3], [4], [8], [12]).

**Theorem 3** *Let a matrix  $A_{k \times k}$  and let  $a_{ij}(n)$  the elements of  $A^n$ . Then the matrix  $B$  with elements the numbers  $a_{ij}(-1)$  is the Drazin inverse of the matrix  $A$ .*

**Proof.** See remark 3.2 of [6].  $\square$

Therefore, one way to compute the Drazin inverse of a matrix is to compute the  $n$ th power of the matrix and then set  $n = -1$ . Of course if the matrix is invertible then setting  $n = -1$  we will get its inverse.

**Remark 2** *In all the above computations we can use the characteristic polynomial, the minimum polynomial or any polynomial  $p(x)$  with the property  $p(A) = 0_{k \times k}$ . The result will be the same as we have seen in [6].*

*Indeed, let  $v(x)$  and  $\hat{v}(x)$  the polynomials that satisfies system 1 using the characteristic polynomial to get  $v(x)$  and a polynomial  $p(x)$  (with  $p(A) = 0_{k \times k}$ ) to get  $\hat{v}(x)$ . Note that the polynomial  $v(x)$  is of degree  $k-1$  and suppose that the polynomial  $p(x)$  is of degree less or equal  $k-1$ . Dividing  $v(x)$  by  $p(x)$  we get*

$$v(x) = \pi(x)p(x) + q(x)$$

*Using the roots (and their multiplicities) of  $p(x)$  we can compute the polynomial  $q(x)$  and see that  $q(x) = \hat{v}(x)$ . Moreover,  $\hat{v}(A) = v(A) = f(A)$ .*  $\square$

**Remark 3** *Under the above point of view two things can be changed in order to increase the accuracy and decrease the time of computations. The first is to find a better way to compute the minimum polynomial and the second is to find a better way to solve the confluent Vandermonde linear system (see for example [13], [14]).*  $\square$

## 4 Examples

We will compute the  $n$ th power of a non invertible matrix and then its Drazin inverse. Let the matrix

$$C = \begin{pmatrix} \frac{1}{5} & \frac{2}{5} & \frac{2}{5} \\ \frac{3}{10} & \frac{3}{5} & \frac{1}{10} \\ \frac{1}{10} & \frac{1}{5} & \frac{7}{10} \end{pmatrix}$$

Note that  $C$  is not invertible. Setting  $f(x) = x^n$  we can use our function `matfun(C,f)` in order to compute the  $n$ th power of  $C$ . We get the following result

$$C^n = \begin{pmatrix} \frac{1}{5} & \frac{2}{5} & \frac{2}{5} \\ \frac{2^n+1}{5 \cdot 2^n} & \frac{2(2^n+1)}{5 \cdot 2^n} & \frac{2 \cdot 2^n - 3}{5 \cdot 2^n} \\ \frac{2^n-1}{5 \cdot 2^n} & \frac{2(2^n-1)}{5 \cdot 2^n} & \frac{2 \cdot 2^n + 3}{5 \cdot 2^n} \end{pmatrix}$$

Setting  $n = -1$  we get the Drazin inverse of the matrix  $C$  which is the following

$$C_D = \begin{pmatrix} \frac{1}{5} & \frac{2}{5} & \frac{2}{5} \\ \frac{3}{5} & \frac{6}{5} & -\frac{4}{5} \\ -\frac{1}{5} & -\frac{2}{5} & \frac{8}{5} \end{pmatrix}$$

Using our function `isdrazin` we verify that the matrix  $C_D$  is indeed the Drazin inverse of  $C$ . We have taken care in our functions so that we can let  $n = -1$  in order to get the Drazin inverse. This is not the case concerning the integrated function `funm`.

Finally we will compute the  $n$ th power and the exponential of a matrix with complex eigenvalues. Let the matrix  $A$

$$A = \begin{pmatrix} 3 & 2 \\ -1 & 3 \end{pmatrix}$$

Setting  $f(x) = x^n$  and  $g(x) = \exp(t * x)$  and typing `matfun(A,f)` and `matfun(A,g)` we get the following results

$$A^n = \begin{pmatrix} 11^{n/2} \cos\left(n \operatorname{atan}\left(\frac{\sqrt{2}}{3}\right)\right) & \sqrt{2} 11^{n/2} \sin\left(n \operatorname{atan}\left(\frac{\sqrt{2}}{3}\right)\right) \\ -\frac{\sqrt{2} 11^{n/2} \sin\left(n \operatorname{atan}\left(\frac{\sqrt{2}}{3}\right)\right)}{2} & 11^{n/2} \cos\left(n \operatorname{atan}\left(\frac{\sqrt{2}}{3}\right)\right) \end{pmatrix}$$

and

$$e^{tA} = \begin{pmatrix} e^{3t} \cos(\sqrt{2}t) & \sqrt{2} e^{3t} \sin(\sqrt{2}t) \\ -\frac{\sqrt{2} e^{3t} \sin(\sqrt{2}t)}{2} & e^{3t} \cos(\sqrt{2}t) \end{pmatrix}$$

In our matlab functions we have transformed the complex roots of the minimum polynomial into their polar forms in order to arrive at the above results. This is not the case with the integrated function `funm`.

Our functions `isan(an,a)` and `isexpt(at,a)` verify if the matrices `an` and `at` are indeed the  $n$ th power of `a` and the exponential matrix of `a`. For the first case we use induction while for the second case we use the fact that the matrix  $e^{tA}$  is the only matrix which satisfies the following matrix differential equation

$$\begin{aligned} (M(t))' &= AM(t) \\ M(0) &= \mathbb{I} \end{aligned}$$

The fact that the above matrix differential equation has a unique solution can be proved as follows. It is easy to see that the exponential matrix  $e^{tA}$  solves the above matrix differential equation by using the Taylor expansion and differentiating term by term since the Taylor series converges absolutely. Next, if there are two or more solutions then their difference will satisfy the following

$$\begin{aligned}(M(t))' &= AM(t) \\ M(0) &= \mathbf{0}\end{aligned}$$

But  $(M(t)e^{-tA})' = M'(t)e^{-tA} - AM(t)e^{-tA} = \mathbf{0}$ . Therefore  $M(t)e^{-tA} = \mathbf{C}$  where  $\mathbf{C}$  is a matrix independent of  $t$ . Multiplying by  $e^{tA}$ , using the fact that  $e^A e^B = e^{A+B}$  for commutative matrices  $A, B$  and since  $M(0) = \mathbf{0}$  it follows that  $M(t) = \mathbf{0}$  for any  $t \in \mathbb{R}$ .

Below we compare our functions with the integrated functions of `Matlab` for large sparse matrices. We see that increasing the dimension of the matrix the ratio of times increases as well.

```
clear; syms x n t f(x)=x^n; g(x)=exp(t*x);
K=45;
H=zeros(K,K);
H(1,1)=1;
for i=2:K H(i,i)=0.7; H(i,1)=0.3;
end
H(4,1)=0.1;
H(4,2)=0.1; H(4,3)=0.2; H(6,7)=0.4;

%Comparing the nth power and the exponential
tic
matfun(H,f);
matfun(H,g);
t3=toc;

tic simplify(funm(H,f)); simplify(funm(H,g)); t4=toc; t4/t3

%comparing the minimum polynomial computation
tic
ppH=poly2sym(minumpoly(H),x);
pH=solve(ppH==0,x);
t1=toc

tic
ppmH=poly2sym(minpoly(H),x);
pmH=solve(ppmH==0,x);
t2=toc

t2/t1
```

The comparison concerning the computation of the minimum polynomial gives the result  $t2/t1 = 348.17$ .

The comparison concerning the computation of the  $n$ th power and the exponential matrix gives the following  $t4/t3 = 1.39$  using the `matlab` online. We believe that the reason that we get better results is the use of the minimum polynomial in order to do these computations for sparse matrices.

## References

- [1] S. Bialas - M. Bialas, *An algorithm for the calculation of the minimal polynomial*, Bulletin Polish Academy Sciences, vol. 56, 2008.
- [2] S. Berberian, *Linear Algebra*, Oxford Science Publications, 1992.
- [3] S. Cambell - C. Meyer, *Generalized Inverses of Linear Transformations*, SIAM, 2009.
- [4] Stephen L. Campbell, Carl D. Meyer, Jr. and Nicholas J. Rose, *Applications of the Drazin Inverse to Linear Systems of Differential Equations with Singular Constant Coefficients*, SIAM Journal on Applied Mathematics, Vol. 31, 1976
- [5] R. Fløve - G. Harris, *A note on generalized Vandermonde determinants*, SIAM J. Matrix Anal. Appl. vol. 14, No. 4, pp. 1146-1151, 1983.
- [6] N. Halidias, *On the Computation of the Minimum Polynomial and Applications*, Asian Research Journal of Mathematics, 2022.
- [7] N. Higham, *Functions of Matrices, Theory and Computation*, SIAM, 2008.
- [8] A. Israel - T. Grevill, *Generalized Inverses, Theory and Applications*, Springer, 2003.
- [9] R. Larson - D. Falvo, *Elementary Linear Algebra*, Houghton Mifflin Harcourt Publishing Company, 2009.
- [10] S. Lipschutz-M. Lipson, *Linear Algebra*, McGraw Hill, 2009.
- [11] C. Moler - C. Van Loan, *Nineteen dubious ways to compute the exponential of a matrix, twenty - five years later*, SIAM Review, 2003.
- [12] R. Penrose, *A Generalized Inverse for Matrices*, Mathematical Proceedings of the Cambridge Philosophical Society, pp. 406-413, vol. 51, 1955.
- [13] J. Respondek, *Numerical recipes for the high efficient inverse of the confluent Vandermonde matrices*, App. Math. Comp. Vol. 218, Issue 5, pp. 2044-2054, 2011.
- [14] J. Respondek, *Matrix black box algorithms - a survey*, Bulletin of the Polish Academy of Sciences, Vol. 70 (2), p. e140535, 2022.
- [15] H. Ricardo, *A Modern Introduction to Linear Algebra*, CRC Press, 2010.
- [16] V. Shoup, *Efficient Computation of Minimal Polynomials in Algebraic Extensions of Finite Fields*, 1999.