

Original Research Article

SARS-CoV-2 Infection and Factors Associated with Mortality in Hospitalized Patients Aged 60 and Over at the Epidemic Treatment Center (ETC) of Saint-Louis (Senegal)

ABSTRACT

Background: SARS-CoV-2 infection typically causes severe pneumonia in elderly patients with high mortality rates.

Aims: This study aimed to describe the epidemiological, clinical, paraclinical, and evolution profile of the SARS-CoV-2 infection in patients aged 60 years and over hospitalized at the Epidemic Treatment Center (ETC) of Saint-Louis (Senegal) and to identify the factors associated with death.

Methodology: This is a cross-sectional, retrospective, descriptive, and analytical study conducted from March 2020 to August 2021. It includes all patients aged 60 years and older hospitalized at the ETC of Saint-Louis (Senegal) for SARS-CoV-2 infection. **The data were entered into Excel and analyzed using EPI INFO version 7 software.**

Results: Of 448 hospitalizations, 188 patient records aged 60 years and older were collected. The prevalence was 42%. The mean age was 70 years (± 8 years). The sex ratio was 1.7 with a male predominance (62.8%). Hypertension was the most common co-morbidity (67%). Severe forms were predominant (64.4%). Dyspnea (60.3%), cough (56.5%), and ageusia (24.5%) were the most common symptoms on admission. Thoracic CT scans showed severe involvement in 51.7% of cases. Hydroxychloroquine-azithromycin combination was prescribed in 98.4% of cases. The mortality rate was 18.6%. The main factors associated with death were age (≤ 75 years) ($p=0.014$) [OR=0.488; 95% CI: 0.201-1.187]; critical forms ($p=0.017$) [OR=0.061; 95% CI: 0.006-0.601]; comorbidity ($p=0.031$) [OR=1.647; 95% CI: 0.280-2.499]; occurrence of complications ($p=0.009$) [OR=1.212; 95% CI: 0.530-2.774]; and SpO₂ $\leq 95\%$ at admission ($p=0.031$) [OR=4.899; 95% CI: 0.752-8.797].

Comment [RS1]: AIM

Comment [RS2]: Microsoft Excel was used to enter data and then analysis was done using EPI INFO version 7 software.

Conclusion: SARS-CoV-2 infection in the elderly is predominantly severe with high mortality. This high-risk group requires special attention.

keywords: SARS-CoV-2; elderly patients; Saint-Louis; Senegal.

1. INTRODUCTION

COVID-19 (coronavirus disease 19), caused by SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2), is a respiratory illness that often results in respiratory failure [1,2]. The World Health Organization declared a pandemic on March 11, 2020, following its fast global expansion [1]. As of September 27, 2020, there were approximately 32,920,218 confirmed cases and nearly one million deaths worldwide. On the same date, Africa recorded 35,440 deaths from 1,459,714 confirmed cases [3]. Like other African countries, Senegal has not escaped the epidemic and the first case was recorded on March 2, 2020. As of September 24, 2020, Senegal reported 14,816 confirmed cases, with 11,818 recovered and 304 deaths [4,5].

Comment [RS3]: Made changes

SARS-CoV-2 infection is typically mild in 80% of cases, with the most typical clinical presentation being acute respiratory infection with fever, dry cough, dyspnea, fatigue, myalgia, and headache [6-9]. However, 15% of cases are severe and 5% are critical [6,7]. The mortality rate is very high in severe and critical cases, reaching up to 50% in some series [6]. Individuals with comorbidities such as advanced age, hypertension, diabetes mellitus, and chronic respiratory diseases are at a higher risk of developing severe forms of the disease or dying from it [9-11]. According to the World Health Organization (WHO), the age-related risk begins at 40 years of age. However, the Centers for Disease Control and Prevention (CDC) states that the risk of developing severe forms is observed in people aged 65 years and over [12]. In the population aged 60 years and older, the mortality rate associated with SARS-CoV-2 infection is estimated to be 6.38%. This is approximately 20 times higher than the mortality rate observed in persons under the age of 60 (0.32%) [12]. In Senegal, few studies have investigated SARS-CoV-2 infection in elderly people. This study aims to describe the epidemiological, clinical, paraclinical, therapeutic, and evolution profile of SARS-CoV-2 infection in patients aged 60 and over hospitalized at the ETC of Saint-Louis (Senegal), and to identify the factors associated with death.

Comment [RS4]: WHO confirmed that 40 years and above will face age-related risks.

2. MATERIALS AND METHODS

2.1. Type, Period of Study, and Study Population.

This was a cross-sectional, retrospective, descriptive, and analytical study conducted from March 2020 to August 2021, carried out on patients aged 60 and over hospitalized at the Saint-Louis ETC (Senegal).

2.2 Inclusion criteria

It included all patients aged 60 years and older hospitalized at the ETC of Saint-Louis (Senegal) for SARS-CoV-2 infection, regardless of severity, during the three waves of COVID-19.

2.3 Non-inclusion criteria

Patients whose medical records were incomplete or could not be found were not included.

2.4 Diagnostic criteria

The diagnosis of SARS-CoV-2 infection was based on

- a. RT-PCR obtained from nasopharyngeal swabs.
- b. Positive Antigen rapid tests
- c. Chest CT scans

2.5 Data collection

Data was collected from medical records using a questionnaire covering the following parameters: epidemiological data (age, gender, marital status, comorbidities, and COVID-19 vaccination status), clinical data (the time between symptom onset and hospitalization, severity of clinical presentation, peripheral oxygen saturation (SpO₂) at admission, and symptoms at admission), paraclinical data (hemogram, transaminases, creatinine, CRP, fasting blood glucose, blood electrolytes, lipids, troponin, prothrombin levels, and thoracic CT scan), and therapeutic data (molecules administered) and evolution data (length of hospital stay and outcome).

2.6 Statistical analysis

Data was entered and analyzed using Excel and EPI INFO version 7 software. The Chi² and Fisher tests were used to compare means and percentages, respectively, based on their applicability. Statistical significance was defined as any difference below 0.05.

Comment [RS5]: Patients with complete and available medical records were considered during the study.

Comment [RS6]: Changes are introduced here for better understanding of the content.

Comment [RS7]: For parameters include a table format. It adds value and clarity to the study.

Comment [RS8]: The study included usage for Microsoft Excel with EPI INFO version 7 software.

Comment [RS9]: Statistical tools employed for calculating means and percentages are Chi² and Fisher tests based on applicability.

3. RESULTS

3.1 Epidemiological Aspects

Out of 448 hospitalizations, 188 patient files of people aged 60 years and over were collected, resulting in a prevalence of 42%. During the first wave, 66 cases out of 256 (25.8%) were recorded, 69 cases out of 102 (67.6%) during the second wave, and 53 cases out of 90 (59%) during the third wave (Fig. 1).




Fig. 1. Prevalence of patients aged 60+ hospitalized at Saint-Louis ETC during the first three waves of COVID-19

The mean age of the study population was 70 years (± 0.8) with a male predominance of 118 (62.8%) compared to 70 females (37.2%), resulting in a sex ratio of 1.7. Comorbidities were present in 115 patients (61.2%) and were mainly hypertension (77 cases, 67%), followed by diabetes mellitus (54 cases, 47%) and asthma (7 cases, 6%). Only 4 patients (2.1%) had received anti-COVID vaccines (Table 1).

Table 1. Distribution of patients according to epidemiological data

Epidemiological data	Number	Percentage (%)
Age		
60-65 years	65	34.6
66-70 years	49	26.1
71-75 years	34	18.1
76-80 years	17	9
81-85 years	14	7.4
86-90 years	6	3.2
91-95 years	3	1.6
Sex		
Males	118	62.8
Females	70	37.2
Marital status (n=145)		
Married	124	85.5
Widowed	16	11
Single	5	3.4
Comorbidities (115)		
HTN	77	67
Diabetes mellitus	54	47
Asthma	7	6
Cardiopathy	6	5.2
Dyslipidemia	6	5.2
Chronic Kidney Disease	3	2.6
Gout	3	2.6
Obesity	2	1.7
Rheumatoid Arthritis	1	0.9
Scleroderma	1	0.9
Sickle Cell Disease	1	0.9
Dementia	1	0.9
Anti-COVID Vaccination (n=188)		
Oui	4	2.1

Non	184	97.9
Type of anti-COVID vaccine(n=04)		
Sinopharm	2	50
AstraZeneca	1	25
Johnson & Johnson	1	25

3.2 Clinical Aspects

The mean time from symptom onset to hospitalization was 9.6 days (± 4.9). The mean SpO₂ was 88.6 % (± 10.2 %). The majority of patients had a severe form of the disease, with 121 cases (64.4%). Dyspnea was the most common symptom upon admission, with 111 cases (60.3%), followed by cough with 104 cases (56.5%), and ageusia with 45 cases (24.5%) (Table 2).

Table 2. Distribution of patients according to clinical data

Clinical data	Number	Percentage (%)
Severity (n=188)		
Mild form	36	19
Moderate form	26	13.8
Severe form	121	64.4
Critical form	5	2.7
Clinical symptoms		
Respiratory symptoms		
Dyspnea	111	60.3
Cough	104	56.5
Chest pain	29	15.2
Hemoptysis	1	0.5
Sneezing	1	0.5
ORL symptoms		
Ageusia	45	24.5
Anosmia	19	10.3
Rhinorrhea	18	9.8
Otalgia	1	0.5

Laryngeal crackles	1	0.5
Digestive symptoms		
Diarrhea	21	11.4
Vomiting	7	3.8
Odynophagia	6	3.3
Abdominal pain	4	2.2
Epigastric pain	3	1.6
Nausea	2	1.1
Hiccups	1	0.5
Constipation	1	0.5
Neurological symptoms		
Headache	44	24
Dizziness	3	1.6
Insomnia	1	0.5
Anxiety	1	0.5
Cardiovascular symptoms		
Precordialgia	1	0.5
General symptoms		
Fever	83	45.4
Asthenia	78	42.6
Arthralgia	59	32.2
Myalgia	54	29.5
Anorexia	54	29.5
Body aches	47	25.7
Lumbago	28	15.3
Chills	14	7.6
Sweat	10	5.5
Loss of weight	4	2.2
Lethargy	1	0.5

3.3 Paraclinical Aspects

The mean leukocyte count was 7898/uL (± 4093). Hyperleukocytosis ($>10,000/uL$) was found in 35 patients (18.6%). Lymphopenia ($<1500/uL$) was found in 108

patients (57.4%). The mean creatinine level was 14.5 mg/dL (± 22.2). Renal failure with a creatinine level ≥ 14 mg/L was found in 34 patients (19.5%). The mean fasting glycemia was 1.1 g/L (± 0.5). Hyperglycemia ≥ 1.26 g/L was found in 32 patients (23%). All patients (100%) underwent a diagnostic PCR test. A chest CT scan was performed in 51 patients (27.1%). The extent of the lesions was specified in 29 cases, with severe involvement predominating in 15 cases (51.7%) (Table 3).

Table 3. Distribution of patients according to paraclinical data

	Number	Percentage (%)
Hemogram (n=188)		
Hyperleukocytosis (>10000/ul)	35	18.6
Lymphopenia (<1500/ul)	108	57.4
Neutropenia (<1500/ul)	7	3.7
Thrombocytopenia (<150000/ul)	9	4.8
Anemia (≤ 10 g/dl)	13	7
Transaminases (n=125)		
ALT (>70UI)	15	12
AST (>60UI)	21	16.8
Creatinine (>14mg/l) (n=174)	34	19.5
CRP (≥ 12 mg/l) (n=38)	32	84.2
Hyperglycaemia ($\geq 1,26$ g/l) (n=139)	32	23
Hemoglobin A1C ($\geq 7\%$) (n=21)	8	38.1
Electrolyte test (n=78)		
Hyponatremia (<130meq/l)	6	7.7
Hypochloremia (<98meq/l)	17	21.8
Hypokalemia (<3,5meq/l)	18	23.1
Lipids (n=106)		
Hypercholesterolemia (>2g/l)	31	29.2
Hypo-HDL-cholesterolemia (<0,4g/l)	35	33
Hyper-LDL-cholesterolemia (>1,6g/l)	18	17

Hypertriglyceridemia (>1,6g/l)	28	26.4
Hypertroponinemia (>0,4ng/ml) (n=80)	80	100
Decrease in prothrombin levels<70% (n=155)	18	11.6
COVID PCR (n=188)	188	100
Thoracic CT scan lesion extent(n=29)		
0 à 25%	3	10.3
25 à 50%	10	34.5
50 à 75%	15	51.7
>75%	1	3.4

3.4 Therapeutic Aspects

The hydroxychloroquine-azithromycin combination was prescribed in 185 cases (98.4%). Antibiotic treatment was initiated in 126 cases (67%). Oxygen therapy was administered in 128 cases (68%) (Table 4).

Table 4. Distribution of patients according to therapeutic data

Therapeutic data	Number	Percentage (%)
Hydroxychloroquine-azithromycin combination	185	98.4
Antibiotherapy	126	67
Oxygen therapy	128	68.1
Non-invasive ventilation	5	2.6
Anticoagulation	179	95.2
Corticosteroid therapy	113	60.1
Analgesic/antipyretic	176	93.6
Vitamin C	181	96.3
Zinc	83	44.1

3.5 Evolutionary Aspects

The mean length of hospital stay was 12.6 days (± 4.7). Just over half of the patients, 100 out of 181 (55.2%), had a hospital stay between 11 to 15 days (Fig. 2). Regarding outcome, there were 153 recoveries (81.4%) and 35 deaths (18.6%). The

main factors associated with death were age (≤ 75 years) ($p=0.014$) [OR=0.488; 95% CI: 0.201-1.187]; critical forms ($p=0.017$) [OR=0.061; 95% CI:]; comorbidities ($p=0.031$) [OR=1.647; 95% CI: 0.280-2.499], complications ($p=0.009$) [OR=1.212; 95% CI: 0.530-2.774], and SpO₂ $\leq 95\%$ at admission ($p=0.031$) [OR=4.899; 95% CI: 0.752-8.797]




Fig. 2. Distribution of patients according to their length of hospital stay

4. DISCUSSION

In our series, the mean age was 70 years (± 8 years). Similar results were found by Wang et al. (China) and Touahri et al. (Algeria) who reported mean ages of 71 and 74 years, respectively [13,14]. In the preliminary study by Zulfiqar et al. (France) on a series of elderly COVID-19 patients, the mean age was slightly higher at 83.97 years [15]. The difference in mean age could be explained by the fact that life expectancy is higher in France (85.4 years for women and 79.3 years for men) [16], whereas in Senegal it is 67.7 years [17].

Comment [RS10]: All et al should be in BOLD

Our series had a male predominance (62.8%) with a sex ratio of 1.7. The same trend was found by Niu et al. (China) and Córdova et al. (Peru), with 56.7% and 51.9% male predominance respectively [18,19]. This male predominance could be explained by the fact that men are more mobile, often finding themselves in gatherings and therefore more exposed to COVID-19.

In our series, co-morbidities were present in 61.2% of patients, with high blood pressure (67%) and diabetes mellitus (47%) being the most common. This is consistent with the results of Wang et al. (China) who found a predominance of hypertension (40.8%) and diabetes mellitus (16%) [13]. Collarino et al. (France) also found a predominance of hypertension (67%) and diabetes mellitus (23%) [20]. This predominance may be explained by the high prevalence of these comorbidities in the general Senegalese population, with rates of 29.8% and 3.4%, respectively [17]. In particular, the Saint-Louis region of Senegal has a high prevalence of hypertension (46%) and diabetes mellitus (10.4%) [17].

In our series, the anti-COVID vaccination coverage rate was 2.1%. According to Masresha B et al, in the first year after vaccine introduction, only 6.8% of the total population was fully vaccinated in the 47 countries of the WHO African Region [21]. Due to the high lethality rate of COVID-19 in the elderly population, they quickly became a priority target for vaccination, despite the evolving indications based on availability and knowledge [22]. This low prevalence of vaccination coverage can be explained by people's reluctance to receive the anti-COVID vaccine.

In our series, most patients had severe COVID-19 (64.4%). The prevalence of severe forms in elderly individuals was corroborated by Mahiat C et al. (Belgium) and Chen et al. (China), who reported 72% and 87.3%, respectively [23,24]. The fact that elderly subjects often have many comorbidities and a weakened immune system may explain this. Indeed, they are more prone to develop severe forms of COVID-19 [11,25] due to senility and decreased IFN-1 production.

In our series, the hydroxychloroquine-azithromycin combination was prescribed in 98.4% of cases. This contrasts with the Lydie C study (France), where only 3% of patients received this combination [26]. The World Health Organisation's decision to discourage the use of drugs such as hydroxychloroquine in COVID-19 patients [27] may explain these results. However, because of its proven results in reducing the median length of hospitalization and the absence of serious side effects, this

Comment [RS11]: Replace In our series in paragraphs with In our subsequent study, Consecutively, Our work defines

molecule was included in the Senegalese treatment protocol at the beginning of the pandemic [28].

In terms of evolution, the mortality rate in our series was 18.6%. A mortality rate of 16.8% was reported by Tam EMY and al. (China) [29]. However, Kone D et al. (Ivory Coast) found a higher fatality rate of 23.7% [30]. This high fatality rate demonstrates the severity of COVID-19 in elderly individuals.

In our series, the main factors associated with death were age, critical forms, comorbidities, the occurrence of complications, and SpO₂ at admission. Many other factors associated with death have been reported in the literature. For instance, Aw et al. (England) found aging, male sex, and a high NEWS-2 score as factors of death [31]. Chen et al. found that factors associated with death were male sex, multiple comorbidities, dyspnea, high creatinine levels, and high AST levels [24]. Thiam et al. found that delirium was the only factor associated with death [32].

5. CONCLUSION

This study has shown that most patients aged 60 and over hospitalized in our CTE were hypertensive men, with very few having been vaccinated against COVID-19. Severe forms of the disease were predominant, and the mortality rate was high and associated with several factors. It is crucial to prioritize COVID-19 vaccination for this population to reduce mortality.

CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the authors.

REFERENCES

1. World Health Organization (2021) WHO Director-General's Opening Remarks at the Media Briefing on COVID-19-March 2020. <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>. Accessed on 06/29/2023.
2. Sohrabi C, Alsafi Z, O'Neill N, Khan M, Kerwan A, Al-Jabir A, et al. World Health Organization declares global emergency: A review of the 2019 novel coronavirus (COVID-19). *Int J Surg* 2020;76:71-6. <https://doi.org/10.1016/j.ijsu.2020.02.034>
3. Lo M, Sy A and Yade S. COVID-19 in Africa: Progress Report and Outlook /COVID-19 in Africa. https://www.policycenter.ma/sites/default/files/RP_20-13_Moubarack_Lo.pdf. Accessed on 06/27/2023.
4. Diouf I, Bousso A and Sonko I. COVID-19 pandemic management in Senegal. *Médecine de Catastrophe - Urgences Collectives* 2020; 4(3): 217-22. <https://doi.org/10.1016%2Fj.pxur.2020.08.009>
5. Omedia. Rapid survey on the effects of COVID-19. A Senegal 2020 gender perspective. https://data.unwomen.org/sites/default/files/documents/Publications/RAPPORT_RGA_UN_WOMEN_SENEGAL.pdf. Accessed on 11/15/2022.
6. Marih L, Sodqi M. Clinical Manifestations of COVID-19. *Batna J Med Sci* 2020;7: S9-S13. <http://dx.doi.org/10.48087/BJMSf.2020.S713>
7. Waechter C. Clinical and paraclinical features of COVID-19, virological diagnosis. *NPG Neurologie – Psychiatrie - Gériatrie* 2021; 21(125): 297-303. <https://doi.org/10.1016%2Fj.npg.2021.05.011>
8. Caumes E. Infection with SARS-CoV-2. *Arch Mal CoeurVaiss Prat* 2020; 2020(291): 2-4. <https://doi.org/10.1016%2Fj.amcp.2020.08.002>
9. Abdelmalek R. Covid-19, chronicles of a pandemic foretold. *Tunis Med* 2020 ; 98 (04): 278-281. <https://applications.emro.who.int/imemrf/455/Tunisie-Med-2020-98-4-295-298-fre.pdf>
10. Lapiere A, Fontaine G, Tremblay PL, Maheu-Cadotte MA and Desjardins M. Coronavirus disease (COVID-19): current state of knowledge. *Soins d'urgence* 2020; 1(1):13-8. https://www.researchgate.net/publication/341355504_La_maladie_a_coronavirus_COVID-19_portrait_des_connaissances_actuelles
11. Hamadouche M. Link between age and severe forms of Covid-19. Application to medical fitness decisions for healthcare personnel. *El Hakim Revue Médicale Algérienne* 2020; 20(30):84-

8. <https://www.researchgate.net/publication/347528153> Lien entre l'age et les formes seves res de Covid-19 Application a la decision d'aptitude medicale du personnel de sante
12. High Authority for Health report in 2020. Immunological and virological aspects of SARS-CoV-2 infection - Genetic variability, immune responses, vaccine platforms and animal models. https://www.has-sante.fr/upload/docs/application/pdf/2020-11/rapport_-_immunit%C3%A9_au_cours_de_l'infection_par_le_sars-cov-2_2020-11-30_17-25-10_860.pdf. Accessed on 11/10/2022.
13. Wang L, He W, Yu X, Hu D, Bao M, Liu H et al. Coronavirus disease 2019 in elderly patients: Characteristics and prognostic factors based on 4-week follow-up. *J Infect* 2020; 80(6): 639-45. <https://doi.org/10.1016/j.jinf.2020.03.019>
14. Touahri R and Ketfi A. Clinical, biological, radiological and evolutionary profile of COVID-19 according to age: a comparative study. *Rev Mal Respir Actual* 2021;13(1):114.
15. Zulfiqar AA, Villalba NL, Severac F, Andrès E. COVID-19 infection in a cohort of elderly patients in France: Results of a preliminary study. *La Presse Médicale Formation* 2020; 1(5): 460–463. <https://doi.org/10.1016%2Fj.lpmfor.2020.10.002>
16. Bagein G, Costemalle V, Deroyon T, Hazo J-B, Naouri D, Pesonel E, et al. The state of health of the population in France-2022. <https://drees.solidarites-sante.gouv.fr/sites/default/files/2022-09/DD102EMB.pdf>. Accessed on 11/01/2022
17. Senegalese Ministry of Health and Social Action. Non-Communicable Diseases Baseline Survey Final Report-2021. <https://openjicareport.jica.go.jp/pdf/12346235.pdf>. Accessed on 12/09/2022
18. Niu S, Tian S, Lou J, Kang X, Zhang L, Lian H et al. Clinical characteristics of older patients infected with COVID-19: A descriptive study. *Arch Gerontol Geriatr* 2020;89(104058):1-5. <https://doi.org/10.1016/j.archger.2020.104058>
19. Córdoba LDS, Vega APM, Luján-Carpio E, Parodi JF, Moncada-Mapelli E, Armacanqui-Valencia I et al. Clinical characteristics of older patients with COVID-19: a systematic review of case reports. *Dement Neuropsychol* 2021;15(1):1-15. <https://doi.org/10.1590/1980-57642021dn15-010001>
20. Collarino R, Vauloup-Fellous C, Allemang-Trivalle A, Mouna L, Duron E, Neiss M et al. Persistence and protection of neutralising antibodies 12 months after SARS-COV-2 infection in elderly subjects. *Infect Dis Now* 2021;51(5):S15. <https://doi.org/10.1016%2Fj.idnow.2021.06.005>

21. Masresha B, Ruiz MAS, Atuhebwe P and Mihigo R. The first year of COVID-19 vaccine roll-out in Africa: challenges and lessons learned. Pan African Medical Journal 2022;41(2):1-10. <https://doi.org/10.11604%2Fpamj.suppl.2022.41.2.33686>
22. Ndiaye M, Sow KD, Ly AB, Diop B, Ba M, Faye A. Epidemiology and response strategies against COVID-19: the Senegalese experience from 2020 to 2021. Pan Afr Med J 2022;43(204):1-13. <https://doi.org/10.11604%2Fpamj.2022.43.204.38290>
23. Mahiat C, Henrard S, Gilard I, Lanthier N, Stärkel P, Brauwer ID, et al. COVID-19: clinical presentation and mortality of the first 50 geriatric patients hospitalized at the Cliniques universitaires Saint-Luc in Spring 2020. Louv Med 2020;139 (09-10): 503-511. https://www.louvainmedical.be/sites/default/files/content/article/pdf/mahiatc_nov2020.pdf
24. Chen T, Dai Z, Mo P, Li X, Ma Z, Song S et al. Clinical characteristics and outcomes of older patients with coronavirus disease 2019 (COVID-19) in Wuhan, China (2019): a single-centered, retrospective study. J Gerontol A Biol Sci Med Sci 2020; 75(9):1788-95. <https://doi.org/10.1093/gerona/glaa089>
25. Bertholom C. Sars-CoV-2-associated immune response. Option/Bio 2021; 32(627): 15-17. [https://doi.org/10.1016%2FS0992-5945\(21\)00018-0](https://doi.org/10.1016%2FS0992-5945(21)00018-0)
26. Lydie C. SARS-Cov2 infection in elderly subjects in the Covid unit of the Cimiez geriatric hospital of the CHU de Nice: prognostic factors for mortality. Médecine humaine et pathologie. 2021. ffdumas-03710502f. <https://dumas.ccsd.cnrs.fr/dumas-03710502/document>
27. World Health Organization (WHO). Treatments against COVID-19: evolutionary orientations 2021. <https://iris.who.int/bitstream/handle/10665/347607/WHO-2019-%20nCoV-therapeutics-2021.3-fre.pdf?isAllowed=y&sequence=1>. Accessed on 12/09/2022
28. Journal of pharmacological information. Special COVID-19. https://www.atrsvv.dz/ressources/fichiers_produits/fichier_produit_1961.pdf. Accessed on 10/10/2022
29. Tam EMY, Kwan YK, Ng YY, Yam PW. Clinical course and mortality in older patients with COVID-19: a cluster-based study in Hong Kong. Hong Kong Med J;28(3): 215-222, 2022 06. <https://doi.org/10.12809/hkmj219367>
30. Kone D, Kadiane-Oussou J, Yapo MT, Akanji IA, Karidioula JM, Sule MA, et al. Comparative study of the Clinical manifestations of COVID-19 in adults versus in

Aged subjects at the Bouake CHU (Ivory Coast).Rev Mali Infect Microbiol 2022 ;17(2) :26-38.

31. Aw D, Woodrow L, Ogliari G and Harwood R. Association of frailty with mortality in older inpatients with Covid-19: a cohort study. Age Ageing 2020;49(6):9. <https://doi.org/10.1093/ageing/afaa184>

32. Thiam CN, Hasmukharay K, Lim WC, Ng CC, Pang GHM, Abdullah A, et al. COVID-19 Infection among Older People Admitted to Hospital: A CrossSectional Analysis. Geriatrics (Basel) 2021;6(25):2-8. <https://doi.org/10.3390/geriatrics6010025>

UNDER PEER REVIEW