

EFFECT OF EXOGENOUS PLANT GROWTH REGULATORS ON GROWTH, YIELD, AND QUALITY OF PARTHENO-CARPIC CUCUMBER/ENGLISH CUCUMBER (*Cucumis sativus* L.) UNDER PROTECTED CONDITION

ABSTRACT

This study was carried out during the Rabi season 2022-2023 to investigate the effect of different levels of NAA, GA3, and TIBA on the growth, yield, and quality of Pusa Parthenocarpic cucumber – 6 and to estimate the economics. Conducted from January to March 2023, the research reveals NAA's significant effect on vine length, earliness parameters, yield, and quality attributes. Treatment 8 (NAA@200ppm) emerges as particularly promising, demonstrating superior growth with an average plant height of 290 cm, accelerated development with an average fruit weight of 252.67 g, and a maximum benefit-cost ratio of 2.39.

Keywords: Parthenocarpic Cucumber, Pusa Variety 6, Exogenous PGR, BC Ratio

1. INTRODUCTION

Cucumber (*Cucumis sativus* L.) is a widely cultivated plant, belonging to the family Cucurbitaceae and having a chromosome number, $2n = 14$ [1]. It is an endemic vegetable in India. It is a fruit that is commonly used as a vegetable in cuisines all over the world. Cucumbers are often eaten raw in salads or pickled in vinegar to make a tasty snack. The cucumber has a crunchy texture and a refreshing taste, making it a great addition to any meal. There are several different varieties of cucumbers available, including English cucumbers, Persian cucumbers, and pickling cucumbers. English cucumbers are longer and thinner than other varieties, while Persian cucumbers are shorter and have a sweeter flavor. Pickling cucumbers are smaller and are ideal for making pickles. Cucumbers are a low-calorie food that is high in vitamins and minerals. They are a great source of vitamin C, vitamin K, and potassium. They are also rich in antioxidants, which can help to protect the body against damage from free radicals. In addition to being a healthy food, cucumbers also have some cosmetic benefits. They are often used in beauty treatments to help soothe and hydrate the skin. Cucumber slices can be placed over the eyes to reduce puffiness and dark circles.

Parthenocarpic Cucumber is gaining importance in Urban and Peri-Urban Farming mostly due to its taste, colour and nature. The promising sweet varieties have gained an overwhelming response from the

market. Parthenocarpy refers to the ability of plants to produce fruit without fertilisation, resulting in seedless fruits. This characteristic is highly desirable in cucumber cultivation as it enhances fruit quality and market value. Parthenocarpic cucumbers are cucumber varieties that can produce fruit without the need for pollination. These varieties are sometimes referred to as self-pollinating, but technically, they do not require pollination to set fruit. The development of cucumbers without fertilization occurs due to genetic factors or the application of plant hormones.

Parthenocarpy in cucumber cultivation offers several advantages. First, it ensures a higher rate of fruit set, as it bypasses the need for pollination and eliminates the dependence on pollinators. This is particularly beneficial in conditions where pollinators are scarce or inefficient. Second, parthenocarpic cucumbers can be harvested earlier compared to pollinated ones. This early harvest allows for an extended harvesting period and increased overall yield. Another benefit of parthenocarpy is the production of seedless cucumbers. These seedless varieties are preferred by consumers as they have a more pleasant texture and taste. [2] Additionally, the absence of seeds eliminates the need for seed removal, making them more convenient. 95% of a raw cucumber (with peel) is water, 4% is carbohydrate, 1% is protein, and very little fat is present. A reference serving of 100 grammes (3+1/2-ounces) has 65 kilojoules (16 kilocalories) of food energy. Its micronutrient content is minimal; only vitamin K, with 16% of the Daily Value, stands out.[3]

Parthenocarpic cucumbers also tend to exhibit greater resilience to environmental factors such as heat, drought, and low light. Their ability to develop fruit without depending on pollination makes them more resistant to adverse conditions, making them suitable for cultivation in challenging climates. However, the occurrence of parthenocarpy is often inconsistent, and external factors such as plant growth regulators can influence its expression.

NAA (naphthaleneacetic acid), GA (gibberellic acid), and TIBA (2,3,5-triiodobenzoic acid) are well-known plant growth regulators commonly used in horticulture and agriculture. NAA is a synthetic auxin that promotes cell elongation and fruit growth. GA stimulates plant growth, including internode elongation and seed germination. Conversely, TIBA is a synthetic auxin transport inhibitor that affects various plant growth processes [4].

By applying different concentrations and combinations of NAA, GA, and TIBA, the study will examine their individual effects on plant growth parameters such as plant height, leaf area, and stem diameter, as well as fruit quality attributes like fruit weight, size, shape, colour, and nutrient content. The findings of this study will contribute to the understanding of how plant growth regulators can influence the growth, yield, and quality of parthenocarpic cucumber plants. This knowledge can help optimize cucumber cultivation techniques, improve crop productivity, and enhance the economic viability of cucumber production under protected conditions.

2. MATERIALS AND METHODS

This experiment was carried out from January to March 2023 in the Horticulture Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj (U.P). Ten treatments were used, including a control (Table 1) and three replications in a randomised block design with a spacing of 60×30 cm were applied at 2 and 4 true leaf stages under protected conditions, Gynoeciousparthenocarpic cucumber (*Cucumis sativus* L.) var. Pusa Parthenocarpic Cucumber-6 was the subject of the experiment. Each treatment group consisted of Five randomly chosen plants for the observational recording of metrics related to growth, earliness, yield, quality, economic and quality parameters viz. The information gathered throughout the inquiry was

exposed to Fisher's description of statistical analysis of variance (ANOVA) (1950).

Table 1. Various levels of TIBA, GA3 and NAA

| Sr. No. | Treatment | Treatment Combination |
|---------|-----------|-----------------------|
| 1 | T0 | Water |
| 2 | T1 | TIBA@50ppm |
| 3 | T2 | TIBA@100ppm |
| 4 | T3 | TIBA@150ppm |
| 5 | T4 | GA@500ppm |
| 6 | T5 | GA@1000ppm |
| 7 | T6 | GA@1500ppm |
| 8 | T7 | NAA@100ppm |
| 9 | T8 | NAA@200ppm |
| 10 | T9 | NAA@300ppm |

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

3.1.1 Plant Height

Table 2 presents notable variations in vine length, with the most substantial measurement observed 45 days after sowing. Specifically, T8 (NAA@200ppm) exhibited the longest vine length, reaching 292.00 cm, followed by T7 (NAA@100ppm) with a measurement of 130.51 cm. In contrast, the control group (T0) recorded a shorter vine length of 171.00 cm according to the data in Table 2. Naphthalene Acetic Acid (NAA) plays a pivotal role in the physiological processes occurring throughout the growth period. It enhances cell division and elongation while also elevating the permeability of the cell wall. This heightened permeability facilitates the entry of a greater quantity of water and dissolved materials into the cell, ultimately contributing to increased cell size and overall growth (Devlin, 1975) [5].

3.1.2 Number of Primary Branches

Table 2 presents notable variations in the mean of the number of branches, with the most substantial measurement observed 45 days after sowing. Specifically, T8 (NAA@200ppm) exhibited the highest number reaching 4.60, followed by T9 (NAA@100ppm) with a number of 3.80. In contrast, the control group (T0) has a minimum number of 3.40 according to the data in Table 2. There exists a direct correlation between the main vine and the formation of primary branches in plants. This means that an increase in the length of the main vine leads to a corresponding increase in the number of primary branches, and vice versa. The observed significance in results could be attributed to the reversible impact on plant mitosis caused by the application of NAA. This, in turn, results in a greater number of primary

branches on the main branches of cucumber. Hence, a longer vine length contributes to a higher count of primary branches. Similar findings were also reported by Dalai et al. 2016 [6].

3.1.3 Number of Nodes

At the 45-day mark post-transplantation, the highest node count was documented in T8 (NAA@200ppm), reaching 11.27, followed by T9 (NAA@300ppm) reaching 11.20. In contrast, the minimum number of nodes was observed in T0 (control), registering at 9.73. The significant rise in the node count could be attributed to the potential stimulation of meristematic tissues by auxin, hastening increased cell division and enlargement in actively growing sections. These observations align with the research outcomes reported by Viradia et al., [7] and El-Soad et al., [8].

3.2 Earliness Parameters

3.2.1 Days to First Flowering

The data regarding the days taken for the first appearance of female flowers reveals a statistically significant outcome. The shortest duration for the emergence of the first female flowering was observed in T8 (NAA@200ppm) at 33.80 days (mean), followed by T1 (TIBA@50ppm) at 34.60 days (mean), while the longest duration was noted in T0 (control) at 43.34 days (mean), as presented in Table 2. Cucumber flowers are typically characterized by their small size and yellow colour, featuring 3-6 petals per blossom. The development of cucumber flowers is significantly influenced by auxin. A detailed study has been reported by Sharif et al. 2022 [9].

3.2.2 Days to First Fruit Setting

The analysis of data revealed that the maximum number of days taken for the first fruit setting was observed in the control group (T0) at 49.62 days (mean), whereas the minimum duration was recorded in treatment T8 (NAA@200ppm) at 40.08 days (mean). Following closely was the treatment T1 (TIBA@50ppm) with a time frame of 40.88 days (mean) for the first fruit setting. Auxin has been widely utilised to achieve successful fruit setting without fertilization. In a study conducted by Qian et al. [10], the advantages of externally applying auxin for the induction of parthenocarpic fruit in cucumbers were demonstrated. Specifically, Naphthaleneacetic acid (NAA) at a concentration of 100 mg/L was sprayed on unpollinated cucumber ovaries. The cucumbers treated with NAA exhibited successful fruit setting, in contrast to untreated ovaries, which did not undergo successful fruit development.

3.2.3 Days to First Fruit Picking

The analysis of data revealed that the maximum number of days taken for the first fruit setting was observed in the control group (T0) at 58.03 days (mean), whereas the minimum duration was recorded in treatment T8 (NAA@200ppm) at 48.5 days (mean). Following closely was the treatment T1 (TIBA@50ppm) with a time frame of 49.3 days (mean) for the first fruit picking. As Auxin is found in higher concentration in the parthenocarpic line of cucumber the NAA treatment reduced the fruit abortion and increased the photosynthetic effect to induce the carbohydrates accumulation in fruit to be ready for harvest earlier. The same case study has been reported by Sharif et al. 2022 [9].

3.3. Yield Parameters

3.3.1 No. of Female Flowers per Plant

The data analysis indicated that the highest number of female flowers per plant was observed in treatment T8 (NAA@200ppm), reaching 16.41. Following closely was T5 (GA@1000ppm) with a count of 16.24. In contrast, the minimum number of female flowers per plant was found in treatment T0 (control) at 13.60. This phenomenon may be attributed to the elevated levels of exogenous auxin, specifically Naphthaleneacetic acid (NAA). The higher concentration of NAA is likely to initiate and sustain additional floral meristems, consequently leading to an augmented number of female flowers. Additionally, auxin plays a crucial role in the growth and differentiation of carpels, fostering the development of more female flowers. A similar study has been reported by Sharif et al. 2022 [9].

3.3.2 Avg. fruit weight (g)

The data analysis indicated that the maximum average fruit weight per plant was observed in treatment T8 (NAA@200ppm), reaching 252.67. Following closely was T9 (NAA@300ppm) with 230.67. In contrast, the minimum number of female flowers per plant was found in treatment T0 (control) at 130.67. Naphthaleneacetic acid (NAA) is a synthetic auxin with a physiological function similar to that of Indole-3-acetic acid (IAA). Due to its lower cost and increased stability, NAA is deemed more suitable for application in crop production. Following the fruit set, the majority of assimilates produced through photosynthesis are allocated to fruit growth, with a portion reserved for essential physiological activities such as growth and respiration. Auxins play a role in enhancing the photosynthetic capacity of leaves and regulating the loading, transport, and distribution of photoassimilates. This, in turn, improves the seed setting rate and contributes to an increase in crop weight. Parallel findings were reported by Chen et. al (2022) in Giant Pumpkin [11].

3.3.3 No. of fruits per plant

The analysis of the data revealed that the maximum number of fruits per plant was recorded in treatment T8 (NAA@200ppm), reaching 9.87. Following closely was T7 (NAA@100ppm) with a 9.27 count of fruits per plant. In contrast, the minimum number of fruits per plant was found in treatment T0 (control) at 8.27. NAA plays a regulatory role in increasing the number of female flowers and decreasing the male-to-female sex ratio. The presence of female flowers at lower nodes on the main vine contributes to a potential reduction in fruit abortion. Consequently, this leads to a proportional increase in the number of fruits per plant. Additionally, the fresh weight per fruit tends to be at its highest under these conditions. A similar study was found by Akter, P., & Rahman, M. A. (2010) in Bitter Gourd [12].

3.3.4 Fruit yield per plot (kg)

The analysis of the data revealed that the maximum number of fruits per plot was recorded in treatment T8 (NAA@200ppm), reaching 10.93 kg. Following closely was T6 (GA@1500ppm) with 9.55 kg of fruits per plant. In contrast, the minimum number of fruits per plant was found in treatment T0 (control) at 6.84. NAA, along with various other plant growth regulators, are recognised for inducing physiological changes in plants, primarily impacting flowering behaviour, sex ratio, increased fruit set, and the enlargement and development of fruits. Additionally, they play a role in influencing the source-sink relationship in plants. The application of growth regulators brings about specific alterations in metabolic processes during fruit and seed development. This, in turn, leads to a greater accumulation of food reserves, ultimately contributing to a higher yield. A similar study was found by Akter, P., & Rahman, M. A. (2010) in Bitter Gourd [12].

3.3.4 Fruit length (cm)

The analysis of the data revealed that the maximum fruit length was recorded in treatment T8 (NAA@200ppm) at 17.68, closely followed by T1 (TIBA@50ppm) with a length of 17.74. In contrast, the minimum fruit length was found in treatment T0 (control) at 15.15. Fruit length is a significant agronomic trait in cucumber breeding. The metabolic effect of NAA associated with this trait results in the acceleration of cell enlargement and the inhibition of cell division. As a consequence, this metabolic influence contributes to the attainment of the highest possible fruit length. A similar study was reported by Dahal et al. (2022) in chilli [13].

3.3.5 Fruit diameter (cm)

The data analysis indicated that the maximum fruit diameter was recorded in treatment T8 (NAA@200ppm) at 4.56, followed by T7 (NAA@100ppm) with a diameter of 4.36. In contrast, the minimum fruit diameter was observed in treatment T0 (control) at 3.59. Similar results were observed by ANBARASI et al (2022) [14]. The observed increase in diameter can be attributed to both, an increase in the number of cells and the elongation of cells, which are characteristic actions of any group of auxins.

Table 2. Effect of PGRs on growth, earliness, and yield parameters of Parthenocarpic Cucumber

| Treatment | Plant Height (cm) | No of Primary Branches | No of Nodes | Days to first flowering | Days to fruit setting | Days to first fruit picking | No. of female flowers per plant | Avg. fruit weight (g) | No. of fruits per plant | Fruit yield per plot (kg) | Fruit length (cm) | Fruit diameter (cm) |
|-----------|-------------------|------------------------|-------------|-------------------------|-----------------------|-----------------------------|---------------------------------|-----------------------|-------------------------|---------------------------|-------------------|---------------------|
| T0 | 171 | 3.40 | 9.73 | 43.34 | 49.62 | 58.03 | 13.60 | 130.67 | 8.20 | 6.84 | 15.15 | 3.59 |
| T1 | 193.33 | 3.80 | 10.33 | 34.60 | 40.88 | 49.30 | 13.69 | 149.67 | 8.33 | 7.76 | 17.74 | 4.06 |
| T2 | 218.33 | 3.73 | 9.93 | 42.67 | 48.95 | 57.36 | 14.92 | 187.67 | 9.00 | 9.50 | 17.24 | 3.95 |
| T3 | 224.33 | 3.80 | 12.07 | 38.34 | 44.62 | 53.04 | 15.25 | 216.33 | 9.20 | 8.48 | 16.93 | 4.36 |
| T4 | 173.33 | 3.73 | 10.13 | 40.08 | 46.36 | 54.78 | 13.77 | 157.00 | 8.27 | 8.10 | 16.18 | 4.01 |
| T5 | 213.67 | 3.60 | 10.80 | 39.98 | 46.26 | 54.68 | 16.24 | 177.00 | 9.80 | 9.29 | 16.57 | 3.79 |
| T6 | 195.00 | 3.60 | 10.93 | 35.08 | 41.36 | 49.78 | 14.26 | 175.33 | 8.60 | 9.55 | 16.91 | 3.83 |
| T7 | 253.33 | 3.67 | 10.40 | 38.19 | 44.47 | 52.88 | 15.42 | 177.00 | 9.27 | 8.45 | 16.79 | 4.36 |
| T8 | 292.00 | 4.60 | 11.27 | 33.80 | 40.08 | 48.50 | 16.41 | 252.67 | 9.87 | 10.93 | 17.68 | 4.56 |
| T9 | 240.00 | 3.80 | 11.20 | 37.67 | 43.95 | 52.37 | 14.11 | 230.67 | 8.60 | 9.07 | 16.42 | 3.42 |
| F Test | S | S | S | S | S | S | S | S | S | S | S | S |
| SE(d)± | 14.39 | 0.35 | 0.59 | 1.45 | 1.45 | 1.54 | 0.42 | 16.45 | 0.21 | 0.92 | 0.69 | 0.42 |
| CD 5% | 3.02 | 0.73 | 1.24 | 3.04 | 3.05 | 3.58 | 0.89 | 3.45 | 0.45 | 1.93 | 1.45 | 0.89 |
| CV | 8.10 | 11.25 | 6.76 | 4.62 | 3.97 | 3.34 | 3.51 | 10.87 | 2.94 | 12.82 | 5.04 | 12.94 |

Table 3. Effect of PGRs on Qualitative Parameters of Parthenocarpic Cucumber

| Treatment | Chlorophyll Content (SPAD Value) | TSS (°Brix) | Vitamin C (mg/100gm) | Juice % | Dry Matter Content |
|-----------|----------------------------------|-------------|----------------------|---------|--------------------|
| T0 | 22.90 | 4.52 | 4.54 | 75.97 | 4.31 |
| T1 | 33.46 | 4.73 | 4.71 | 78.21 | 5.57 |
| T2 | 33.57 | 5.23 | 4.65 | 82.79 | 6.18 |
| T3 | 30.87 | 5.91 | 5.64 | 89.30 | 5.44 |
| T4 | 26.07 | 5.47 | 5.69 | 89.08 | 5.81 |
| T5 | 29.70 | 5.87 | 5.16 | 80.22 | 5.88 |
| T6 | 26.07 | 5.33 | 5.30 | 82.39 | 5.54 |
| T7 | 28.87 | 5.33 | 5.08 | 84.16 | 5.26 |
| T8 | 36.83 | 6.03 | 5.70 | 91.50 | 6.83 |
| T9 | 25.73 | 5.28 | 5.16 | 84.28 | 5.11 |
| F Test | S | S | S | S | S |
| SE(d)± | 0.31 | 0.38 | 0.31 | 3.10 | 0.71 |
| CD 5% | 0.65 | 0.81 | 0.65 | 6.50 | 1.50 |
| CV | 1.29 | 8.76 | 7.31 | 4.52 | 15.63 |

3.4 Quality Parameters

3.4.1 Chlorophyll Content (SPAD Value)

While taking the readings for Chlorophyll content per leaf using a SPAD meter, the following observations were recorded. After 15 days, the maximum reading was recorded in treatment T8 (NAA@200ppm) at 40.90, followed by T3 (TIBA@150ppm) at 38.63. The minimum recording was recorded in treatment T0 (control) at 34.77. After 30 days, the maximum reading was recorded in treatment T8 (NAA@200ppm) at 38.87, followed by T1 (TIBA@50ppm) at 35.78. The minimum recording was recorded in treatment T0 (control) at 29.15. After 45 days, the maximum reading was recorded in treatment T8 (NAA@200ppm) at 36.83, followed by T2 (TIBA@100ppm) at 33.57. The minimum recording was recorded in treatment T0 (control) at 22.90. Leaves play a crucial role in various plant activities, including photosynthesis, respiration, and transpiration. The exogenous application of Naphthaleneacetic acid (NAA) effectively enhances leaf photosynthetic performance, potentially through the increase in leaf area and chlorophyll content. This improvement may also be associated with enhancements in both stomatal and non-stomatal restrictions, along with the reinforcement of photosynthetic system II, as compared to other treatments. Parallel findings were reported by Chen et. al (2022) in Giant Pumpkin [11].

3.4.2 TSS (°Brix), Vitamin C, Juice Percentage and Dry Matter Content

Various readings were taken while determining the qualitative parameters. TSS (°Brix) was maximum in treatment T8 (NAA@200ppm) at 6.03, followed closely by treatment T3 (TIBA@150ppm) at 5.91, and the lowest in treatment T0 (control) at 4.52.

Vitamin C content (mg/100gm) was recorded highest in the treatment T8 (NAA@200ppm) at 5.70, followed very closely by the treatment T4 (GA@500ppm) at 5.69, and the lowest in the treatment T0 (control) at 4.54. Ascorbic acid, being a highly unstable vitamin, is susceptible to changes in its content due to exposure to heat, oxygen, and light. Additionally, the manner of washing, whether in still or turbulent water, can induce mechanical stress, leading to a reduction in the nutrient concentration (Dewhirst et al., 2017) [15]. However, the study was done under a protected environment, hence the exposure to external environmental factors was very minimised due to which the Vitamin C content might be unaffected.

Juice percentage was recorded highest in treatment T8 (NAA@200ppm) at 91.50%, followed very closely by treatment T3 (TIBA@150ppm) at 89.30%, and the lowest in the treatment T0 (control) at 75.97%.

Dry matter content (in grams) was recorded maximum in the treatment T8 (NAA@200ppm) at 6.83 grams, followed closely by the treatment T (TIBA@100ppm) at 6.18 grams, and the lowest in the treatment T0 (control) at 4.31 grams.

Auxins play an important role in the metabolic rates and increase the photosynthetic activities in the plants which led to the higher nutrient contents, so these parameters might have been affected by it. Similar results were found by Vidyullatha et al., 2022 [16] and Sravika et al., 2021 [17].

3.5 Economics of various treatments

Table 4. Cost of cultivation of Parthenocarpic Cucumber under protected condition

| Sr. No. | Particulars | Unit | Rate/Unit | Quantity | Cost (Rs/ha) |
|---------|--|----------|-----------|----------|-----------------|
| 1 | Land Preparations | Man-days | 500 | 6 | 3,000 |
| 2 | Mulching | KG | 150 | 100 | 15,000 |
| 3 | Vermicompost | Ton | 2000 | 10 | 20,000 |
| 4 | Seeds | KG | 2 | 10,000 | 20,000 |
| 5 | Sowing | Man-days | 500 | 7 | 3,500 |
| 6 | Irrigation (Electricity) | KWH | 3 | 1,200 | 3,600 |
| 7 | Training Wire (steel) | KG | 120 | 20 | 2,400 |
| 8 | Training Wire (nylon) | KG | 180 | 5 | 900 |
| 9 | Harvesting | Man-days | 500 | 25 | 12,500 |
| 10 | Land (inc. poly house and irrigation system) | Months | 10,000 | 2 | 20,000 |
| 11 | Misc. Labour | Man-days | 500 | 4 | 2,000 |
| | Total | | | | 1,02,900 |

Table 5. Economies of different treatments done on Parthenocarpic Cucumber for this study

| Treatments | Fruit Yield (t/ha) | Cultivation Cost (INR) | Gross Revenue (INR) | Net Returns (INR) | Benefit:Cost Ratio |
|------------|--------------------|------------------------|---------------------|-------------------|--------------------|
| T0 | 3.42 | 1,02,900 | 1,53,900 | 51,000 | 1.50 |
| T1 | 3.88 | 1,02,900 | 1,74,600 | 71,700 | 1.70 |
| T2 | 4.75 | 1,02,900 | 2,13,750 | 1,10,850 | 2.08 |
| T3 | 4.24 | 1,02,900 | 1,90,800 | 87,900 | 1.85 |
| T4 | 4.05 | 1,02,900 | 1,82,250 | 79,350 | 1.77 |
| T5 | 4.64 | 1,02,900 | 2,08,800 | 1,05,900 | 2.03 |
| T6 | 4.77 | 1,02,900 | 2,14,650 | 1,11,750 | 2.09 |
| T7 | 4.23 | 1,02,900 | 1,90,350 | 87,450 | 1.85 |
| T8 | 5.47 | 1,02,900 | 2,46,150 | 1,43,250 | 2.39 |

| | | | | | |
|----|------|----------|----------|----------|------|
| T9 | 4.54 | 1,02,900 | 2,04,300 | 1,01,400 | 1.99 |
|----|------|----------|----------|----------|------|

The highest gross returns were observed in Treatment 8 (NAA@200ppm) at Rs. 2,46,150, followed by T6 (GA@1500ppm) at Rs. 2,14,650. Conversely, the lowest returns, amounting to Rs. 1,53,900, were noted in T0 (control).

In terms of net returns, Treatment 8 (NAA@200ppm) demonstrated the highest figure at Rs. 1,43,250, followed by T6 (GA@1500ppm) at Rs. 1,11,750. The lowest net returns, totalling Rs. 51,000, were registered in T0 (control).

Regarding the Benefit-Cost (BC) ratio, Treatment 8 (NAA@200ppm) exhibited the highest ratio at 2.39, followed by T6 (GA@1500ppm) at 2.09. The lowest BC ratio, recorded at 1.50, was associated with T0 (control). All these data are shown in the table 5 above.

4. CONCLUSION

This study concludes that NAA (200 ppm) performed best in terms of growth (vine length at 292 cm, chlorophyll content at 36.83), yield (5.47 t/ha), and quality (TSS value at 6.03 °Brix, dry matter content at 6.83). The highest BC ratio was at 2.39 in the treatment T8 (NAA @ 200 ppm).

REFERENCES:

1. Kaur, M., & Sharma, P. (2022). Recent advances in cucumber (*Cucumis sativus* L.). *The Journal of Horticultural Science and Biotechnology*, 97(1), 3-23.
2. Homestead Crowd. "Parthenocarpic Cucumbers (13 Best Varieties to Grow Now)". Available: <https://homesteadcrowd.com/parthenocarpic-cucumbers-best-varieties/>. Accessed on December 11, 2023.
3. by HEALTH JADE TEAM "Cucumber nutrition". Available: <https://healthjade.com/cucumber/> Accessed on December 13, 2023
4. Kumar, R., Dongariyal, A., Bhatt, R., & Raturi, P. (2020). Chapter-2 Plant Growth Regulators. Chief Editor Manoj Kumar Ahirwar, 49, 23.
5. Devlin, R. M. (1975). *Plant Physiology* 3rd. Edition D. Van Nostrand Comp. New York, P. 600.
6. Dalai, S., Singh, M. K., Kumar, M., Singh, K. V., & Kumar, V. (2016). Growth, flowering and yield of cucumber (*Cucumis sativus* L.) as influenced by different levels of NAA and GA3. *Journal of Plant Development Sciences*, 8(9), 445-450.
7. Viradia RR. Effect of growth regulators on growth, yield and quality of tomato cv. Pusa ruby. M. Sc. (Ag.) Thesis, Gujarat Agricultural University, S.K. Nagar, GUJARAT (INDIA);1982.
8. El-Soad IAA, Omran AF, Ashour NI. Stimulatory effects of 2,4-D on growth and yield of tomato. *Egyptian J. Hort.* 1976;3(2):149-155.
9. Sharif, R., Su, L., Chen, X., & Qi, X. (2022). Involvement of auxin in growth and stress response of cucumber. *Vegetable Research*, 2(1), 3-4.
10. Qian C, Ren N, Wang J, Xu Q, Chen X, et al. 2018. Effects of exogenous application of CPPU, NAA and GA4+7 on parthenocarpy and fruit quality in cucumber (*Cucumis sativus* L.). *Food Chemistry* 243:410–13.

11. Chen, C., Wu, X. M., Pan, L., Yang, Y. T., Dai, H. B., Hua, B., ... & Zhang, Z. P. (2022). Effects of exogenous α -naphthaleneacetic acid and 24-epibrassinolide on fruit size and assimilate metabolism-related sugars and enzyme activities in giant pumpkin. *International Journal of Molecular Sciences*, 23(21), 13157.
12. Akter, P., & Rahman, M. A. (2010). Effect of foliar application of IAA And GA on sex expression, yield attributes and yield of bitter gourd (*Momordica Charantia* L.). *Chittagong University Journal of Biological Sciences*, 55-62.
13. Dahal, M., Kumar, J., Silas, V. J., Devkota, S., & Dahal, A. (2022). Effect of naphthaleic acetic acid (NAA) on growth and yield attributing characteristics of chilli (*Capsicum annum* L.), Pusa Jwala
14. ANBARASI, D., & VENKATRAMAN, M. (2022). Effect of plant growth regulators on growth, yield and quality characters of chilli (*Capsicum annum* L.). *Annals of Plant and Soil Research*, 24(4), 543-546.
15. Dewhirst, R. A., Clarkson, G. J., Rothwell, S. D., & Fry, S. C. (2017). Novel insights into ascorbate retention and degradation during the washing and post-harvest storage of spinach and other salad leaves. *Food Chemistry*, 233(15), 237-246. <http://dx.doi.org/10.1016/j.foodchem.2017.04.082>. PMID:28530571.
16. Vidyullatha, L. M. M., & Topno, S. E. (2022). Effect of Naphthalene Acetic Acid & Indole Acetic Acid on Growth, Yield and Quality of Muskmelon (*Cucumis melo* L.). *International Journal of Plant & Soil Science*, 34(22), 1460-1469.
17. Sravika, B., Patro, K. K. K., Kumari, K. U., Emmanuel, N., & Suneetha, S. (2021). Effect of growth regulators on yield and quality parameters of summer squash (*Cucurbita pepo* L.). *The Pharma Innovation Journal*, 10(8), 708-711.