

Original Research Article

Exploring the In-Vitro Antibacterial Efficacy of Aqueous Extracts from Leaves, Bark, and Stem of *Celtis timorensis* Span: A Comprehensive Study

ABSTRACT

The current study evaluated the in-vitro antibacterial activity of *Celtis timorensis* Span. (*C. timorensis*) against *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853), and *Staphylococcus aureus* (ATCC 25923) in response to the global healthcare challenge of infectious diseases caused by Multidrug Resistance organisms. The study was motivated by the inappropriate use of existing antimicrobials and the insufficient discovery of new agents contributing to this crisis. Drawing inspiration from historical evidence in plant-based Ayurveda and traditional medicine, the study focused on *C. timorensis*, a plant known for its historical use in treating infectious diseases.

Crude extracts from the leaves, bark, and stem of *C. timorensis* were prepared using the cold maceration process, and their antibacterial activity was assessed against the aforementioned bacterial strains. The cylinder plate method was employed to measure zones of inhibition at various concentrations (250 µg/ml, 500 µg/ml, 750 µg/ml, and 1000 µg/ml). Positive and negative controls included Gentamicin and distilled water respectively. The diameter of inhibition zones was measured after 24 hours of incubation.

Results indicated positive antibacterial effects of all three extracts against *E. coli* (ATCC 25922) and *P. aeruginosa* (ATCC 27853). Notably, aqueous extraction from the stem exhibited the highest inhibitory zones against *E. coli* and *P. aeruginosa*. However, none of the concentrations of the extracts showed positive antibacterial effects against *S. aureus* (ATCC 25923). Statistical analysis confirmed the significance of the observed antibacterial activity against *E. coli* and *P. aeruginosa*.

Dose-response study results highlighted the efficacy and potency of aqueous extractions from bark and leaves against *Escherichia coli*. In contrast, leaves and bark demonstrated the highest efficacy and potency, respectively against *Pseudomonas aeruginosa*. In conclusion, the study scientifically validated the hypotheses formulated at the study's outset, utilizing evidence from plant-based Ayurveda and traditional medicine. The findings underscore the potential of *C. timorensis* as a source of antibacterial agents in the context of addressing multi-drug resistance-related infectious diseases.

Keywords: *Celtis timorensis*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, antibacterial activity

1. INTRODUCTION

Microbial diseases, constituting a substantial percentage exceeding 20% of global mortality, persist as a formidable challenge despite the advancements in antimicrobial chemotherapy (Saga, T. and Yamaguchi, K., 2009). The World Health Organization (WHO) actively addresses this challenge through initiatives like the Global Antibiotic Research and Development Partnership (GARDP), which advocates for innovative antibiotic treatments and alternative approaches. Despite the routine application of plant-derived remedies by Ayurvedic practitioners for therapeutic interventions, there exists a notable underutilization of numerous plant-derived antimicrobial agents within Western medical practices. This discrepancy is ascribed, in part, to the prevailing inclination toward using plant-derived medicine, attributed to their perceived diminished side effects compared to their synthetic counterparts. Consequently, the scope of research findings on plant-derived antimicrobial agents has experienced a notable surge. These agents exhibit both bactericidal and bacteriostatic effects, with the diverse array of secondary metabolites in medicinal plants serving as viable alternatives to resistance-modifying agents, thereby impeding the adaptive response of bacteria, viruses, and fungi (Guptha and Birdi, 2017).

The present study focuses on *Celtis timorensis*, colloquially known as "Gurenda" (www.kew.org, 2023) a flowering plant extensively distributed in humid regions encompassing India, Sri Lanka, Burma, and the Malay Islands (<https://indiabiodiversity.org>, 2023). *C. timorensis* belongs to the Cannabaceae family and is recognized for its medicinal properties, notably in wound healing (Rajaneekar Dasari, Kunchibhotla, Sathyavathi, & Reddy, 2013). Despite extensive global studies on its therapeutic attributes, limited information is available concerning its antimicrobial effects. *C. timorensis* comprises various phytoconstituents, including alkaloids, carbohydrates, proteins, sterols, phenols, flavonoids, gums, mucilage, and saponins. Sinhala Ayurveda literature attests to the historical use of *C. timorensis* as a remedy for various epidemic diseases, a fact reinforced by explicit acknowledgment from the Institute of Ayurveda in Sri Lanka (Ayurvedic Medicinal Plants of Sri Lanka Compendium Version 3, 2003). This study endeavors to comprehensively evaluate the antibacterial efficacy of aqueous extracts derived from the leaves, bark, and stem of *C. timorensis* Span, thereby contributing valuable insights into its potential therapeutic applications.



Fig. 1. *Celtis timorensis* plant Fig. 2. *Celtis timorensis* leaves

2. MATERIALS AND METHODS

2.1. Sample Collection

Well-grown and fully expanded mature fresh leaves stem, and bark parts of *Celtis timorensis* were collected during daytime within the natural flowering months of the plant from an estate in Peradeniya district in Central Province, Sri Lanka (latitude of 7° 15' 9.00" N and longitude of 80° 35' 28.79" E).

2.2. Identification and Authentication of the plant

Properly rinsed, dried, and pressed specimens were sent to the National Herbarium, Peradeniya, Sri Lanka for authentication.

2.3. Preparation of crude plant material extracts

20g of each part of *C. timorensis* was ground into a fine powder and added to 100 ml of solvent. The prepared samples were filtered, and the filtrate was considered the stock crude solution. The extracts were dissolved in distilled water and a dilution series was prepared (200 mg/ml, 150 mg/ml, 100 mg/ml, and 50 mg/ml).



Fig. 3. Maceration of Bark, stem and root extracts of *C.timorensis*

2.4 Evaluation of in vitro antibacterial activity

2.4.1 Collection and sub-culturing of test microorganisms

Pathogenic strains of *Escherichia coli* (*E. coli*)(ATCC 25922), *Pseudomonas aeruginosa* (*P. aeruginosa*) (ATCC 27853), and *Staphylococcus aureus* (*S. aureus*) (ATCC 25923) were acquired from the Medical Research Institute, Colombo 08, Sri Lanka (Hwang, Jung Ho, Sang-Young Lee,2021). Subsequently, these strains were sub cultured on nutrition agar and stored at a temperature of 4°C for maintenance purposes.

2.4.2 Preparation of test solutions

The test solutions underwent the preparation of a concentration gradient, comprising concentrations of 1000 µg/ml, 750 µg/ml, 500 µg/ml, and 250 µg/ml

2.4.3 Preparation of Gentamicin antibiotic for positive control

Gentamicin was used as a positive control in the study. A 125 µl volume of Gentamicin was measured and transferred into a volumetric flask. The solution was replenished to a final volume of 100 ml with sterile distilled water, resulting in a final concentration of 50 µg/ml. (Senadeera, Nimesha & Fernando, K & Wickramasekara, W *et al.*, 2022).

2.4.4 Preparation of McFarland Standards

A 0.5 McFarland standard is a bacterial suspension containing 1x10⁸ to 2x10⁸ CFU/ml. It was prepared in the lab by adding BaCl₂ to H₂SO₄ with constant stirring. (Muthusarayanan, S., Sivarajasekar, N., Vivek, J.S *et al.*, 2018).

2.4.5 Preparation of bacterial broth

Bacterial broths were prepared using subcultures 24 hours prior. Saline was used as the medium and bacterial colonies were acquired using an inoculating loop and dissolved in saline under aseptic conditions. (Hwang, Jung Ho, Sang Young Lee, 2021).

2.4.6 Preparation of culture media

38.04 grams of Mueller-Hinton Agar were dissolved in distilled water, heated, and aliquoted into 100 ml conical flasks. The flasks were autoclaved at 121°C for 15 minutes, and the glassware was sterilized in a hot air oven at 170°C for 60 minutes (Vinny R. Sastri, 2022). These steps collectively produce a microbiologically pristine Mueller-Hinton Agar medium suitable for various microbiological applications, particularly in antibiotic susceptibility testing (Yao, Tatsuma and Asayama, Yuta, 2017).

2.5 Determination of antimicrobial assay

2.5.1 Inoculum standardization

All bacterial strains were inoculated in Müller-Hinton broth for about 24 hours (Vaseekaran, S., Balakumar, S. and Arasaratnam, V., 2010). Bacterial strains were cultured at 37°C on slopes of nutrient agar. All the procedures were conducted under aseptic conditions. The glassware that was used underwent sterilization.

2.5.2 Assay of antibacterial activity using agar well diffusion method

Agar was applied to petri plates, followed by a bacterial-seeded agar layer. The plates were treated with plant extract concentrations and positive/negative controls. After incubation, the inhibition zones were measured using a digital vernier caliper, and the data was averaged from three independent trials for precision (Abdelaziz, A. A. *et al.*, 2012, Souza-Filho, F.J.D., Soares, A.D.J., Vianna, M.E., *et al.*, 2008).

2.6 Statistical analysis

The acquired data underwent statistical analysis using SPSS version 23, and dose-response curves were generated using GraphPad Prism 8.0.1 software.

3.0 Results

3.1 Evaluation Of Levels Of Antimicrobial Efficacy Of *Celtis timorensis* Span. Plant Parts

3.1.1 Antibacterial effect of *Celtis timorensis* Span. against *Escherichia coli* (ATCC 25922)

Table 1. Antibacterial effect of aqueous extracts of *C. timorensis* leaves, bark and stem against *E. coli*

Antimicrobial effect of <i>Celtis timorensis</i> Span.	Zone of inhibition (mm)					
	1000 µg/ml	750µg/ml	500 µg/ml	250 µg/ml	Positive control	Negative control
Leaves	14.05±0.13	13.10±0.09	10.96±0.04	9.75±0.10	17.06±0.15	8.00±0.0
Bark	14.20±0.07	12.16±0.08	10.82±0.02	9.38±0.08	16.98±0.23	8.00±0.0
Stem	14.85±0.13	13.55±0.13	11.00±0.05	10.05±0.09	17.18±0.14	8.00±0.0

As depicted in table 01, the highest antibacterial efficacy against *E. coli* was observed in the aqueous stem extract at a concentration of 1000 µg/mL, resulting in a substantial inhibitory zone of 14.85 mm. In contrast, the aqueous leaf extract exhibited the lowest antibacterial activity, while the bark extract demonstrated a moderate effect. Mean inhibition zone measurements for the bark and leaf extracts at a concentration of 1000 µg/mL were recorded as 14.20 mm and 14.05 mm, respectively.

At a concentration of 1000 µg/mL, all three extracts—leaves, bark, and stem—exhibited their most significant antibacterial effects. Furthermore, these extracts demonstrated consistent antibacterial activity across all tested concentrations (1000 µg/mL, 750 µg/mL, 500 µg/mL, 250 µg/mL), underscoring their efficacy against *E. coli*.

3.1.2 Antibacterial effect of *Celtis timorensis* Span. against *Pseudomonas aeruginosa* (ATCC 27853)

Table 2. Antibacterial effect of aqueous extracts of *Celtis timorensis* leaves, bark and stem against *P. aeruginosa*

Antimicrobial effect of <i>Celtistimo rensis</i> Span.	Zone of inhibition (mm)					
	1000 $\mu\text{g/ml}$	750 $\mu\text{g/ml}$	500 $\mu\text{g/ml}$	250 $\mu\text{g/ml}$	Positive control	Negative control
Leaves	11.80 \pm 0.33	10.70 \pm 0.35	9.62 \pm 0.51	8.00 \pm 0.0	15.12 \pm 0.42	8.00 \pm 0.0
Bark	11.76 \pm 0.20	10.25 \pm 0.07	8.46 \pm 0.21	8.00 \pm 0.0	15.91 \pm 0.15	8.00 \pm 0.0
Stem	11.91 \pm 0.39	10.38 \pm 0.15	8.38 \pm 0.14	8.00 \pm 0.0	16.08 \pm 0.30	8.00 \pm 0.0

As depicted in table 03, the evaluation of antibacterial activity against *Pseudomonas aeruginosa*, the aqueous extraction from the stem of the plant displayed the highest efficacy, producing a notable 11.91 mm zone of inhibition at a concentration of 1000 $\mu\text{g/ml}$. In contrast, the bark extraction exhibited the least substantial effect, with the leaf extract falling in between at 11.76 mm and 11.80 mm, respectively. Interestingly, all three extracts demonstrated discernible antibacterial effects at concentrations of 750 $\mu\text{g/ml}$ and 500 $\mu\text{g/ml}$, yet none exhibited activity at the lower concentration of 250 $\mu\text{g/ml}$.

3.1.3 Antibacterial effect of aqueous extracts of *Celtis timorensis* leaves, bark and stem against *Staphylococcus aureus*

Table 3. Antibacterial effect of aqueous extracts of *C. timorensis* leaves, bark and stem against *S. aureus*

Antimicrobial effect of <i>Celtistimorensis</i> Span	Zone of inhibition (mm)					
	1000 $\mu\text{g/ml}$	750 $\mu\text{g/ml}$	500 $\mu\text{g/ml}$	250 $\mu\text{g/ml}$	Positive control	Negative control
Leaves	8.00 \pm 0.0	8.00 \pm 0.0	8.00 \pm 0.0	8.00 \pm 0.0	19.58 \pm 0.37	8.00 \pm 0.0

Bark	8.00±0.0	8.00±0.0	8.00±0.0	8.00±0.0	19.08±0.15	8.00±0.0
Stem	8.00±0.0	8.00±0.0	8.00±0.0	8.00±0.0	19.75±0.14	8.00±0.0

3.1.4 Antibacterial effect of *Celtis timorensis* Span. against *Staphylococcus aureus* (ATCC 25923)

None of the extracts of leaves or bark or stem exhibited any discernible antimicrobial effect against *S. aureus*.

Dose-Response Study

3.1.5 Dose-response curves and EC₅₀ values of aqueous extractions of *Celtis timorensis* Span. plant parts against *Escherichia coli* (ATCC 25922)




Fig. 4. Dose-response curves of *C. timorensis* leaves, bark and stem extracts against *E. coli*.

3.1.6 Dose-response curves and EC₅₀ values of aqueous extractions of *Celtis timorensis* Span. plant parts against *Pseudomonas aeruginosa* (ATCC 27853)

log concentration vs Zone of Inhibition against *Pseudomonas aeruginosa*




Fig. 5. Dose-response curves of *C. timorensis* leaves, bark and stem extractions against *P. aeruginosa*.

According to the data obtained by EC_{50} values and Dose-Response curves, aqueous extract of leaves exhibited the highest efficacy and lowest potency and the aqueous extract of bark exhibited the highest potency and lowest efficacy against *P. aeruginosa*.

4. DISCUSSION

The current study represents the inaugural investigation into the antibacterial properties of aqueous extracts from various components (leaves, stem, and bark) of *Celtis timorensis*. Grounded in existing literature, this research endeavors to assess the in vitro antibacterial activity of these aqueous extracts, marking the first-of-its-kind exploration in the scientific domain.

The previous studies evaluated the unexplored antibacterial properties of alcoholic extractions of *C. timorensis*, renowned in Sinhala Ayurveda for its strong antimicrobial effects (Gunawardena, D. C., 1975; Hewage, C.M., Bandara, B.M.R., Karunaratne *et al.*, 1998). Despite recognition, the antibacterial activity of aqueous macerations from specific plant parts remains scientifically untested. Notably, Ayurvedic literature underscores *C. timorensis*'s traditional use in treating various skin infections, corroborated by recent studies emphasizing its wound healing attributes due to antimicrobial properties (PKumar, P.M., Suba, V. and Reddy, R.B., 2017). The primary aim of this study was to conduct an antibacterial assay against wound-infecting bacteria, focusing on *Escherichia coli* (ATCC 25922), *P. aeruginosa* (ATCC 27853), and *Staphylococcus aureus* (ATCC 25923).

As depicted in table 01, the antibacterial assay against *E. coli* revealed that the 1000 $\mu\text{g/ml}$ concentration of all three plant extracts (Leaves, Bark, and Stem) exhibited the highest zone of inhibition, demonstrating a direct proportional relationship between plant extract concentration and inhibition zone size. The aqueous extraction of the stem displays the maximum zone of inhibition (14.85 mm), while bark and

leaves show zones of 14.20 mm and 14.05 mm, respectively. All three parts exhibit statistically significant differences ($p < 0.05$) in their mean zones of inhibition against *E. coli*, although all are lower than the positive control.

Similarly, the antibacterial assay against *P. aeruginosa* indicates that the aqueous extraction of the stem achieves the highest zone of inhibition (11.91 mm), while bark and leaves display zones of 11.80 mm and 11.76 mm, respectively (table 02). All three parts exhibit statistically significant differences ($p < 0.05$) in their mean zones of inhibition against *P. aeruginosa*, with values below the positive control.

As delineated in table 03 against *S. aureus*, none of the concentrations for any plant part show positive antibacterial responses, in contrast to the positive control (approximately 19.4 mm). Consequently, aqueous extractions of leaves, bark, and stem of *C. timorensis* exhibit positive antibacterial responses against gram-negative bacteria but lack efficacy against gram-positive *S. aureus*.

In summary, the study concludes that aqueous extractions of *C. timorensis* leaves, bark, and stem demonstrate positive antibacterial responses against gram-negative bacteria (*E. coli* and *P. aeruginosa*) but lack efficacy against gram-positive *S. aureus*. The potency and biological effect vary among plant parts, with leaves displaying the highest potency against *E. coli* and bark against *P. aeruginosa*.

5. CONCLUSION

The current study investigated the in-vitro antimicrobial activity of *C. timorensis* Span. plant components (leaves, bark, and stem) against wound infecting bacteria—*Escherichia coli*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*. While gram-negative bacteria were susceptible to aqueous extracts, *Staphylococcus aureus*, gram-positive bacteria showed resistance. The cylinder plate method revealed the highest inhibition zones against *Escherichia coli* and *Pseudomonas aeruginosa* with 1000 g/ml concentrated aqueous extraction of stem and the inhibition zones were 14.85 mm and 11.91 mm respectively. Concentration and the inhibition zones are directly correlated with all three plant sections. Statistical analysis indicated significant differences among plant components against *E. coli* and *P. aeruginosa*. Dose-response data identified aqueous extraction of bark as most efficacious against *E. coli* and aqueous extraction of leaves against *P. aeruginosa*. Overall, *C. timorensis* leaves and bark displayed robust antibiotic activity against gram-negative bacteria, particularly *E. coli* and *P. aeruginosa*.

References

- Saga, T. and Yamaguchi, K., 2009. History of antimicrobial agents and resistant bacteria. *Jmaj*, 52(2), pp.103-108.
- Gupta, P.D. and Birdi, T.J., 2017. Development of botanicals to combat antibiotic resistance. *Journal of Ayurveda and integrative medicine*, 8(4), pp.266-275.
- *celtistimorensis*- Medicinal Plant Names Services(2023). <https://mpns.science.kew.org/mpns-portal/backToSearch?query=celtis+timorensis&filter=&fuzzy=false&nameType=all>.
- <https://indiabiodiversity.org/species/show/264069>, 2023
- Dasari, R., Kunchibhotla, A., Sathyavathi, D. and Reddy, P.J., 2013. Evaluation of methanolic extract of *Celtis timorensis* for its antidepressant activity. *Indo Am J Pharm Res*, 3, pp.2533-8.
- http://www.instituteofayurveda.org/plants/plants_detail.php?i=1313&s=Scientific_name, 2023
- Senadeera, Nimesha & Fernando, K & Wickramasekara, W & Fernando, M & Ranaweera, Chathuranaga & Rajapaksha, Weranga & Silva, A. (2022). In vitro Anti-inflammatory Activity of

Endemic *Artocarpus nobilis* Thw Found in Sri Lanka. *Asian Plant Research Journal*. 116-122. 10.9734/APRJ/2021/v8i430192.

- Muthusaravanan, S., Sivarajasekar, N., Vivek, J.S., Paramasivan, T., Naushad, M., Prakashmaran, J., Gayathri, V. and Al-Duaij, O.K., 2018. Phytoremediation of heavy metals: mechanisms, methods and enhancements. *Environmental chemistry letters*, 16, pp.1339-1359.
- Hwang, Jung Ho, Sang Young Lee *et al.* 2021. "Microscopic Analysis of Bacterial Inoculum Effect Using Micropatterned Biochip" *Antibiotics* 10, no. 3: 300. <https://doi.org/10.3390/antibiotics10030300>
- Vinny R. Sastri, 2022. Material Requirements for Plastics Used in Medical Devices. In *Plastics Design Library, Plastics in Medical Devices*, 3, pp. 65-112.
- Yao, Tatsuma; Asayama, Yuta (2017). *Animal-cell culture media: History, characteristics, and current issues. Reproductive Medicine and Biology*, 16(2), 99–117. doi:10.1002/rmb2.12024
- Vaseekaran, S., Balakumar, S. and Arasaratnam, V., 2010. Isolation and identification of a bacterial strain producing thermostable alpha-amylase.
- Abdelaziz, A.A., Elbanna, T.E. and Gamaledeen, N.M., 2012. validated microbiological and HPLC methods for the determination of moxifloxacin in pharmaceutical preparations and human plasma. *Brazilian journal of microbiology*, 43, pp.1291-1301.
- Souza-Filho, F.J.D., Soares, A.D.J., Vianna, M.E., Zaia, A.A., Ferraz, C.C.R. and Gomes, B.P.F.D.A., 2008. Antimicrobial effect and pH of chlorhexidine gel and calcium hydroxide alone and associated with other materials. *Brazilian dental journal*, 19, pp.28-33.
- Gunawardena, D. C. (1975). Medicinal and Economic Plants of the Musaeum Zeylanicum of Paul Hermann. *Journal of the Sri Lanka Branch of the Royal Asiatic Society*, 19, 33–48. <http://www.jstor.org/stable/23728486>
- Hewage, C.M., Bandara, B.M.R., Karunaratne, V., Wannigama, G.P., Pinto, M.R.M. and Wijesundara, D.S.A., 1998. Antibacterial activity of some medicinal plants of Sri Lanka.
- Kumar, P.M., Suba, V. and Reddy, R.B., 2017. Wound healing activity of *Celtis timorensis* Span.(Cannabaceae) leaf extract in Wistar albino rats.