

Nutrient Status of coastal soils-24 Parganas (South), West Bengal, India

ABSTRACT

The study conducted in the year 2023-24 at the Department of Soil Science and Agricultural Chemistry, The Neotia University, 24 Parganas (South), situated in coastal region of West Bengal, aimed to assess the nutrient status of the soils in the instructional farm. Soil samples (105 No) were collected at 0-15 cm depth and analyzed for pH, electrical conductivity (EC), organic carbon (OC), and major nutrients such as available nitrogen, phosphorus, and potassium. The result showed that the soils were neutral in reaction and saline in nature, whereas organic carbon and available nitrogen content were at a medium status, but available phosphorus and potassium status were high. Moreover organic carbon showed negative correlation with soil electrical conductivity (-0.384, $p < 0.05$). The nutrient index values for available nitrogen and available phosphorus were between 1.66-2.33 range, categorizing it as medium fertile. Conversely, the nutrient index value for available potassium exceeded 2.33, classifying it as high fertile.


Keywords: Coastal region; saline soil; major nutrients; nutrient index; fertility.

1 INTRODUCTION

Coastal soils play a vital role in the agricultural productivity of West Bengal, as they are heavily influenced by their proximity to the ocean and unique environmental conditions [1] (Mallik, 2020). These soils are generally deficient in organic matter and nutrients, poor physical properties and saline in nature [2] (Elayaraja *et.al*, 2023), [3] (Mandal *et.al*, 2019), as well as high levels of soil salinity due to factors like sub-soil saline water, poor drainage, and intrusion of sea water. Moreover, it is found that the suitability of coastal soils for specific crops is varied, for example, while paddy crops were found to be well-suited for these soils, other crops like chilli, mustard, sunflower, and vegetables may face challenges in terms of nutrient availability and tolerance to the high salinity levels. However, there is a lack of detailed information regarding the coastal soils of The Neotia University School of Agriculture and Allied Sciences in West Bengal. This study aims to assess the nutrient status of these coastal soils to provide valuable insights for effective soil management strategies.

2 MATERIALS AND METHODS

The research was conducted at Instructional Farm of The Neotia University (TNU), Sarisha, Diamond Harbour, West Bengal, India, located at 22°48' N latitude and 88°31' E longitudes with an average altitude of 8 m above the mean sea level (MSL) (Figure 1). The main crops grown within the study area are rice, maize, sunflower, groundnut, mustard, mungbean and black gram. The farm encompasses seven blocks viz. Uncultivated land (S1), Cultivated land (S2), Net house (S3), Poly house (S4), Upland (S5), Lowland (S6) and Orchard field (S7).



32

33

Figure 1: Soil Sampling area in TNU

34 For the study, 105 soil samples (15 representative soil samples from each seven blocks)
 35 from surface soil (0-15 cm depth) were collected from the instructional farm. These samples
 36 were air-dried, crushed using a wooden mallet, and sieved. After labeling, the samples were
 37 stored in plastic container for subsequent analysis. The analysis involved standard
 38 procedures to determine the available nutrient status. This included measuring of soil
 39 reaction (1:2.5; Soil: water), electrical conductivity (1:2.5; Soil: water), organic carbon via
 40 wet chromic acid digestion [4](Walkley and Black, 1934), available nitrogen through alkaline
 41 permanganate method [5] (Subbiah and Asija, 1956), available phosphorus via 0.5 M
 42 sodium bicarbonate [6] (Olsen et al., 1954), and available potassium using the neutral
 43 normal ammonium acetate method [7] (Stanford and English, 1949). Soil nutrient index was
 44 evaluated for the soil samples analyzed based on the formula suggested by [8] (Parker *et*
 45 *al.*, 1951) as given below:

46

$$\text{Soil Nutrient Index (SNI)} = \frac{N_1 \times 1 + N_2 \times 2 + N_3 \times 3}{N_T}$$

47

where,

48

N_1 = Number of samples falling in low class of nutrient status;

49 N_2 = Number of samples falling in medium class of nutrient status; and
50 N_3 = Number of samples falling in high class of nutrient status.
51 N_T = Total number of samples.

52

53 Separate indices were calculated for different nutrients like N, P and K. The soils were rated
54 as per the SNI values as low (<1.67), medium (1.67 to 2.33) and high (>2.33) (Parker *et al.*
55 1951).The database on analysis of soil available nutrient content was developed by using
56 Microsoft Excel. Descriptive statistical parameters viz., mean, range, standard deviation and
57 correlation of various soil parameters were computed using SPSS 22.0.

58

59

60 **3RESULTS AND DISCUSSION**

61

62 **3.1 Soil pH**

63

64 Soil pH, a critical property influencing nutrient availability and plant growth, was measured
65 across seven blocks (S1 to S7) of the study area. The pH values varied within and between
66 these blocks, ranging from 6.46 to 7.89 (Table1; Figure 2). Specifically, S1 recorded pH
67 values between 7.25 and 7.71, S2 from 6.56 to 7.47, S3 from 6.46 to 7.27, S4 from 6.46 to
68 7.66, S5 from 6.53 to 7.89, S6 from 6.75 to 7.80 and S7 from 6.95 to 7.58. The mean pH
69 values for these blocks were 7.55, 6.78, 6.94, 6.97, 7.01, 7.3 and 7.21 respectively. These
70 results indicated that the soils fall under neutral to slightly alkaline category. These results
71 are in agreement with [9] (Sarkar *et al.*, 2001) who reported the pH of the coastal region
72 ranged from 5.3 to 8.1.

73

74 **3.2 Electrical Conductivity**

75

76 In the study, the electrical conductivity (EC) of soil was analyzed across seven blocks (S1 to
77 S7). The EC values varied as follows: S1 ranged from 0.218 to 0.481 dSm^{-1} , S2 from 2.30 to
78 2.99 $dS m^{-1}$, S3 from 0.58 to 0.77 $dS m^{-1}$, S4 from 2.22 to 2.55 $dS m^{-1}$, S5 from 1.30 to 1.85
79 $dS m^{-1}$, S6 from 1.028 to 1.63 $dS m^{-1}$ and S7 from 2.081 to 2.367 $dS m^{-1}$. The mean EC
80 values for these blocks were recorded as 0.388, 2.608, 0.676, 2.346, 1.555, 1.285 and 2.223
81 $dS m^{-1}$, respectively (Table1; Figure 2). These measurements indicate that the mean
82 electrical conductivity of the surface soil across all blocks fell within the range of 0.388 to
83 2.608 dSm^{-1} categorizing the soils as non saline to saline in nature. The study revealed that
84 all soil samples had high electrical conductivity (EC) except for S1, S3 and S6 samples,
85 indicating a saline nature with high salt concentration. These findings are consistent with
86 those reported by [10, 11, 12] (Bandyopadhyay *et al* ,1988, and 2003, Muhr *et al*, 1965).

87

88 **3.3 Organic Carbon**

89

90 The recorded organic carbon content in the soil varied across different blocks. In S1, the
91 range was 0.52 to 0.59 percent; in S2 it was 0.44 to 0.58 percent; in S3 0.45 to 0.58 percent;
92 in S4 0.45 to 0.59 percent; in S5 and S6 0.41 to 0.59 percent respectively and in S7, 0.44 to
93 0.59 percent. The average organic carbon content across these seven blocks ranged from
94 0.41 to 0.59 percent (Table1; Figure 2). This indicates that the soil organic carbon levels in

95 these blocks generally fell into the low to medium category. The correlation analysis
96 revealed a negative association between electrical conductivity (EC) and organic carbon ($r =$
97 -0.384 ; $p < 0.05$) (Table 2). Tripathi *et al.* (2006) [13] found a decrease in soil organic carbon
98 content with increasing salinity ($r = -0.38$; $p < 0.01$), a trend also observed by [14] (Kaur *et*
99 *al.* 1998), who reported a significant negative relationship between organic carbon and EC.
100 The organic carbon content in the studied area was found to be in medium range, consistent
101 with findings by [15] (Joshi and Kadrekar, 1987), who observed variations in organic carbon
102 in coastal soils ranging from $< 0.5\%$ to $> 0.75\%$.

103

104 **3.4 Available Nitrogen**

105

106 In the present study, the average available nitrogen content in the surface soil across
107 different blocks showed varied result. The mean values recorded for each block were as
108 follows: $235.18 \text{ kg ha}^{-1}$ in S1, $339.04 \text{ kg ha}^{-1}$ in S2, $318.05 \text{ kg ha}^{-1}$ in S3, $324.36 \text{ kg ha}^{-1}$ in S4,
109 $329.35 \text{ kg ha}^{-1}$ in S5, $327.48 \text{ kg ha}^{-1}$ in S6 and $316.72 \text{ kg ha}^{-1}$ in S7. The range of available
110 nitrogen in these blocks also showed considerable variation, with S1 ranging from 210.48 to
111 $259.24 \text{ kg ha}^{-1}$, S2 from 300.43 to $386.50 \text{ kg ha}^{-1}$, S3 from 286.2 to $374.01 \text{ kg ha}^{-1}$, S4 from
112 280.23 to $370.40 \text{ kg ha}^{-1}$, S5 from 258.62 to $381.62 \text{ kg ha}^{-1}$, S6 from 281.48 to $398.63 \text{ kg ha}^{-1}$
113 and S7 from 278.74 to $376.50 \text{ kg ha}^{-1}$ (Table 1; Figure 3). Notably, S2 had the highest mean
114 available nitrogen content of 339.04 kg/ha , while S1 had the lowest content of $235.18 \text{ kg ha}^{-1}$.
115 The soils in S1 were categorized as low in terms of available nitrogen status, whereas all
116 other blocks were classified as medium in available nitrogen status. These findings are
117 consistent with those reported by Ray *et al.*, 2014 [16]

118

119 **3.5 Available Phosphorus (P_2O_5)**

120

121 The study revealed considerable variation in the available phosphorus content in the soil
122 across different blocks. The mean available phosphorus content in each block also varied,
123 with S1 having a mean of 20.24 kg ha^{-1} , S2 54.98 kg ha^{-1} , S3 33.10 kg ha^{-1} , S4 35.03 kg ha^{-1} ,
124 S5 51.75 kg ha^{-1} , S6 50.99 kg ha^{-1} and S7 25.11 kg ha^{-1} (Table 1; Figure 3). Interestingly, S2
125 recorded the highest mean available phosphorus content at 54.98 kg/ha , while S1 had the
126 lowest at 20.24 kg ha^{-1} . In terms of classification, the available phosphorus content was low
127 in S1 and S7, while it was categorized as medium in the remaining five blocks. Shahandeh *et*
128 *al.*, (2003) [17] stated that the reduced condition of soils increased the availability of soil
129 phosphorus due to increased solubility of Fe-associated P and Mn-associated P.

130

131 **3.6 Available Potassium (K_2O)**

132

133 The study highlighted variations in the soil's available potassium content across different
134 blocks. The mean available potassium content in each block was as follows: S1 had 267.72
135 kg ha^{-1} , S2 $312.26 \text{ kg ha}^{-1}$, S3 $290.63 \text{ kg ha}^{-1}$, S4 $311.57 \text{ kg ha}^{-1}$, S5 $364.28 \text{ kg ha}^{-1}$, S6
136 $412.03 \text{ kg ha}^{-1}$, S7 $310.43 \text{ kg ha}^{-1}$ (Table 1; Figure 3). S6 recorded the highest mean available
137 potassium content at $412.03 \text{ kg ha}^{-1}$, while S1 had the lowest value at $267.72 \text{ kg ha}^{-1}$.
138 Despite these variations, all the blocks were categorized as having a high level of available
139 potassium status except S1 fell under low category. The results were consistent with the
140 findings by Mahajan *et al.*, (2015) [18].

141
142

Table: 1 Soil properties of different blocks of the Instructional farm, TNU (mean±sd)

Block No.	pH	EC (dSm ⁻¹)	OC (%)	Available Nitrogen (kg ha ⁻¹)	Available P ₂ O ₅ (kg ha ⁻¹)	Available K ₂ O (kg ha ⁻¹)
1	7.55±0.14	0.39±0.10	0.55±0.02	235.18±15.68	20.24±0.18	267.72±2.33
2	6.78±0.24	2.61±0.25	0.52±0.05	339.04±30.73	54.98±0.35	312.26±4.56
3	6.94±0.28	0.68±0.07	0.53±0.04	318.05±28.25	33.10±0.32	290.63±4.19
4	6.97±0.37	2.35±0.08	0.52±0.05	324.36±32.15	35.03±0.36	311.57±4.77
5	7.01±0.44	1.56±0.21	0.49±0.06	329.35±36.46	51.75±0.41	364.28±5.41
6	7.30±0.38	1.28±0.14	0.52±0.06	327.48±37.76	50.99±0.42	412.03±5.60
7	7.21±0.21	2.22±0.09	0.53±0.06	316.72±36.47	25.11±0.41	310.43±5.41


143

144 **Table 2: Pearson's Correlation of analysed soil properties of TNU**

Parameter	pH	EC (dS m ⁻¹)	OC (%)	Available Nitrogen (kg ha ⁻¹)	Available P ₂ O ₅ (kg ha ⁻¹)	Available K ₂ O (kg ha ⁻¹)
pH	1.00					
EC (dS m ⁻¹)	-0.372*	1.00				
OC (%)	0.463*	-0.384*	1.00			
Available Nitrogen (kg ha ⁻¹)	-0.738*	0.675*	-0.698*	1.00		
Available P ₂ O ₅ (kg ha ⁻¹)	-0.408*	0.397*	-0.820*	0.730*	1.00	
Available K ₂ O (kg ha ⁻¹)	0.017*	0.161*	-0.653*	0.553*	0.710*	1.00

145 *. Correlation is significant at the 0.05 level (2-tailed).


146
147



148

149

Figure 2. Physico-chemical parameters of different soil in TNU



150

151

152

Figure 3. Macro nutrient content of different soil in TNU

153 3.7 Soil Nutrient Index

154

155 A comprehensive understanding of soil nutrient levels is crucial for ensuring optimal crop
156 production and soil health. By analyzing the nutrient index report, farmers and land
157 managers can make informed decisions regarding fertilizer application, crop selection, and
158 soil management practices. This report aims to provide an in-depth analysis of the soil's
159 nutrient levels, helping to optimize agricultural productivity and sustainability. The values for
160 available N, P, and K worked out from SNI were 1.83, 1.71 and 2.86 respectively, against
161 the nutrient index values < 1.67 for low, 1.67 to 2.33 for medium and > 2.33 for high fertility
162 status of the area[8] (Parker *et al*,1951). Table 3 presents the calculated Nutrient Index
163 Values and Fertility Ratings for the soils of Instructional Farm of The Neotia University.

164

165 **Table 3: Nutrient Index and Fertility Status of Soil**

Available nutrients	Nutrient Index	Fertility Status
Nitrogen (kg ha ⁻¹)	1.83	Medium
P ₂ O ₅ (kg ha ⁻¹)	1.71	Medium
K ₂ O(kg ha ⁻¹)	2.86	High

166

167

168 4CONCLUSION

169

170 The physico-chemical characteristics and nutrient status of coastal soil of Instructional farm,
171 The Neotia University revealed that the soil were neutral in soil reaction and saline. While
172 organic carbon and available nitrogen content were medium, available phosphorus and
173 potassium status were high. The nutrient index for available nitrogen and available
174 phosphorus was medium and for available potassium it was high.Hence to enhance soil
175 fertility, it is imperative to replenish nutrients through the application of organic matter, green
176 manures, and inorganic fertilizers. Adopting a comprehensive nutrient management
177 approach, can ensure balanced nutrition for crops, thereby sustaining soil health and
178 maximizing crop yields in this region.

179

180

181 AUTHORS' CONTRIBUTIONS

182

183 This work was carried out in collaboration among all authors. All authors read and approved
184 the finalmanuscript.

185

186 REFERENCES

187

188

189 1. Mallik TK. Salient features of West Bengal coastal ecosystem and effective
190 management of environment. *MedCrave Group*, 2020.

- 191 2. Elayaraja D, Senthilvalavan P and Kamalakannan P. Fortified Organic Manure and
192 NPK Fertilizer Levels: Influence on Soil Properties, Yield and Micronutrients
193 Biofortification of Brinjal in Coastal Soil International Journal of Plant & Soil Scienc.
194 2023; 35(19): 2027-2041.
- 195 3. Mandal UK, Burman D, Bhardwaj AK, Nayak DB, Samui A, Mullick S, Mahanta KK,
196 Lama TD, Maji B, Mandal S, Raut S, Sarangi SK.. Waterlogging and coastal salinity
197 management through landshaping and cropping intensification in climatically
198 vulnerable Indian Sundarbans. *Agricultural Water Management*. 2019; 12–26.
- 199 4. Walkley A, Black CA. An examination of digestion method for determining soil
200 organic matter and the proposed modification of the chromic acid titration method.
201 *Soil Science*. 1934;37:29–38.
- 202 5. Subbiah V, Asija GL. A rapid procedure for estimation of available nitrogen in soil
203 *Current Science*. 1956;5:259-260
- 204 6. Olsen SR, Watanabe FS, Cole CV, Dean LA. Estimation of Available P in Soils by
205 Extraction with Sodium Bicarbonate. U.S.D.A. Circular No. 939;1954.
- 206 7. Stanford S, English L. Use of flame photometer in rapid soil tests of K. *Canadian*
207 *Journal of Agronomy*. 1949;41:446-447.
- 208 8. Parker FW, Nelson WL, Winters E, Miles IE. The broad interpretation and
209 application of soil test information. *Agronomy Journal*. 1951;43(3):105-112.
- 210 9. Sarkar D, Sahoo AK, Sah KD and Gajbhiye KS (2001). Coastal soils of eastern
211 India-their charactersistics,
212 potentials and limitations towards alternate land use. *Journal of Indian Society of*
213 *Coastal Agricultural Research* **19(1&2)**: 80-83
- 214 10. Bandyopadhyay AK, Bhargava GP, Rao KGK, Sen HS, Sinha TS, Bandyopadhyay
215 BK, Biswas CR, Bal AR, Dutt SK and Mandal RC (1988). *Coastal Saline Soils of*
216 *India and Their Management*, Bull No. **13**. Central Soil Salinity Research Institute,
217 RRS Canning, West Bengal, pp. 158.
- 218 11. Bandyopadhyay BK, Maji B, Sen HS and Tyagi NK (2003). *Coastal Soils of West*
219 *Bengal- Their Nature, Distribution and Characteristics*. Bull. No. **1/2003**. Central Soil
220 Salinity Research Institute, Regional Research Station, Canning Town, West
221 Bengal, India, pp 1-62.
- 222 12. Muhr GR, Dutta NP, Sankara SH, Liley VK, and Donahue RR. (1965). *Soil Testing in*
223 *India*. US Agency for International Development, New Delhi.
- 224 13. Tripathi S, Kumari S, Chakroborty A, Gupta A, Chakroborty K and Bandyopadhyay
225 BK.(2006). Microbial biomass and its activity in salt affected coastal soils. *Biology*
226 *and Fertility of Soils* **42**: 273-277.
- 227 14. Kaur B, Aggarwal AK and Gupta SR (1998). Soil microbial biomass and nitrogen
228 mineralization in salt affected soils. *International Journal of Ecology and*
229 *Environmental Sciences* **24**: 103-111.
- 230 15. Joshi RG and Kadrekar SB (1987). Fertility status of coastal salt affected soils of
231 Maharashtra. *Journal of Indian Society of Coastal Agricultural Research* 5(1):111-
232 116.
- 233 16. Ray, P., Meena, B.L. and Nath, C. P. (2014). Management of Coastal Soils for
234 Improving Soil Quality and Productivity. *Popular Kheti*. 2(1):95-99.
- 235 17. Shahandeh, H., Hossner, L. R and Turner, F. T. (2003). Phosphorus relationships
236 to manganese and iron in rice soils. *Soil Science*. 168(7): 489-500.

- 237 18. Mahajan, G. R., B.L.Manjunath, A.M. Latore, Ruenna D'Souza, Shashi Vishwakarma,
238 N.P. Singh. 2015. Fertility Status of the Unique Coastal Acid Saline Soils of Goa.
239 Journal of the Indian Society of Soil Science, 63(2): 232-237
240
241