

## Original Research Article

### **Effect of organic manures and bio-fertilizers on growth, yield and quality of kharif cowpea [*Vigna unguiculata* (L.)]**

#### **ABSTRACT**

A field experiment was conducted at Agronomy Instructional Farm of SDAU, Sardarkrushinagar during *kharif* season of 2020-2021. The experiment consisting nine treatments viz., T<sub>1</sub>-Absolute control, T<sub>2</sub>-100% RDF (20 kg N + 40 kg P<sub>2</sub>O<sub>5</sub>/ha), T<sub>3</sub>-FYM @ 5.0 t/ha, T<sub>4</sub>-FYM @ 2.5 t/ha + *Rhizobium* + PSB, T<sub>5</sub>-vermicompost @ 2.0 t/ha, T<sub>6</sub>-vermicompost @ 1.0 t/ha + *Rhizobium* + PSB, T<sub>7</sub>-castor cake @ 2.0 t/ha, T<sub>8</sub>-castor cake @ 1.0 t/ha + *Rhizobium* + PSB, T<sub>9</sub>-*GhanJeevamrut* @ 250 kg/ha at sowing time + Seed treatment with *Beejamrut* @ 200 ml/kg seed + *Jeevamrut* @ 500 lit/ha with irrigation at sowing and 30 DAS were tested in randomized block design with four replications. The result revealed that the application of T<sub>8</sub>-castor cake @ 1.0 t/ha + *Rhizobium* + PSB showed significant improvement on plant height, number of branches per plant, dry matter accumulation, number of pods per plant, number of seeds per pod, length of pod and seed yield per plant whereas, application of T<sub>4</sub>-FYM @ 2.5 t/ha + *Rhizobium* + PSB gave significantly higher number of root nodules per plant, fresh and dry weight of root nodules.

*Keywords:* Vermicompost, nodule, castor cake, Jeevamrut and dry matter accumulation

#### **1. INTRODUCTION**

Pulses are wonder gift of nature to the living universe and the real gateway of sustainable agriculture. Cowpea is highly responsive to fertilizer application. The dose of fertilizer depends on the initial soil fertility status and moisture availability conditions. Nitrogen plays important role in various metabolic process of the plant growth. Nitrogen is an essential constituent of protein and chlorophyll [1]. The manures are organic in nature, plant or animal origin and contain organic matter in large proportion and plant nutrients in small quantities and used to improve soil productivity by correcting soil physical, chemical and biological properties. Manures contribute to the fertility of the soil by adding organic matter and nutrients, such as nitrogen, that are trapped by bacteria in the soil. To alleviate the problem, included nutrient management is the fine method in the protection of soil fertility and supply of plant vitamins to a foremost stage for maintaining the desired crop productivity through optimization of blessings from all feasible sources of plant nutrients in an incorporated way [2]. There is growing interest in using organic manures as a source of nutrient supply to crop production for long-term soil productivity, ecological stability and reducing the need for chemical fertilizer. Among

the various organic manure sources, FYM and vermicompost are readily available on the market. Integration of inorganic and organic manures not only sustains crop production but also improves soil health and nutrient use efficiency [3]. Organic manures enhances the soil fertility and yield of crops by rendering unviable sources of elemental nitrogen bound, phosphate and decomposed plant residues into available form in order to facilitate the plant to absorb the nutrients [4].

## 2. MATERIALS AND METHODS

A field experiment was conducted at Agronomy Instructional Farm, Chimanbhai Patel College of Agriculture, SardarkrushinagarDantiwada Agricultural University, Sardarkrushinagar to study the “Effect of organic manures and bio-fertilizers on growth and yield of *kharif* cowpea [*Vigna unguiculata* (L.) Walp]” during *kharif* season of 2020-2021. The soil of experimental field was loamy sand in texture with low in organic carbon (0.18 %) and available nitrogen (138.0 kg/ha), medium in available phosphorus (38.6 kg/ha) and high in available potassium (215.2 kg/ha) having pH value of 7.56. The experiment consisting nine treatments *viz.*, T<sub>1</sub>-Absolute control, T<sub>2</sub>-100% RDF (20 kg N + 40 kg P<sub>2</sub>O<sub>5</sub>/ha), T<sub>3</sub>- FYM @ 5.0 t/ha, T<sub>4</sub>-FYM @ 2.5 t/ha + *Rhizobium* + PSB, T<sub>5</sub>-vermicompost @ 2.0 t/ha, T<sub>6</sub>-vermicompost @ 1.0 t/ha + *Rhizobium* + PSB, T<sub>7</sub>-castor cake @ 2.0 t/ha, T<sub>8</sub>-castor cake @ 1.0 t/ha + *Rhizobium* + PSB, T<sub>9</sub>-*GhanJeevamrut* @ 250 kg/ha at sowing time + Seed treatment with *Beejamrut*@ 200 ml/kg seed+ *Jeevamrut* @ 500 lit/ha with irrigation at sowing and 30 DAS were tested in randomized block design with four replications. Cowpea variety GC 4 was used as a test crop.

### Method for preparation of different bio-enhancers

Different bio enhancers used for the application as per treatments were prepared at SDAU, Sardarkrushinagar. The method of preparation are described below.

**1. Bijamrut**-Take 5 kg cow dung + 5 liter cow urine + 50 gm lime and handful soil from the bund of farm + 20 liter water. Preparation method: Take 5 kg local cow dung in cloth and bound it by tape. Hang this in the 20 liter water up to 12 hours. Take 1 liter water and add 50 gm lime in it, let it stable for a night. Then next morning, squeeze this bundle of the cow dung in that water thrice continuously, so that all essence of cow dung is accumulated in that water. Add a handful of soil in that water solution and stir it well. Finally add 5 liter deshi cow urine in that solution and add the lime water and stir it well. Keep this mixture for 12-16 hours. Filter it and use for seed treatment.

**2. Jivamrut**-Take 10 kg cow dung (fresh) + 10 liter cow urine + 2 kg jaggery + 2 kg pulse flour (cow pea) + 1 kg *sajiv* soil + 200 liter water in a container and stirred well. Allowed the mixture to ferment for 7 days under tree shade. The mixture was stirred twice (morning and evening) every day in a

clockwise direction. The container was kept under well ventilated open shed. The mouth of container was tied with thin cotton cloth to enable proper aeration in the container. *Jivamrut* was applied @ 500 lit/ha with irrigation at sowing and 30 DAS as per the treatments.

**3. Ghan Jivamrut-** *Ghanjivamrut* is dry or solid *jivamrut* which can be dried and stored for later use. Take 200 kg of desi cow dung manure and spread uniformly on the ground to form a layer. Now add 20 liter of liquid *jivamrut* on the manure. Mix the mixture well and store it as a heap for 48 hrs in a shade. After 48 hrs, spread the mixture on a clean surface and sundry it. Flip it upside down during the day so that all particles get exposed to sunlight and quickly dries up. Once it is completely dried up break the lumps into powder form with the wooden bat and fill them into jute bags and store it in a cool and dry place.

### **Seed treatment**

The required quantity of healthy cowpea seeds of cultivar GC 4 was inoculated with *rhizobium* and PSB @ 20 ml/kg seeds and required quantity of seeds inoculated with *bijamrut* @ 200 ml/kg seeds as per treatments.

### **Statistical Analysis**

The data recorded for various parameters during the course of investigation were statistically analysed [5] The significance of difference was tested by “F” test at 5 per cent level. The critical difference were calculated when the differences among treatments were found significant under “F” test. In remaining cases, only standard error of mean was worked out.

## **3. RESULTS AND DISCUSSION**

### **3.1 Effect on plant population (per metre row length)**

The data presented in Table 1 showed that the plant population per meter row length at 20 DAS and at the time of harvest were found not significant due to different integrated nutrient management treatments, which indicated that there was no any adverse effect of bio fertilizers and organic sources of nutrients on germination as well as on survival of the cowpea plants.

### **3.2 Effect on growth parameters**

Data on plant height, number of branches per plant, dry matter accumulation, number of root nodules per plant, fresh and dry root nodules weight per plant (mg) and days to physiological maturity are presented in Table 1.

#### **3.2.1 Plant height (cm)**

The data on plant height of cowpea recorded at the time of 30 DAS, 60DAS and at harvest are presented in Table 1 clearly indicated that the plant height was increased progressively and linearly up to the harvest with the advancement of crop growth and also significantly affected by the different

integrated nutrient management treatments. The mean data on plant height presented in Table 1 indicated that different treatments had significant influence on plant height at 30 DAS. Significantly higher plant height (21.58 cm) was recorded by the application of 100% RDF (T<sub>2</sub>), but it was at par with FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>) and vermicompost @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>6</sub>). At 60 DAS and at harvest, plant height was significantly influenced by different integrated nutrient management treatments. Significantly higher plant height at 60 DAS (48.07 cm) and at harvest (52.26 cm) was recorded by the application of castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>), but it was at par with 100% RDF (T<sub>2</sub>), FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>), vermicompost @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>6</sub>) and castor cake @ 2.0 t/ha (T<sub>7</sub>) at the time of 60 DAS and FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>), vermicompost @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>6</sub>) at the time of harvest. Significantly the lower plant height at 30 DAS (17.08 cm), 60 DAS (36.68 cm) and at harvest (44.97 cm) was observed by the absolute control (T<sub>1</sub>). The probable reason might be positive effect of nutrient on growth character due to augment of cell division and cell expansion [6]. The increase in the plant growth in terms of plant height under treatment T<sub>8</sub> (Castor cake @ 1.0 t/ha + *Rhizobium* + PSB) is the combined use of organic manure and bio-fertilizer might be increased the fertilizer use efficiency and also nutrient availability leading to the higher growth of plant in terms of plant height by higher cell multiplication and cell elongation. Moreover, organic manures also contain almost all essential elements in variable quantities, which had synergistic effect with other essential elements for their availability [7].

#### 4.3.2 Number of branches per plant

Maximum number of branches per plant with castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>) might be due to the integrated use of organic manures and bio fertilizers offers more balanced nutrition and favourable soil conditions for better growth of the plant. Moreover, higher availability of nutrients might be reflected on higher number of branches per plant [8,9].

#### 3.2.3 Number of root nodules per plant

An appraisal of data presented in Table 1 indicated that the number of root nodules per plant was significantly influenced by different integrated nutrient management treatments. An application of FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>) to cowpea crop recorded significantly higher number of root nodules per plant (25.00), but it was at par with vermicompost @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>6</sub>), Castor cake @ 2.0 t/ha (T<sub>7</sub>) and Castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>). Significantly less number of root nodules per plant (18.32) was noted under absolute control (T<sub>1</sub>). Improvement in number of root nodules per plant could be attributed due to favorable aeration and moisture regime with the integration of organic manures with bio-fertilizers. Moreover, seed inoculation of bio-fertilizers (*Rhizobium* + PSB) may be provided more conducive environment for better root growth and respiration along with higher soil biological activity. The PSB liquid bio-fertilizer might also have enhanced the availability of phosphorus to plants which must have utilized in greater root development and nodulation [10].

#### 3.2.4 Dry matter accumulation per plant (g)

The mean data on dry matter accumulation of cowpea presented in Table 1 indicated that different treatments had significant influence on dry matter accumulation at 30 DAS. Treatment T<sub>2</sub> (100% RDF)

registered significantly higher dry matter per plant (37.30 g) which was at par with FYM @ 5.0 t/ha(T<sub>3</sub>), FYM @ 2.5 t/ha + *Rhizobium* + PSB(T<sub>4</sub>), vermicompost @ 1.0 t/ha + *Rhizobium* + PSB(T<sub>6</sub>), castor cake @ 2.0 t/ha(T<sub>7</sub>) and castor cake @ 1.0 t/ha + *Rhizobium* + PSB(T<sub>8</sub>). Data presented in Table 1. Showed that significantly higher dry matter per plant at 60 DAS (63.44 g) and at harvest (92.02 g) was observed with treatment T<sub>8</sub> (castor cake @ 1.0 t/ha + *Rhizobium* + PSB) which was at par with treatment 100% RDF(T<sub>2</sub>), FYM @ 2.5 t/ha + *Rhizobium* + PSB(T<sub>4</sub>), vermicompost @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>6</sub>) and castor cake @ 2.0 t/ha(T<sub>7</sub>). Significantly the lower dry matter accumulation at 30 DAS (30.43 g), 60 DAS (47.85 g) and at harvest (60.14 g) was observed by the absolute control (T<sub>1</sub>). Adequate major nutrients might have helped in harvesting of solar energy as reflected by increased dry matter accumulation. The beneficial effect of organic manures and bio-fertilizers in growth attributes are close agreement with the findings reported by Pargiet *al.* [11].

### **3.2.5 Fresh and dry root nodules weight per plant (mg) at 45 DAS**

Results presented in Table 1 indicated that the fresh and dry root nodules weight per plant was significantly influenced by different integrated nutrient management treatments. Significantly maximum fresh nodules weight per plant (69.80 mg) and dry nodules weight per plant (26.77 mg) was recorded with treatment T<sub>4</sub> (FYM @ 2.5 t/ha + *Rhizobium* + PSB), but it was at par with castor cake @ 1.0 t/ha + *Rhizobium* + PSB(T<sub>8</sub>). Significantly the minimum fresh nodules weight per plant (53.71 mg) and dry nodules weight per plant (15.99 mg) was registered under the treatment of absolute control (T<sub>1</sub>). Improvement in fresh and dry root nodules weight per plant could be attributed due to favorable aeration and moisture regime with the integration of organic manures with bio-fertilizers. Moreover, seed inoculation of bio-fertilizers (*Rhizobium* + PSB) may be provided more conducive environment for better root growth and respiration along with higher soil biological activity. The positive response of organic manures and bio fertilizer to increase the nodulation might be due to the organic matter provides energy for nodulation. The PSB liquid bio-fertilizer might also have enhanced the availability of phosphorus to plants which must have utilized in greater root development and nodulation [12].

### **3.3 Effect on yield and yield attributes**

#### **3.3.1 Number of pods per plant**

The set of data furnished in Table 1 confirms that the number of pods per plant was influenced significantly by different treatments. Significantly the more number of pods per plant (12.80) was recorded under the application of castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>). Treatment 100% RDF(T<sub>2</sub>) and FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>) remained at par with treatment T<sub>8</sub>. The increment in number of pods per plant under T<sub>8</sub> and T<sub>4</sub> was to the tune of 55.15 and 51.51 per cent higher than treatment T<sub>1</sub> (absolute control), respectively. Significantly the less number of pods per plant (8.25) was recorded under the treatment T<sub>1</sub> (absolute control). The data in Table 1 displayed that the application of treatment T<sub>8</sub> of castor cake @ 1.0 t/ha + *Rhizobium* + PSB on number of pods per plant was found significant over rest of the treatments except treatment 100% RDF (T<sub>2</sub>) and FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>). More number of pods/plant might be due to more survival of flowers under high supply of photosynthates under treatment castor cake @ 1.0 t/ha + *Rhizobium* + PSB(T<sub>8</sub>) as compared to other treatments. Higher number of pods per plant might be owing to better vegetative growth and photosynthetic activity ascribed by larger uptake of nutrients and water from

the soil by well-developed root system and profuse nodulation which caused an increase in nitrogen supply from symbiotically fixed nitrogen in root nodules [11].

### 3.3.2 Number of seeds per pod

The data presented in Table 1 disclosed that the number of seeds per pod was significantly influenced by application of different treatments. Significantly the higher number of seeds per pod (10.39) was obtained under treatment T<sub>8</sub> i.e. application of castor cake @ 1.0 t/ha + *Rhizobium* + PSB which remained at par with treatment FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>), 100% RDF (T<sub>2</sub>), vermicompost @ 2.0 t/ha (T<sub>5</sub>), vermicompost @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>6</sub>) and castor cake @ 2.0 t/ha (T<sub>7</sub>). Significantly the lower number of seeds per pod (8.61) was gained under absolute control (T<sub>1</sub>). The increment in number of seeds per pod under T<sub>8</sub> and T<sub>4</sub> was to the tune of 20.67 and 18.93 per cent higher than treatment T<sub>1</sub> (absolute control), respectively. The highest number of seeds per pod was obtained with application of castor cake @ 1.0 t/ha + *Rhizobium* + PSB probably due to the fact that supplying of ample amount of nutrients in form of organic manure and bio-fertilizer must have increased carbohydrate accumulation and their remobilization to reproductive parts of the plant, being closest sink and hence, resulted into increased flowering, fruiting and seed formation and thus more number of seeds per pod [13].

### 3.3.3 Length of pod (cm)

The results from Table 1 revealed that different treatments including organic manure and bio-fertilizer significantly influenced on length of pod of cowpea. Significantly higher length of pod (11.56 cm) was obtained through the application of castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>). Treatment 100% RDF (T<sub>2</sub>), FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>), vermicompost @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>6</sub>) and castor cake @ 2.0 t/ha (T<sub>7</sub>) remained at par with treatment castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>) and the lower value of pod length (9.78 cm) was obtained by absolute control. The length of pod measured under castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>) and FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>) was 18.20 and 17.58 per cent higher than treatment T<sub>1</sub> (absolute control), respectively. The length of cowpea pod was significantly influenced by different organic manures as well as bio fertilizers (Table 2. and Fig. 1). Lengthier pods were observed with application of castor cake @ 1.0 t/ha + *Rhizobium* + PSB probably due to the fact that increased supply of N and P and their higher uptake by plants might have stimulated the rate of various physiological processes in plant and resulted in increased pod length. The increase in pod length might be due to better partitioning of assimilates from source to sink. Resultant utilization of accumulated photosynthates influenced the growth and development of yield attributes [14].

### 3.3.4 Seed yield per plant (g)

An appraisal of data presented in Table 2 clearly indicated that seed yield per plant of *kharif* cowpea was differed significantly due to various integrated nutrient management treatments. Significantly higher seed yield per plant (8.73 g) was obtained under treatment castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>), but it remained at par with treatment of 100% RDF (T<sub>2</sub>), FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>) and vermicompost @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>6</sub>). The magnitude of increase in seed yield per plant due to castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>) and FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>) over T<sub>1</sub> (absolute control) was to the tune of 40.80 and 38.87 per cent,

respectively. Significantly the lower seed yield per plant (6.20 g) was noted under treatment absolute control (T<sub>1</sub>). Castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>) recorded higher seed yield per plant than other treatments. The seed yield per plant was higher due to higher pod length, seeds per pod, number of pods per plant and also higher test weight due to better growth of plant. This result is in conformity with the findings of Vikrant *et al.*[15].

### 3.3.5 Seed index (g)

The data presented in Table 2 disclosed that seed index was not varied significantly due to different treatments. However, marginally higher seed index was observed in treatment castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>) (9.98 g) than rest of treatments.

### 3.3.6 Seed yield (kg/ha)

A close examination of result indicated that application of organic manure and bio-fertilizer manifest significant influence on seed yield. Significantly the higher seed yield (1109kg/ha) was recorded under the treatment T<sub>8</sub> (castor cake @ 1.0 t/ha + *Rhizobium* + PSB) which was at par with treatment 100% RDF (T<sub>2</sub>), FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>) and vermicompost @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>6</sub>). Whereas, absolute control (T<sub>1</sub>) gave the lower seed yield (707kg/ha) among the treatments. The magnitude of increase in seed yield of cowpea due to castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>) and FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>) over T<sub>1</sub> (absolute control) was the extent of 56.85 and 49.08 per cent and over T<sub>2</sub> (100% RDF) was the extent of 6.94 and 1.64 per cent, respectively. Seed yield was significantly higher with treatment castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>) might be due to the fact that combined application of organic manure and Bio-fertilizer must have produced excess of assimilates which are first stored in leaves and later translocate into seeds at the time of senescence, which might have ultimately led to higher seed yield. Second reason for higher seed yield might be due to increase in photosynthetic activity of plant and root system and thus enabled plant to extract more water and nutrients from the soil depth, resulting into better development of plant growth, yield attributes and ultimately the higher seed yield. Third probable reason for increased seed yield under the INM may be the higher number of pods per plant and number of seeds per pod which gradually increased the seed yield under this treatment [16].

### 3.3.7 Stover yield (kg/ha)

A close surveillance of data presented in Table 2 showed that application of organic manure and bio-fertilizer manifest significant influence on Stover yield. Treatment T<sub>8</sub> (castor cake @ 1.0 t/ha + *Rhizobium* + PSB) being at par with treatment 100% RDF (T<sub>2</sub>), FYM @ 5.0 t/ha (T<sub>3</sub>), FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>) and vermicompost @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>6</sub>) recorded significantly the higher stover yield (1986kg/ha). Whereas, absolute control (T<sub>1</sub>) produced lower stover yield (1367kg/ha) among the different treatments. The magnitude of increase in stover yield due to castor cake @ 1.0 t/ha + *Rhizobium* + PSB (T<sub>8</sub>) and FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>) over T<sub>1</sub> (absolute control) was to the tune of 45.28 and 34.60 per cent, respectively. Treatment T<sub>8</sub> i.e. application of castor cake @ 1.0 t/ha + *Rhizobium* + PSB gave the highest Stover yield probably because of the fact that under this treatment the synthesis of photosynthates is more owing to adequate supply of nitrogen and phosphorus which increased the photosynthesis activity and

production of more biomass, which ultimately resulted in higher yield of stover. Similar results in cowpea were given by [17].

#### **4.5 Effect on quality parameters**

##### **4.5.1 Protein content in seed (%)**

The set of data concerning to the protein content in seeds of cowpea which are provided in Table 2 indicates that the different treatments succeed in having their significant influence on protein content. Significantly higher protein content (22.33 %) was gained through the treatment T<sub>8</sub> (Castor cake @ 1.0 t/ha + *Rhizobium* + PSB), which was at par with treatment 100% RDF (T<sub>2</sub>), FYM @ 2.5 t/ha + *Rhizobium* + PSB (T<sub>4</sub>) and *GhanJeevamrut* @ 250 kg/ha at sowing time + Seed treatment with *Beejamrut* @ 200 ml/kg seed + *Jeevamrut* @ 500 lit/ha with irrigation at sowing and 30 DAS (T<sub>9</sub>). Among all the treatments observed the lower protein content (18.41 %) was obtained under treatment T<sub>1</sub> (absolute control). As a whole, treatment T<sub>8</sub> (castor cake @ 1.0 t/ha + *Rhizobium* + PSB) and T<sub>4</sub> (FYM @ 2.5 t/ha + *Rhizobium* + PSB) were correspondingly increased of protein content was to the tune of 21.29 and 16.89 per cent over treatment T<sub>1</sub> (absolute control), respectively. Increase in protein content in seed might be due to the fact that higher nitrogen content in seed is directly associated to higher availability of nitrogen to plants. Higher nitrogen in seeds are directly responsible for higher protein content because it is a primary component of amino acid which constitutes the basis of protein content. Another reason for increasing the protein content might be due to fact that cowpea is a leguminous crop and the application of bio-fertilizers must have activated the microbial population responsible for root nodulation and efficient nodulation which must have enhanced nitrogen fixation by the plant and ultimately increased the protein content [18].

#### **4. CONCLUSION**

Based on field experimentation, it is concluded that higher yield from *kharif* cowpea could be obtained by applying either castor cake @ 1.0 t/ha or FYM @ 2.5 t/ha along with *Rhizobium* + PSB in loamy sand of North Gujarat Agro-climatic conditions.

#### **REFERENCES**

1. Meena, L. R. and Chand, R. Response of fodder cowpea to varying levels of nitrogen and phosphorus under rainfed conditions of Rajasthan. *Indian Journal of Small Ruminants*, 2014;20(2): 121-123.
2. Jagdale A, Dhamak A, Pagar B, Wagh P. Effect of different organic formulations on growth and yield of soybean. *International Journal of Chemical Studies*. 2020; 8(4): 1634-1638.
3. Gowthamchand NJ, Ganpathi, Soumya TM. Effect of bulky manures and fermented liquid organics on growth, yield, nutrient uptake and economics of French bean (*Phaseolus vulgaris* L.) under rainfed condition. *International Journal of Agriculture, Environment and Biotechnology*. 2019; 12(4):361-368.
4. Verma TS. Research Bulletin No. 2, AICRP-LTFE, IISS, Bhopal (MP) and Department of Soils, CSK HPKV, Palampur (HP). 2005;1-88
5. Panse, V.G. and Sukhatme, P.V. Statistical methods for agricultural workers, ICAR pub. New Delhi. 1985; pp. 361.

6. Khandelwal, R., Choudhary, S.K., Khangarot S.S., Jat, M.K. and Singh, P. Effect of inorganic and biofertilizers on productivity and nutrients uptake in cowpea [*Vigna unguiculata*(L.)Walp]. *Legume Research*, 2012; 35(3): 235-238.
7. Kulkarni, M.V.; Patel K.C.; Patil D.D. and Madhuri Pathak. Effect of organic and inorganic fertilizers on yield and yield attributes of groundnut and wheat. *International Journal of Chemical Studies*.2018; 6(2): 87-90.
8. Birla J., Patel, B.M., Patel, P.M., Tamboli, Y.A. and Patil D. Yield and quality of cowpea [*Vigna unguiculata* (L.)Walp] as influenced by organic sources of nitrogen. *Legume Research*,2017; 41(6): 899-902
9. Kalegore, N. K., Gavhane, M. A., Bhusari, S. A., Kastle, S. V. and Dhamane, R. S. Response of cowpea [*Vigna unguiculata* (L.)Walp.] to inorganic and biofertilizers. *International Journal of Economic Plants*, 2018; 5(4):167-169.
10. Lyngdoh, C., Bahadur, V., David, A.A., Prasad, V.M., and Jamir, T. Effect of manures, organic supplements and bio-fertilizers on growth and yield of cowpea (*Vigna unguiculata* L. Walp). *International Journal of Current Microbiology and Applied Sciences*.2017; 6(8): 1029-1036.
11. Pargi, K. L., Leva, R. L., Vaghasiya, H. Y. and Patel H. A. Integrated nutrient management in summer cowpea [*Vigna unguiculata* (L.) Walp.] under south Gujarat condition. *International Journal of Current Microbiology and Applied Sciences*, 2018;7(9): 1513-1522.
12. Khandelwal, R., Choudhary, S.K., Khangarot S.S., Jat, M.K. and Singh, P. Effect of inorganic and biofertilizers on productivity and nutrients uptake in cowpea [*Vigna unguiculata*(L.)Walp]. *Legume Research*, 2012; 35(3): 235-238.
13. Gupta Shivangini. Effect of Nitrogen, Phosphorus and bio-fertilizer on green pod yield of summer cowpea (*Vigna unguiculata*L. Walp) under middle Gujarat condition, M.Sc. (Agri.) Thesis (Unpublished) submitted to Anand Agricultural University, Anand, Gujarat.
14. Chauhan, J., Paithankar D.H, Khichi, P., Ramteke, V., Jogdhande, S., and Manohar, M. Effect of various sources of nutrients including organic, inorganic, bio- fertilizer and its combination on growth and protein content of cowpea (*Vigna unguiculata* L. Walp). *Research Journal of Agricultural Sciences*.2016; 7(2): 256-259.
15. Vikrant, V.R., Singh, H., Malik, C.V.S. and Singh, B.P. Influence of farm yard manure and phosphorus application on grain yield and economics of cowpea. *Crop Research*, 2005; 29(1): 87-90.
16. Joshi, J. R., Patel, V. M., Barad, H. L., Macwan, S. M. and Javid, Ehsas. Effect of land configuration and fertilizer management practices on growth, yield and yield attributes and economics of summer cowpea [*Vigna unguiculata*(L.) Walp.] under South Gujarat condition. *International Journal of current Microbiology and Applied Sciences*, 2018; 7(1):1148-1155.
17. Dekhane, S.S., Khafi, H.R., Raj, A.D., and Parmar, R.M. Effect of bio fertilizer and fertility levels on yield, protein content and nutrient uptake of cowpea [*Vigna unguiculata* (L.) Walp.]. *Legume Research*, 2011; 34(1):51-54.
18. Joshi, D.,Gediya, K.M., Patel, J.S., Birari, M.M., and Gupta, S.Effect of organic manures on growth and yield of summer cowpea [*Vigna unguiculata*(L.) Walp] under middle Gujarat conditions. *Agricultural Science Digest*, 2016; 36(2): 134-137.

**Table: 1. Effect of different integrated nutrient sources on plant population, plant height, number of branches, root nodules, dry matter and growth parameters of *kharif* cowpea.**

Treatments	Plant population (per meter row length)		Plant height (cm)			Number of branches per plant	Number of root nodules /plant at 45D AS	Dry matter/plant(g)			Fresh root nodules weight per plant (mg)	Dry root nodules weight per plant (mg)	Number of pods per plant	Number of seeds per pod	Length of pod (cm)
	20 DAS	At harvest	30 DAS	60 DAS	At harvest			30 DAS	60 DAS	At harvest					
T <sub>1</sub> - Absolute control	8.51	7.89	17.08	36.68	44.97	3.86	18.32	30.43	47.85	60.14	53.71	15.99	8.25	8.61	9.78
T <sub>2</sub> -100% RDF	8.77	8.13	21.58	44.18	46.97	5.36	19.82	37.30	59.30	83.32	62.20	21.67	11.45	10.12	11.28
T <sub>3</sub> -FYM @ 5.0 t/ha	8.60	8.06	19.08	42.67	45.37	4.89	21.19	33.60	56.74	80.51	63.83	22.76	10.65	9.07	10.19
T <sub>4</sub> -FYM @ 2.5 t/ha + <i>Rhizobium</i> + PSB	8.88	8.24	19.82	47.53	51.73	6.17	25.00	36.90	61.30	86.17	69.80	26.77	12.50	10.24	11.50
T <sub>5</sub> - Vermicompost @ 2.0 t/ha	8.58	8.16	18.88	42.27	45.26	4.78	20.81	33.10	54.10	76.67	62.47	21.86	9.70	9.64	10.20
T <sub>6</sub> - Vermicompost @ 1.0 t/ha + <i>Rhizobium</i> + PSB	8.78	8.14	19.77	47.07	50.40	5.52	24.43	36.70	59.40	86.07	66.26	24.40	11.25	10.05	11.46
T <sub>7</sub> - Castor cake @ 2.0 t/ha	8.70	8.09	19.18	44.23	46.86	5.23	21.81	35.60	58.42	84.00	63.97	22.87	10.75	9.78	10.86
T <sub>8</sub> - Castor cake @ 1.0 t/ha + <i>Rhizobium</i> + PSB	8.98	8.24	19.08	48.07	52.26	6.23	24.46	33.30	63.44	92.02	68.96	26.21	12.80	10.39	11.56
T <sub>9</sub> - <i>GhanJeevamrut</i> @ 250 kg/ha at sowing time + Seed treatment with <i>Beejamrut</i> @ 200 ml/kg seed+ <i>Jeevamrut</i>	8.53	7.90	18.79	41.98	45.17	4.49	20.70	32.23	53.40	66.04	62.33	21.77	9.30	9.29	9.99

<i>t</i> @ 500 lit/ha with irrigation at sowing and 30 DAS															
S.Em. ±	0.21	0.16	0.67	1.57	1.65	0.25	1.15	1.42	1.76	3.14	1.55	0.74	0.47	0.36	0.33
C.D. at 5 %	NS	NS	1.97	4.59	4.82	0.73	3.34	4.15	5.13	9.16	4.53	2.17	1.36	1.05	0.98
C.V. %	4.87	4.08	7.00	7.17	6.93	9.67	10.49	8.27	6.15	7.90	4.87	6.56	8.70	7.40	6.22

**Table: 2. Yield parameters, Harvest index and protein content of *kharif* cowpea as influenced by different integrated nutrient management treatments.**

Treatments	Seed yield per plant (g)	Seed yield (kg/ha)	Stover yield (kg/ha)	Harvest index (%)	Days to physiological maturity	Protein content (%)
T <sub>1</sub> -Absolute control	6.20	707	1367	34.16	78.25	18.41
T <sub>2</sub> -100% RDF	8.15	1037	1825	36.20	82.50	20.73
T <sub>3</sub> -FYM @ 5.0 t/ha	7.24	945	1733	35.36	82.75	20.11
T <sub>4</sub> -FYM @ 2.5 t/ha + <i>Rhizobium</i> + PSB	8.61	1054	1840	36.34	84.75	21.52
T <sub>5</sub> -Vermicompost @ 2.0 t/ha	6.91	901	1699	34.75	82.25	20.41
T <sub>6</sub> -Vermicompost @ 1.0 t/ha + <i>Rhizobium</i> + PSB	8.15	1033	1813	36.36	84.75	20.66
T <sub>7</sub> -Castor cake @ 2.0 t/ha	7.57	958	1721	35.84	82.75	20.30
T <sub>8</sub> -Castor cake @ 1.0 t/ha + <i>Rhizobium</i> + PSB	8.73	1109	1986	35.79	84.50	22.33
T <sub>9</sub> - <i>GhanJeevamrut</i> @ 250 kg/ha at sowing time + Seed treatment with <i>Beejamrut</i> @ 200 ml/kg seed+ <i>Jeevamrut</i> @ 500 lit/ha with irrigation at sowing and 30 DAS	6.80	891	1596	35.93	80.50	21.11
S.Em. ±	0.31	50	88	1.38	2.22	0.56
C.D. at 5 %	0.90	146	257	NS	NS	1.65
C.V. %	8.08	10.41	10.18	7.72	5.38	5.48