

Original Research Article

Risk Assessment and Distribution of Heavy Metals in Water and Sediments along Ibelebiri Axis of Kolo Creek, Ogbia Local Government Area, Niger Delta Region of Nigeria.

ABSTRACT

This study was conducted to investigate the risk and distribution of heavy metals (Zn, Ni, Cr, Cu, Pb, Co, and Cd) in water and sediments along Ibelebiri axis of Kolo creek in Ogbia L.G.A., Niger Delta region of Nigeria. Water and sediment samples were also collected from a control site. The samples were digested using acid mixture and the filtrates obtained were analysed using atomic absorption spectrophotometer fitted with inductive coupled plasma atomic emission spectroscopy (ICP-AES). Generally, heavy metal concentrations showed a decreasing trend of Cu > Zn > Pb > Cr > Cd > Co > Ni for water samples. The mean concentration of these metals: Ni, Cr, Zn and Cu recorded in this study is below international recommended limits. But, Pb, Cd levels in locations LC1, LC3 and LC4 are above the international recommended limits. The heavy metals in the sediments showed a decreasing trend of Zn > Pb > Cu > Co > Cr > Ni > Cd. The concentrations of these metals - Cd, Pb, Ni, Cr, Cu and Zn are below the permissible limit set by DPR. However, Co values at locations LC3 and LC4 are slightly above the DPR target value of 20 mg/kg for sediments. Using the geoaccumulation index (Igeo) classification, the sediments across the four locations are uncontaminated with Cd, Cr, Cu, Zn, Co and Ni. But, the sediments in locations LC1, LC3 and LC4 are moderately contaminated with Pb. Similarly, the contamination factor (CF) classification, revealed low contamination of Cd, Cr, Cu, Zn, Ni and Co across the four locations; moderate contamination of Pb in the four locations. Finally, principal component analysis (PCA) identified two major factors: uncontrolled combustion processes such as artisanal refinery activities and gas flaring from nearby flow stations, may have contributed to the pollution of the water and sediments in the study area.

Key words: Heavy metals, risk, pollution, contamination, accumulation.

Comment [OM1]: What warranted this research? Full justification is needed.

Comment [OM2]: Delete

Comment [OM3]: was

Comment [OM4]: in full

Comment [OM5]: Recommendations?

Comment [OM6]: arrange in alphabetical order

1. INTRODUCTION

Rivers are a crucial conduit for the entry of sediment, salt, and chemical components into lakes and seas in the hydrologic cycle of the planet (1). Every year, a significant amount of heavy metals are extracted from the earth's crust in order to advance the economy and satisfy daily human needs (2). Heavy metals are released into rivers during the mining and production process through atmospheric sedimentation or industrial wastewaters. The bulk is concentrated in sediments, and only a minor portion is dissolved in water bodies. The foundation of food webs and an essential component of aquatic environments is sediments. However, they also act as pollution sinks (3). Heavy metals are accepted and transported by rivers, and river sediment not only acts as a significant sink and transporter of heavy metals but also as a possible source of secondary pollution, which can also indicate the level of contamination (4). Additionally, heavy metal pollution often has a long half-life, a

wide distribution, and is easily accumulated and stabilized in the environment, endangering human health either directly or through the food chain (5,6). One of the greatest environmental issues in the world today is heavy metal contamination in rivers brought on by social, economic, and industrial activity (7), which has caught the attention of academics. A study (8) was carried out on the effect of refinery and petrochemical effluent on water quality of Ubeji Creek Warri, Southern Nigeria. The study showed obvious pollution of the creek but that there has been an improvement in the treatment of Warri Refinery and Petrochemical effluent before it is discharged. However, pollution arising from uncontrollable combustion processes from artisanal refineries, gas flaring and biomass which releases atmospheric particulates especially black carbon has been a major concern to environmentalist in recent times. These particulates are easily washed down and get into the sink. The buildup of heavy metals in the sediment poses a long-term threat to the aquatic habitats (4).

Comment [OM7]: scientific community

In Bayelsa State, due to poor management of municipal and industrial effluents, water bodies including the Ibelebiri stream which discharges into Kolo creek have been adversely affected. The unsustainable environmental practices, socio-economic issues have worsened spatially in the last two decades. Consequently, there has been increase in water-borne diseases, a decline in water value, various ecological and human health risks. A study (9) was carried out on the heavy metal distribution and assessment of ecological risk in surface waters and sediments within the flow stations in Kolo creek, Nigeria. Severally other studies investigated the concentration of heavy metals in water and sediments of Kolo creek and some aquatic animals found in the sediments (10,11,12). The attention of the creek is obviously because of the presence of Shell Petroleum flow station at Imirigi which is one the communities through which the Kolo creek traverses. Previous studies on the Nemberi river have focused on the river water chemistry and physicochemical properties in the river water (13,14) and few studies on seasonal and spatial distribution of heavy metals (15,16). However, in the scope of our knowledge, there has hardly been any detailed studies on the heavy metal contamination of Kolo creek at Ibelebiri watershed which traverses through other communities including Imirigi where Shell Petroleum flow station is located. Therefore, this study is aimed at investigating other sources of heavy metals in the creek beside the flow station or pollution due to oil and gas production.

Comment [OM8]: state the name of country

Comment [OM9]: in Nigeria

2. MATERIALS AND METHODS

2.1 Study Area

Ibelebiri is a community in Ogbia Local Government Area of Bayelsa State. The climate of the area is characterized by a relatively dry season (November – March) period with occasional precipitation and wet season (April – October). The annual rainfall varies between 2500 – 3000mm. Mean temperatures are as high as 32°C with humidity of about 81% in the dry season and about 92% in the wet season (17). The inhabitants are predominantly farmers and fishermen. Some are engaged in petty trading, and some are civil servants. The major source of water is the creek; rainwater stored in tanks and a few boreholes owned by some individuals.

Comment [OM10]: Include GPS points




Fig. 1: Map of Study Area showing sampling locations (black dots)
(source: www.mapcarta.com/17024300)

Comment [OM11]: Get a better Map from Google maps with proper coordinates

2.2 Sample Collection and Preparation

Periodic seasonal sampling (November and December 2021 – January 2022), (March – May 2022) and (June – August 2022) was done using standard procedure. Watersamples were collected from Kolo creek at Ibelebiri community axis- (location 1: before Ibelebiri community, location 2: within Ibelebiri community, location 3: after Ibelebiri community and location 4: used as control- at approximately 500m apart) into pre-cleaned 1L plastic containers, acidified with few drops of nitric acid, and stored in a refrigerator until analysis. Sediment samples were collected using plastic grabs from the same locations where water samples were collected from, and transferred into well-labeled, pre-cleaned polythene bags. Samples were then transported to the laboratory and kept in a

Comment [OM12]: You must state how the stations were established. What warranted the selection of these stations?

Comment [OM13]: What % or molarity?

Comment [OM14]: Pre-washed

Comment [OM15]: conveyed

refrigerator at $-4\text{ }^{\circ}\text{C}$ before analysis. Similarly, a sample of water and its sediment were also collected from the university town of Otuoke, presumably a more 'pristine' environment to serve as the control. The sediment samples were freeze-dried, crushed in an agate mortar, passed through a nylon 100-mesh sieve ($<0.15\text{ mm}$) and stored in pre-cleaned polythene bags at $4\text{ }^{\circ}\text{C}$ in the dark before analysis.

Comment [OM16]: References needed

2.3 Digestion of samples

2.3.1 Water samples

100 ml of water samples was placed in a 250 ml conical flask, 10 ml of 1M HNO_3 was added, placed in a digestion block and heated at 150°C until the mixture reduced to about 25 ml. The mixture was removed from the block, cooled, filtered using a Whatman No. 40 filter paper, transferred to a 100 mL volumetric flask and made up to mark with 1M HNO_3 (18).

2.3.2 Digestion of sediment samples

Wet digestion of sediment samples was carried out as follows: 2.00 g sample of sediment sample was placed in 50 mL conical flasks. 10 mL of aqua regia (1:3 HNO_3 :HCl) mixture was added, followed by 5 ml of perchloric acid and covered with a dish. This mixture was then placed in a digestion block and heated at 150°C until the acid mixture reduced to about 5 ml. The mixture was removed from the block, cooled, filtered using a Whatman No. 40 filter paper, transferred to a 100 mL volumetric flask and made up to mark with 1 M HNO_3 (18).

The filtrates obtained from digestion of water and sediment were analysed using an Inductively Coupled Plasma-Atomic Emission Spectroscopy (ICP-AES).

Comment [OM17]: What were the units for quantification of metals?

2.4 Ecological Risk Assessment of Heavy Metals Pollution in Sediment

Comment [OM18]: Delete

2.4.1 Geo-accumulation Index (I_{geo})

The I_{geo} has been applied widely to assess soil and sediment contamination, which is defined by the following equation (19):

$$I_{geo} = \log_2 \left(\frac{C_n}{1.5B_n} \right)$$

Where C_n is the measured concentration of heavy metal (n) assayed in sediment samples and B_n is the geochemical background value of heavy metal (n). The value of 1.5 is introduced to minimize the possible variations in the background values due to lithogenic effects.

Comment [OM19]: factor

A scale of I_{geo} indicates differences in the contamination level where it denotes uncontaminated sediments when $I_{geo} \leq 0$ (class 1); uncontaminated to moderately contaminated when $0 < I_{geo} \leq 1$ (class 2); moderately contaminated when $1 < I_{geo} \leq 2$ (class 3); moderately to heavily contaminated when $2 < I_{geo} \leq 3$ (class 4); heavily contaminated when $3 < I_{geo} \leq 4$ (class 5); heavily to extremely contaminated when $4 < I_{geo} \leq 5$ (class 6); and extremely contaminated when $I_{geo} > 5$ (class 7).

2.4.2 Contamination Factor (CF)

The CF is the ratio of the measured concentration of a given metal in the sediment to its background value, which has been classified into four grades as low contamination ($CF < 1$), moderate contamination ($1 \leq CF < 3$), considerable contamination ($3 \leq CF < 6$), and very high contamination ($CF \geq 6$) (20).

2.4.3 Pollution Load Index (PLI)

The PLI of a single site is the n th root of n number of multiplied together Contamination factor (CF) values. The CF is the ratio obtained by dividing the concentration of each metal in the sediment by the baseline or background value.

$$PLI = (CF_1 \times CF_2 \times CF_3 \times \dots \times CF_n)^{1/n}$$

Site indices can be treated in the same way to give a zone or area index. Therefore, PLI for a zone is the n th root of n number of multiplied together PLI values. A PLI value of zero indicates perfection, a value of one indicates the presence of only baseline levels of pollutants, and values above one would indicate progressive deterioration of the site and estuarine quality (21). $PLI > 1$ suggests that pollution exists; otherwise, there is no pollution.

2.5 Correlation Matrix and Factor Analysis

Correlation matrix and principal component analysis was performed on the obtained data using SPSS package version 20.0. Only values which is 0.5 and above were retained. Also, factors with eigen values equal or greater than unity was considered.

2.6 STATISTICAL ANALYSIS

Statistical analysis was carried out using SPSS package version 20.0.

Comment [OM20]: Before this section, you must state the experimental design applied.

Comment [OM21]: Not the best for such studies! GENSTAT will give you better results.

3. RESULTS AND DISCUSSION

3.1 Heavy Metals Concentration in Water

The result of seasonal average of heavy metals (Ni, Cr, Cu, Pb, Zn, Co, Cd) concentration collected from water samples from the Kolo creek is presented in Table 1. The heavy metal with the highest concentration is Cu with a mean value of 0.283 ± 0.064 mg/l at LC1. The range of Cu concentration in LC1 is from 0.210 to 0.330 mg/l. Contrarily, the lowest measured concentration is that of Ni at locations LC3 and LC4, with mean values of 0.001 ± 0.000 mg/l and 0.001 ± 0.000 mg/l respectively. The range of Ni concentrations at locations LC3 is from 0.000 to 0.001 mg/l and in LC4, from 0.000 to 0.001 mg/l respectively. The table also showed that some heavy metals have significant differences in distribution and concentrations across different locations. For example, the mean value of Cu at LC1 is 0.283 ± 0.064 mg/l, which is significantly higher than the mean value of 0.016 ± 0.007 mg/l measured at LC4. The same could be observed in Cr at LC1, with mean value of 0.005 ± 0.004 mg/l significantly higher than the 0.002 ± 0.001 at LC4. This probably suggests that there could be some factors that affect the distribution of heavy metals in the Kolo creek. The heavy metals generally, showed a decreasing trend in concentration: Cu > Zn > Pb > Cr > Cd > Co > Ni. The mean concentration of these metals: Ni, Cr, Zn and Cu recorded in this study are below international recommended limits (22,23). However, Pb and Cd concentrations in locations LC1, LC3 and LC4 recorded in this study, may have exceeded the international recommended limits (22,23). This suggests a potential health risk of lead poisoning in the study area. Elevated level of Pb can cause adverse effect in aquatic and human lives in the area. Chronic exposure to lead at a very low level could lead to neurological and reproductive problems (24). Exposure to lead (Pb) causes loss of memory, nausea, insomnia, anorexia, and weakness of joints, irritation and could make tumour to arise (25,26). There is also risk of cadmium poisoning in the study area. The mean concentration of Cd obtained in this study is well below the range of 0.01 to 0.03 mg/l obtained in Bomadi Creek in the Niger Delta of Nigeria (27). The mean concentration of Cd, Pb, Zn and Cu obtained in this study are below the range obtained in river Benue (28). Again, the mean concentration of Zn, Cd, Ni and Cr obtained in this study are below the values, but Cu and Pb are comparable with the values obtained in river Ureje, Ado Ekiti (29). The level of cadmium, lead and nickel in this study are greater than the value determined in a study by Uzoekwe and Aigberua (9) in the surface waters within the flow stations in Kolo Creek, Nigeria. Furthermore, zinc (Zn) concentrations from this study of Kolo creek at Iberiberi axis is comparable with the value range observed in the Orashi river (30).

Comment [OM22]: It is best that you report using ranges at each station to make the work more presentable .

Comment [OM23]: Information on significant differences must be included and interpreted.

Table 1: Heavy metals concentration (mg/l) in Kolo creek at Ibelebiri community

Location	Ni	Cr	Cu	Pb	Zn	Co	Cd
LC1	0.002 ^a ±0.002 (0.000-0.003)	0.005 ^a ±0.004 (0.001-0.008)	0.283 ^a ±0.064 (0.210-0.330)	0.020 ^b ±0.000 (0.019-0.022)	0.044 ^a ±0.005 (0.039-0.048)	0.002 ^a ±0.001 (0.001-0.002)	0.004 ^a ±0.006 (0.001-0.011)
LC2	0.002 ^a ±0.001 (0.001-0.002)	0.002 ^a ±0.002 (0.000-0.004)	0.026 ^b ±0.002 (0.025-0.028)	0.020 ^b ±0.002 (0.018-0.021)	0.038 ^b ±0.002 (0.036-0.040)	0.002 ^a ±0.000 (0.002-0.002)	0.003 ^a ±0.004 (0.001-0.008)
LC3	0.001 ^a ±0.000 (0.000-0.001)	0.002 ^a ±0.001 (0.001-0.003)	0.015 ^b ±0.001 (0.010-0.022)	0.027 ^a ±0.005 (0.022-0.032)	0.040 ^{ab} ±0.003 (0.038-0.043)	0.001 ^a ±0.001 (0.001-0.002)	0.006 ^a ±0.005 (0.001-0.010)
LC4	0.001 ^a ±0.000 (0.000-0.001)	0.002 ^a ±0.001 (0.001-0.002)	0.016 ^b ±0.007 (0.011-0.024)	0.025 ^{ab} ±0.003 (0.022-0.028)	0.040 ^{ab} ±0.002 (0.038-0.041)	0.001 ^a ±0.001 (0.001-0.002)	0.006 ^a ±0.004 (0.001-0.009)

NB: Values are expressed as mean ± standard deviation (range in parenthesis). Means in the same column bearing different superscripts are significantly ($p < 0.05$) different. LC1: location 1 before Ibelebiri community. LC2: Location 2 within Ibelebiri Community. LC3: Location 3 after Ibelebiri community. LC4: Location used as Control.

3.2 Heavy Metals Concentration in the Sediments of Kolo Creek at Ibeleberi Community

The result of seasonal average of heavy metals (Ni, Cr, Cu, Pb, Zn, Co, Cd) concentrations in sediments sampled from Kolo creek at Ibeleberi axis is presented in table 2 below. The results showed that Zn at LC1 with a mean value of 35.865 ± 0.000 mg/kg had the highest concentration. The Zn concentration in the sediment at LC1 ranged from 0.048 to 35.865 mg/kg. The lowest concentration measured in the sediments at LC1 (Cd with a mean value of 0.016 ± 0.000 mg/kg). The Cd concentrations at LC1 ranged from 0.011 to 0.016 mg/kg. The results also showed that some of the trace metals estimated showed a significant spatial variation. For example, the mean value of Ni at LC4 is 0.032 ± 0.002 mg/kg, while a lower value of 0.025 ± 0.000 mg/kg was obtained at LC1. Thus, there could be factors responsible for the distribution of heavy metals in the sediments. Generally, the heavy metals concentrations in the sediments showed a decreasing trend of $Zn > Pb > Cu > Co > Cr > Ni > Cd$. The results also revealed that measured values for Cd, Pb, Ni, Cr, Cu and Zn are below the permissible limit (31, 32). However, a measured value for Co at locations LC3 and LC4 appears to be slightly above the permissible limit value of 20 mg/kg for sediments (31,32). This suggests the sediments in these locations may have been contaminated with Co. The levels of concentrations of nickel and cadmium observed in this study are below the concentrations reported by Uzoekwe and Aigberua (9) for surface waters within the vicinity of a flow station in Kolo Creek, Nigeria. Though, the mean value of Pb in LC2 is below the permissible level, the concentration obtained at LC1 and LC3 are higher than the level observed by Uzoekwe and Aigberua (9) in the surface waters within the flow station in Kolo creek, Nigeria. The mean concentration of Zn obtained from the sediments in locations LC1 and LC3 are above the value while the mean concentration of Cu obtained in this study are above the value measured at river Benue (28). The mean concentration of Cd, Cr, and Ni are found to be lower than the levels got in the study of dredged and non-dredged rivers in Lagos, Nigeria (33). The mean concentration of Cu in LC 1 and 3 are higher than the levels obtained in Meghna River ecosystem (34). Furthermore, the mean levels of Cr, Ni and Cd are lower than the levels obtained in Meghna River ecosystem (34). Finally, Pb concentration obtained from sediments collected from sampling locations LC1 and LC3 in this study are relatively higher than the levels observed in Belawan Harbor (35).

Comment [OM24]: Report using ranges from each station.

Comment [OM25]: No information on significant differences and interpretation!

Comment [OM26]: Mean.....

Table 2: Heavy metals concentration (mg/kg) in sediments of Kolo creek at Ibelebiri community

Location	Ni	Cr	Cu	Pb	Zn	Co	Cd
LC1	0.025 ^a ±0.000 (0.003-0.025)	0.041 ^a ±0.000 (0.008-0.041)	20.251 ^b ±0.000 (0.330-20.251)	35.553 ^a ±0.000 (0.020-35.865)	35.865 ^a ±0.000 (0.048-35.865)	16.325 ^a ±0.000 (0.002-16.325)	0.016 ^b ±0.000 (0.011-0.016)
LC2	0.027 ^a ±0.009 (0.002-0.033)	0.045 ^a ±0.012 (0.004-0.053)	23.560 ^a ±0.630 (0.026-24.006)	31.601 ^b ±1.330 (0.021-32.538)	30.675 ^{ab} ±3.280 (0.036-32.995)	18.217 ^a ±5.500 (0.002-22.105)	0.021 ^a ±0.007 (0.008-0.0021)
LC3	0.035 ^a ±0.020 (0.001-0.036)	0.036 ^a ±0.000 (0.002-0.036)	24.006 ^a ±0.000 (0.022-24.006)	30.664 ^b ±0.000 (0.028-30.664)	28.355 ^b ±0.000 (0.043-28.355)	22.105 ^a ±0.000 (0.001-22.105)	0.021 ^a ±0.000 (0.010-0.021)
LC4	0.032 ^a ±0.002 (0.001-0.033)	0.037 ^a ±0.001 (0.002-0.038)	23.117 ^a ±1.260 (0.024-24.006)	31.331 ^b ±0.940 (0.022-31.998)	30.311 ^{ab} ±2.770 (0.041-32.266)	21.104 ^a ±1.420 (0.001-22.105)	0.020 ^a ±0.001 (0.009-0.021)

NB: values are expressed as mean ± standard deviation (range in parenthesis). Means in the same column bearing different superscripts are significantly (p<0.05) different. LC1: location 1 before Ibelebiri community. LC2: Location 2 within Ibelebiri Community. LC3: Location 3 after Ibelebiri community. LC4: Location used as Control

Comment [OM27]: delete

Table 3: Therange of geoaccumulation index (mean in parenthesis) of the heavy metals in sediments studied.

Sampling	Geoaccumulation index						
Location	Pb	Cd	Cr	Cu	Zn	Co	Ni
LC1	-10.040 to - 0.760 (-2.840)	-4.770 to -4.230 (-4.230)	-14.190 to - 11.840 (-12.620)	-7.830 to - 1.890 (-3.870)	-11.190 to -1.650 (-4.830)	-13.870 to - 0.881(-5.210)	-15.290 to - 12.230 (-13.250)
LC2	-9.970 to - 0.630 (-2.930)	-5.230 to -3.840 (-4.320)	-15.190 to - 11.470 (-12.900)	-11.490 to - 1.640 (-4.950)	-11.610 to -1.770 (-5.120)	-13.870 to - 0.440 (-5.130)	-15.870 to - 11.830 (-13.420)
LC3	-9.550 to 0.550 (-2.820)	-4.910 to -3.840 (-4.190)	-16.190 to - 12.020 (-13.410)	-11.740 to - 1.640 (-5.010)	-11.350 to -1.990 (-5.110)	-14.870 to - 0.440 (-5.250)	-16.870 to - 11.700 (-13.470)
LC4	-9.900 to 0.610 (-2.910)	-5.060 to -3.840 (-4.290)	-16.190 to - 11.950 (13.390)	-11.610 to - 1.640 (-5.000)	-11.420 to -1.800 (-5.070)	-14.870 to - 0.440 (-5.300)	-16.870 to - 11.830 (-13.560)

LC1: location 1 before Ibelebiri community. LC2: Location 2 within Ibelebiri Community. LC3: Location 3 after Ibelebiri community. LC4: Location used as Control

Table 4: The range of contamination factor (mean in parenthesis) of the heavy metals in sediments studied.

Location	Contamination factor						
	Pb	Cd	Cr	Cu	Zn	Co	Ni
LC1	0 to 2.540 (1.690)	0.060 to 0.080 (0.070)	0	0.010 to 0.410 (0.270)	0 to 0.480 (0.320)	0.000 to 0.820 (0.540)	0
LC2	0 to 2.320 (1.510)	0.040 to 0.110 (0.080)	0	0 to 0.480 (0.310)	0 to 0.440 (0.270)	0 to 1.110 (0.610)	0
LC3	0 to 2.190 (1.460)	0.050 to 0.110 (0.090)	0	0 to 0.480 (0.320)	0 to 0.380 (0.250)	0 to 1.110 (0.740)	0
LC4	0 to 2.290 (1.490)	0.050 to 0.110 (0.080)	0	0 to 0.480 (0.310)	0 to 0.430 (0.270)	0 to 1.110 (0.700)	0

LC1: location 1 before Ibelebiri community. LC2: Location 2 within Ibelebiri Community. LC3: Location 3 after Ibelebiri community. LC4: Location used as Control

Comment [OM28]: delete

Table 5: Pollution load index (PLI) of the studied area

Sample location	PLI
LC1	0.371
LC2	0.393
LC3	0.441
LC4	0.418

LC1: location 1 before Ibelebiri community. LC2: Location 2 within Ibelebiri Community. LC3: Location 3 after Ibelebiri community. LC4: Location used as Control

Comment [OM29]: delete

3.3 Ecological Risk Assessment of Heavy Metals Pollution in Sediments

Comment [OM30]: delete

3.3.1 Geoaccumulation Index

The result of geoaccumulation index (I_{geo}) in sediments collected from Kolo creek transversing through Ibelebiri community in Ogbia L.G.A. is presented in table 3. Using the geoaccumulation index classification, the sediments across the four locations are uncontaminated with Cd, Cr, Cu, Zn, Co and Ni. The sediments collected from some locations LC1, LC3 and LC4 are moderately contaminated with Pb. The uncontaminated sediments imply that it is likely that these sediments heavy metal concentrations are within natural background range or have minimal anthropogenic influence. The negative values obtained for the geoaccumulation index calculation implies that the sediments have low ecological risk to the aquatic environment and living organism. However, sediments moderately contaminated with Pb due to anthropogenic activities pose serious ecological and human health risk.

Comment [OM31]: hydrobionts

Comment [OM32]: Recast this to include "potential risk"

3.3.2 Contamination Factor (CF) of Heavy Metals in the Sediments.

The results of the contamination factor of heavy metals in the studied sediments samples are shown in table 4. According to the CF classification, there is low contamination of the ecosystem by Cd, Cr, Cu, Zn, Ni and Co across the four locations. Finally, there is moderate contamination of Pb across the locations studied. This implies that Pb has been introduced into the sediments from human activities, such as industrial emissions, vehicular exhaust, leaded gasoline.

Comment [OM33]: Back up with references

3.3.3 Pollution Load Index (PLI)

A $PLI < 1$ indicates there is no pollution in the 4 locations studied or that the concentration of the pollutants is below detection limit as shown in Table 5. This suggests that the study area is free from heavy metal contamination or that the contamination is very low. However, using the PLI alone, may not reflect the true status of the sediments (36) because it does not consider the bioavailability, toxicity and speciation of the metals.

Comment [OM34]: And mobility

3.4 Principal Component Analysis

The result of principal component (PCA) is represented in table 6. Two major factors were identified; uncontrolled combustion process, such as artisanal refinery activities and gas flaring from nearby flow

station. The percentage variance and cumulative results further highlighted the emergence of a pollution source other than gas flaring.

Table 6: PCA of Heavy Metals in Sediments

Component	Eigen value	% Variance	Cumulative %
1	5.663	80.896	80.896
2	1.321	18.866	99.762

Two factors identified: Uncontrolled combustion processes and activities from nearby oil and gas production including gas flaring.

The choice of Ibelebiri community through which Kolo creek flows before it gets to Imiringi is in order to investigate if there are other pollution determinants at Ibelebiri or a pollution source other than activities due to hydrocarbon production such as gas flaring. Although wind direction can determine eventual deposit of particulate matter arising from gas flaring, plume of gas flaring at Imiringi flow station shows otherwise.

Comment [OM35]: Support with references

4. CONCLUSION

This study showed that water flowing from Kolo creek along Ibelebiri community axis had Pb and Cd in concentrations deemed unsafe, and with sediments that had Co in concentrations above approved level. This should raise concerns as this could lead to serious health issues in the community. PCA result identified two major factors; uncontrolled combustion processes such as artisanal refinery activities and gas flaring from nearby flow stations. These contributed to the pollution of the water and sediments in the studied area. Though several studies attributed pollution of the creek to mainly oil and gas production at Imiringi, this study reveals an emerging source of heavy metal contaminants that ultimately contribute to the pollution load of Kolo creek.

Comment [OM36]: ????

Comment [OM37]: revealed

REFERENCES

1. Dey, M., Akter, A., Islam, S., Dey, S.C., Choudhury, T.R., Faterma, K.J. and Begum, B.A. (2021) Assessment of contamination level, pollution risk and source apportionment of heavy metals in the Halda River water, Bangladesh. *Heliyon*, 2021;7, e08625.
2. Avkopashvili, M., Avkopashvili, G., Avkopashvili, I., Asanidze, L., Matchavariani, L., Gongadze, A. and Gakhokidze, R. Mining - Related Metal Pollution and Ecological Risk Factors in South-Eastern Georgia. *Sustainability*, 2022;14, 5621.
3. Xiao, H., Shahab, A., Xi, B.D., Chang, Q.X., You, S.H.; Li, J.Y., Sun, X.J., Huang, H.W. and Li, X.K. Heavy metal pollution, ecological risk, spatial distribution, and source identification in sediments of the Lijiang River, China, *Environ. Pollut.* 2020; 269: 116-189.
4. Jiunan, G. and Jia, W. Heavy metals in Yinma River sediment in a major Phaeozems zone, Northeast China: Distribution, chemical fraction, contamination assessment and source apportionment: *SCientifiCRePorTS*2018;8:12231.
5. Zhang, J., Hua, P. and Krebs, P. Influences of land use and antecedent dry-weather period on pollution level and ecological risk of heavy metals in road-deposited sediment. *Environ. Pollut.*2017;228: 158–168.
6. Wei, H., Hu, D. and Li, K.B. Sediments contamination levels of organochlorine pesticides in Weihe River, Northwestern China. *Environ. Earth Sci.* 2016;75, 1–8.
7. Usman, Q.A., Muhammad, S., Ali, W., Yousaf, S. and Jadoon, I. Spatial distribution and provenance of heavy metal contamination in the sediments of the Indus River and its tributaries, North Pakistan: Evaluation of pollution and potential risks. *Environ. Technol.Innovation.*2020; 21:101184.
8. Uzoekwe, S. A. and Oghosanine, T. A. The Effect of Refinery and Petrochemical Effluent on Water Quality of Ubeji Creek, Warri, Southern Nigeria. *Ethiopian J. of Environ. Stud. and Mgmt.*2011; 4:107-112.

Comment [OM38]: provide DOI for journals cited

9. Uzoekwe, S. A. and Aigberua, A. O. Heavy metal distribution and assessment of ecological risk in surface waters and sediments within the flow stations in Kolo Creek, Nigeria. *J. chem. soc. Nig.*, 2019;44 (5): 809-824.
10. Ogamba, E. N., Ebere, N. and Izah, S. C. Heavy metal Concentration in water, sediment and Tissues of *Eichhornia Crassipes* from Kolo Creek, Niger Delta, *Greener J. Environ. Manage. PublicSafety*, 2017;6(1):1-5.
11. Enevide, C. and Chukwuma, K. C. Oil spillage and heavy metals toxicity in the Niger Delta. *J. Health Pollution* 2018;19: 180905.
12. Agoro, E. S. and Ikimi, C. G. Translocation capacity of some heavy metals in aquatic food chain of crude oil impacted community of Imiringi in Bayelsa State, Nigeria. *Toxic. Research and Appl.* 2022;6. doi:[10.1177/23978473221146222](https://doi.org/10.1177/23978473221146222)
13. Ali, M. Y., Amin, M. N. and Alam, K. Ecological Health Risk of Buriganga River, Dhaka, Bangladesh. *Hydro Nepal*, 2008;3, 25-28.
14. Moniruzzaman, M., Elahi, S. F. and Jahangir, M. A. A. Study on Temporal Variation of Physico-chemical Parameters of Buriganga River Water through GIS (Geographical Information System) Technology. *Bangladesh J. Sci. Ind. Res.* 2009;44 (3), 327-334.
15. Alam, A. M. S., Islam, M. A., Rahman, M. A., Siddique, M. N. and Matin, M. A. Comparative study of the toxic metals and non-metal status in the major river system of Bangladesh. *Dhaka Univ. J. Sci.*, 2003;51 (2): 201-208.
16. Ahmad, M. H., Dhiaa, N. J. and Arkan, R. A. Spatial distribution and evaluation of heavy metals in surface sediments of the Al-Najaf Sea depression reservoir, Iraq. *Alex. Engr. J.* 2020;09.049.
17. Bassey, S. E., Ohimain, E. I. and Angaye, T. C. N. The Molluscicidal Activities of Methanolic and Aqueous Extracts of *Jatropha curcas* Leaves against *Bulinus globosus* and *Bulinus rholfsi*, Vectors of Urinary Schistosomiasis. *J. of Parasitology*, 2013;103: 115-122.
18. Csuros, M. and Csuros, C. Environmental Sampling and Analysis for Metals. 1st edition. CRC Press; 2002.
19. Loska, K., Wiechula, D., Barska, B., Cebula, E. and Chojnecka, A. Assessment of Arsenic Enrichment of Cultivated Soils in Southern Poland. *Pol. J. Environ. Stud.*, 2003;12:187-192

20. Chakravarty, I. M. and Patgiri, A. D. Metal pollution assessment in sediments of the Dikrong River, N.E. India. *J. Hum. Ecol.* 2009;27 (1),63–67.
21. Tomlinson, D. C., Wilson, J. G., Harris, C. R. and Jeffrey, D.W. Problems in the assessment of heavy metals in estuaries and the formation of a pollution index. *HelgolWissMeeresunters.*1980; 33:566-575
22. WHO (World Health Organization). Guidelines for drinking-water quality, 4th edition, incorporating the 1st edition addendum; 2017. <https://www.who.int/publications/i/item/9789241549950>
23. Nigerian Standard for Drinking Water Quality (NSDWQ). Standard Organisation of Nigeria. Nigeria Industrial Standard, NIS-554-2015; 2015
24. Ogbonna, P. C., Ukpai, N and Obasi, K. O. Assessment of metal contamination in Ubeyi River and accumulation in fish and sediment. *J. of Applied Sci. and Environ. Managmt.* 2018;22(8): 1151–1157.
25. Adelekan, B. A. and Abegunde, K. D. Heavy metals contamination of soil and groundwater at automobile mechanic villages in Ibadan, Nigeria. *Int. J. Physic. Sci.* 2011;6: 1045 – 1058.
26. Al Hagibi, H. A., Al-Selwi K. M., Nagi, H. M. and Al- Shwafi, N. A. Study of Heavy Metals contamination in Mangrove sediments of the red sea coast of Yemen from Al-Salif to Bab-el-Mandeb Strait. *J. Eco. & Natural. Resorc.* 2018;2(1):121.
27. Iwegbue, C. M. A., Faran, T. K., Iniaghe, P. O., Ikpefan, J. O., Tesi, G. O., Nwajei, G. E. and Martincigh, B. S. Water quality of Bomadi Creek in the Niger Delta of Nigeria: assessment of some physicochemical properties, metal concentrations, and water quality index. *Appli. Water Sci.* 2023; 13:36. Doi: 10.1007/s13201-01804-2.
28. Egesi, O. C., Alum-Udendi, O., Ogbonna, P. C. and Ugor, N. N. Assessment of Heavy metals in water, Fish, and Sediment of River Benue, Benue State, Nigeria. *Nig. Agric. J.*2023;54: 55-67.
29. Ibigbami, T. B., Odeja, O. O., Akinyeye, O. J., Sunkanmi, A. A. and Isaac, B. O. Advanced Indexed Model Analysis of Heavy metals pollution of River Ureje, Ado Ekiti, Nigeria. *J. Environ. Pollut. and Contr.* 2023;6(1):103.

30. Odoemelam, S. A., Edori, M. N. and Ogbuagu, M. N. Assessment of heavy metal status of Orashi river along the Engenni axis, Rivers State of Nigeria. *Comm. Physic. Sci.* 2019;4(2): 74-80.
31. DPR, Department of Petroleum Resources. Environmental Guidelines and Standards for the Petroleum Industry in Nigeria, EGASPIN 2 nd Ed;2002; p14, 280.
32. DPR, Department of Petroleum Resources. Environmental Guidelines and Standards for the Petroleum Industry in Nigeria, EGASPIN 3rd Ed;2018; p22.
33. Jolaosho, T. L., Elegbede, I. O., Ndimele, P. E., Mekuleyi, G. O., Oladipupo, I. O. and Mustapha, A. A. Comprehensive geochemical assessment, probable ecological and human health risks of heavy metals in water and sediments from dredged and non-dredged rivers in Lagos, Nigeria. *J. Hazardous Mat. Advc.* 2023; 12:100379.
34. Islam, S. Md., Hassan, K. Md., Bhuiyan, I. D. Md. and Sajeed, M. I. Heavy metal contamination in surface water and sediment of the Meghna River ecosystem. *Int. j. adv. Res. Biol. Sci.* 2023;10(4):22-45.
35. Sulistyowati, L., Yolanda, Y. and Andareswari, N. Harbor water pollution by heavy metal concentrations in sediments. *Global J. Environ. Sci. and Managmnt.* 2023;9(4):885-898.
36. Frascareli, D., Silva, S. C., Mizael, J. D. O. S. S., Rosa, A. H., Pompeo, M. L. M., Lopez-Doval, J. C. and Moschini-Carlos, V. M. Spatial distribution, Bioavailability, and Toxicity of metals in surface sediments of tropical reservoirs, Brazil. *Environ. Moni. Assessmt.*2018;190(4) DOI:10.1007/s10661-018-6515-8.