Gastrointestinal helminth parasites, stomach content analysis and condition factor of *Bostrychus africanus* (sleeper gobies) and *Periophthalmuspapilio*(mudskipper)of Ikpukulu-Ama Creek, Port Harcourt, Rivers State, Nigeria

Abstract

Bostrychus africanus and *Periophthalmuspapilio* are important fishes ~~of~~parts of southern Nigeria. However, there is dearth of information on the gastrointestinal helminths of these gobiids, especially from the Ikpukulu-Ama Creek, Port Harcourt,Rivers State, Nigeria. This research was, therefore, necessary to provideinformation on the parasites as well as stomach content analysis and condition factor of both fish species ~~for academic and reference purposes~~. A total of 62 *Bostrychusafricanus* and thirty-six (36) *Periophthalmuspapilio* were caught ~~for analysis~~ using basket traps on the mudflats of the creek. Sampling was done in September 2023. In the laboratory, samples were identified and examined for parasites using standard techniques. Nematodes, *Raphidascaroides africanus*(16.1%), were isolated from *B. africanus*,whereasacanthocephalans,*Neoechinorhynchus* sp.(2.8%), were isolated from *P. papilio*. The result of this study suggests a parasite preference of *B.africanus* over *P. papilio*.Stomach content analysis showed presence of crab parts and muddy sediments in *B. africanus*. Condition factor was better in *B. africanus* (1.1 to 1.8) than in *P. papilio* (0.7 to 1.1). There is a dearth of information on the stomach content of *P. papilio*and therefore, is suggested for future studiesand scientific reasons for parasite preference of *B.africanus* over *P. papilio* should be elucidated.

Keywords: Acanthocephalans, Bonny River, gastrointestinal helminths, gobiids, mudflats.

Introduction

Fishes are a significant food source in Africa, particularly in the Niger Delta of Nigeria (Ezenwaka and Nweke, 2021), playing a crucial ecological and economic role in the food chain (Abowei and Ezekiel, 2013). Fish is known to provide vital nutrient such as omega-6 oil and the cholesterol level is relatively very low (Abu and Agarin, 2016). Gobiids, such as *Bostrychus africanus* and*Periophthalmuspapilio*, are carnivoresthat feed on crustaceans and other benthic invertebrates, small fishes and insects. Many species pass through a marine larval stage during which they feed on planktons (Ugbomeh*etal.*, 2018). The analysis of fish stomach content, which

Comment [JS1]: Gastrointestinal Helminth Parasites, Stomach Contents and Condition Factors of *Bostrychus africanus* (Sleeper gobies) and *Periophthamuspapilie* (Mudskipper) found in Ikpukulu-Ama Creek, Port Harcort, Rivers State, Nigeria

Comment [JS2]: in

Comment [JS3]: a dearth

Comment [JS4]: delete

Comment [JS5]: delete

Comment [JS6]: state the names of the techniques used for the identification of the specimens and sample analysis for the detection of the parasite

Comment [JS7]: examined

Comment [JS8]: observed

Comment [JS9]: the presence

Comment [JS10]: the parasite

Comment [JS11]: nutrients

Comment [JS12]: plankton

UNDER PEER REVIEW

depends on food availability and varying by location, provides information about their feeding habits and culture potential (Chukwu and Princewill, 2019).

Comment [JS13]: varies

Nonetheless, fish health can be negatively impacted by parasites, especially helminth parasites, which leave fish susceptible to infections (Kaur, 2014). Fish health is determined by their condition factor, which varies by species and season (Seiyaboh (2018), and is influenced by anthropogenic activities, food abundance, diet, ideal temperature, and salinity (Abowei, 2009; Atama *etal.*, 2013; Ugbomeh*et al*., 2018). Fish regularly ingest larval stage of helminth parasites because of the abundance and diversity of these parasites in the aquatic ecosystem (Lagrue*etal.*, 2011). These parasites can obstruct the intestinal tract of fish, affecting food digestion and absorption, leading to morbidity and mortality, economic losses, and a potential risk of zoonosis (Kayode *etal.*, 2012; Soham and Gadahar, 2016).

Comment [JS14]: the larval

Gobiidsare commonly infected by gastrointestinal helminths.However, scientific reporting of parasitic infections in these fish species is scanty. Ugbomeh*etal*. (2018) found Ascaridida nematodes in *B. africanus* and *P. papilio*, with higher parasite infection prevalence in *B. africanus*. Robert *etal*. (2022) found *Raphidascaroidesafricanus* in Rumuolumeni and Afikpo markets in Port Harcourt, Nigeria; while, Elele and Aziaka (2019) reported the presence of monogenean, digenean, cestode and nematode parasites in *Periophthalmus* species obtained from some waterfronts in Rivers State, Nigeria. A few authorsin Nigeria have reported helminth parasite infestations in freshwater fish in Nigeria, but there doesn't seem to be much information available about fish parasites in the brackish water habitat, with only a few publications on brackish water (Akinsanya *etal.*, 2007; Oribhabor*etal.*, 2010; Worgu and Okaka 2012; Oribhabor*etal*., 2012). The Ikpukulu-Ama creek has been examined for some studies including assessment of cadmium and lead concentrations in barnacles (*Semibalanusbalanoides*) and

Comment [JS15]: "Creek" with capital C

UNDER PEER REVIEW

oysters (*Crassostrea gasar*) (Ugbomeh and Okereke, 2015), study of phytoplankton communities (Okoseimiema*et al*., 2020), heavy metal concentrations of surface water, sediment samples, fish and plankton from same creek (Ijeoma and Joshua, 2019; Okoseimiema and Vincent-Akpu, 2020), etc. There is paucity of information regarding aspects of the parasite communities infecting fish inhabiting the Ikpukulu-Ama Creek. This research was therefore conducted to examine two gobiids, *B. africanus* and *P. papilio* for their gastrointestinal helminth parasites, their stomach content analysis and condition factor, contributing to biological research data related to thecreek.

Comment [JS16]: a paucity

Materials and Methods

Study Location

Ikpukulu-Ama Creek (Figure1) is located at Latitude N 4° 44'6.65320" and Longitude E 7°1'40.61060". It is a small tidal creek in the Niger Delta region that flows through a small fishing settlement behind Government Comprehensive Secondary School, Borikiri in Port Harcourt City Local Government Area of Rivers State, Nigeria. The creek is also a tributary of the Bonny River that flows into the Atlantic Ocean. It has salinity values ranging from 5 ppt to 35ppt and it is surrounded by dredging facilities and some petroleum oil servicing companies (Okoseimiema*et al*., 2020).

Comment [JS17]: provide with the map indicating the study location

Comment [JS18]: Remove the space

Sampling Protocol

Fish samples (sixty-two specimens of *Bostrychusafricanus* and thirty-six *Periophthalmuspapilio)* were caught from the Ikpukulu-Ama Creek at Borokiri by the use of basket traps set in the

UNDER PEER REVIEW

muddy part of the creek and examined for gastrointestinal helminth parasites, stomach content analysis and condition factor in this study.Sampling was done weekly in September 2023. The fish samples were transported in ice chest to the Parasitology Laboratory, Department of Animal and Environmental Biology, Rivers State University, Port Harcourt, for identification and parasitological examination.

Comment [JS19]: in an ice chest



Fig. 1: Photograph of Ikpukulu-Ama Creek, Borikiri, Port Harcourt, showing its geographic coordinates

Fish samples were identified using the taxonomic keys described by Ssentongo *et al*. (1986) and Schneider (1990). The total length was measured using a meter rule while the body weight of each fish specimen was measured using a sensitive weighing balance (model: Camry EK5350).Microsoft Excel was used to computeFulton's condition factor (CF) according to Zhelev *et al.* (2015) as follows: $CF= BW /TL^3 \times 10^2$, where BW is the wet body weight of fish in g and TL is total length of fish in cm.

Comment [JS20]: a total

UNDER PEER REVIEW

For parasitological examination, a longitudinal section was made through the anal pore of each fish specimen. The exposed gastrointestinal tract was excised, slit longitudinally and examined microscopicallyfor parasites in Petri dishes containing physiological saline. The contents of the Petri dishes were carefully examined for analysis of the stomach content.Parasite species presentwere identified according to Paperna (1996) and Moravec (2019) and were fixed in 70% ethanol.

Data analysis

Parasite prevalence and mean intensity of infection were calculated according to the formula of Bush *et al*., (1997).

Results

Morphometric characteristics and Condition Factor of *Bostrychusafricanus* and *Periophthalmuspapilio* ofIkpukulu-Ama Creek, Borikiri, Port Harcourt

*Bostrychus africanus*specimens examined in this study had wet body weight ranging from 10.0g to 21.0g with a mean value of about 12.9g. Their total length values ranged from 8.8cm to 11.3cm, and their condition factor, 1.1 to 1.8 with a mean value of 1.4 (Table 1).

The *P. papilio* samples examined had wet body weight ranging from 4.0g to 15.0g. Their total length was in the range of 7.9cm to 11.5cm, with a mean value of 9.6cm. The condition factor of these *P. papilio* specimens ranged from 0.7 to 1.1 with a mean value of 0.9 (Table1).

Comment [JS21]: weights

Comment [JS22]: weights

Comment [JS23]: remove the space

Table 1: Morphometric Values and Condition Factor of*Bostrychusafricanus* and *Periophthalmuspapilio*, Ikpukulu-Ama Creek, Borikiri, Port Harcourt, Rivers State.

	Bostrychusafricanus	*Periophthalmuspapilio*

UNDER PEER REVIEW

Parameters	Range	Mean	Range	Mean
Wet Body Weight (g)	10.0 – 21.0	12.9 ± 2.5	4.0 – 15.0	8.0 ± 2.9
Total Length (cm)	8.8 – 11.3	9.8 ± 0.5	7.9 – 11.5	9.6 ± 1.0
Condition Factor	1.1 – 1.8	1.4 ± 0.2	0.7 – 1.1	0.9 ± 0.1

± standard deviation

Comment [JS24]: remove the space

Parasitic Helminths Isolated from *Bostrychus africanus* and *Periophthalmuspapilio*ofIkpukulu-Ama Creek, Borikiri, Port Harcourt

Two parasitic helminth species, *Rhaphidascaroides africanus* (Nematoda) and*Neoechinorhynchus*sp. (Acanthocephala),were encountered in this study.

Ten (16.1%) of *B. africanus* examinedwere infected with *Rhaphidascaroides africanus*with mean intensity of approximately one parasite per infected host. Only one (1) *P.papilio* specimen was infected with the acanthocephalan, *Neoechinorhynchus*sp, showing aprevalence of 2.8% (Table2).

Comment [JS25]: a mean intensity

Table 2: Prevalence and mean intensity of parasites of *Bostrychus africanus* and *Periophthalmuspapilio*, Ikpukulu-Ama Creek, Borikiri, Port Harcourt

Parasites	**Fish Species**			
	Bostrychus africanus		*Periophthalmuspapilio*	
	Prevalence (%)	Mean Intensity (±Stdev)	Prevalence (%)	Mean Intensity (±Stdev)
Raphidascaroides africanus	16.13	1.4 ± 1.0	-	-
*Neoechinorhynchus*sp.	-	-	2.78	1.0 ± 0.0

UNDER PEER REVIEW

Stomach content analysis of *Bostrychus africanus* and *Periophthalmuspapilio*, Ikpukulu-Ama Creek, Borikiri, Port Harcourt

The stomach content of *B. africanus*included parts of crabs and muddy sediments. The crabs were present in thestomach of ten (10)*B. africanus* while others were mostly composed of muddy sediments. The feeding habit of the *P. papilio* from this River could not be ascertained as there were no fragments of particles left in the stomach of the samples examined, except for mud particles.

Discussion

According to the study, the mean length of the *B. africanus* and *P. papilio* samples was 9.8 cm and 9.6 cm, respectively. On the other hand, the samples analyzed by Ugbomeh*etal*. (2018) were not significantly less in length, but they were heavier.Abowei and Hart (2008) and Ogamba*etal*. (2014) hypothesized that heavier fish samples of a given length are in better condition than less weighty fish.

This study result revealed the presence of the gastrointestinal nematode, *Raphidascaroides africanus* infective in *B. africanus*, and the acanthocephalan, *Neoechinorhynchus* sp. infective in *P. papilio*. Elsewhere, occurrence of nematode parasite had been recorded in *B. africanus* and *P. papilio*, though of different species. Ugbomeh*et al*. (2018) recovered Ascarid nematodes in their work which is different from the species recorded in this study.*Raphidascaroides africanus*infestation was also reported by Robert *et al.* (2022) in *B. africanus*. These parasites are commonly associated with *B. africanus* in Africa (Moravec, 2019). Acanthocephalan infection in gobiids had not been well documented as there ispaucity of information on the subject. Nematodes can obstruct the intestinal tract of fish, affecting food digestion and absorption, and

Comment [JS26]: the occurrence

Comment [JS27]: a paucity

UNDER PEER REVIEW

their presence in fish indicates a potential risk of zoonosis (Kayode *etal.*, 2012;Soham and Gadahar, 2016).

It was observed that prevalence of parasite infection and mean intensitywas higher in *B. africanus*than*P. papilio*, which is similar to the report of Ugbomeh*et al*. (2018). Both gobiids hereby examined have a parasitic mode of life and occupy similar ecological niches (Chukwu and Princewill, 2019; Abiaobo*et al*., 2021), so differences in parasite prevalence could simply be as a result of host preference by the parasites and the intensity depend on prevailing environmental conditions.

Several of the samples examined presented with empty stomach especially, the *P. papilio*, but the few with stomach content (which were ten (10) of *B. africanus*) all had crab parts and muddy sediments. This is in consonance with the report of Chukwu and Princewill (2019) that crab parts are the most important food items of *B. africanus*. However, it is important to note that stomach content depends on food availability and this would vary by sample location. Surprisingly, information on the stomach content analysis of *P. papilio* in Nigeria is sparse.

The mean condition factor for *P. papilio* was slightly below 1.0, though values ranged from 0.7 to 1.1. Abiaobo*et al.* (2021) reported similar condition factor values, between 0.9 and 1.2. Similarly, Moslen and Daka (2017) reported condition factor of *P. papilio* ranging from 1.12 to 1.45. In this present study, *B. africanus* had condition factor which indicated good health status, having values ranging from 1.1 to 1.8, and a mean of 1.4. Chukwu and Ansa (2017) reported mean condition factor of 1.35 for the same species from New Calabar River, Nigeria.Condition factor of fish, which varies by species and season according to Seiyaboh (2018), is a measure of the physiological state of the fish and is dependent on anthropogenic activities, food abundance,

Comment [JS28]: the prevalence

Comment [JS29]: stomachs

Comment [JS30]: the condition

Comment [JS31]: a condition

Comment [JS32]: a mean

Comment [JS33]: The condition

UNDER PEER REVIEW

diet, the ideal temperature and salinity for growth (Abowei, 2009; Atama *etal*., 2013;Ugbomeh*et al*., 2018). Fish health is also impacted by parasites which leaves them vulnerable to subsequent infections by pathogenic agents.

Conclusion

The study revealed that the gastrointestinal helminth parasites of *Bostrychusafricanus* and *Periophthalmuspapilio*of Ikpukulu-Ama Creek, Rivers State, Nigeria are composed of nematodes and acanthocephalans. *B.africanus* had a higher prevalence of parasite infection. Stomach content analysis was only possible in *B. africanus*, which was mainly comprised of Crab parts, while P. papilio had an empty stomach.Condition factor was better in *B.africanus* than in *P. papilio*. Literature on the stomach content of *P. papilio* is scarce and is suggested for future studies. Reasons for parasite preference of *B. africanus* over *P. papilio* should be elucidated.

Comment [JS34]: italicize

References

Abiaobo, N.O., Asuquo, I.E., Ejiogu, I.N. and James, E.J. (2021). Aspects of the biology of *Periophthalmusbarbarus* (Mudskipper), from Jaja Creek, Niger Delta, Nigeria. *Ecology and Evolutionary Biology*, 6, 15-22.

Abowei, J. F. N (2009), The Abundance Condition Factor and Length-Weight Relationship of Some *Sardinella maderensis* (Jenyms, 1842) from Nkoro River, Niger Delta, Nigeria. *Advanced Journal of Food Science and Technology*, 1(1):65-70.

Abowei, J.F.N and Hart A.I (2008). Artisanal fisheries characteristics of the fresh water reaches of the lower Nun River, Niger Delta, Nigeria. Journal of Applied Science and Environmental *Management*, 12, 5-7.

Abu, O. M. G. and Agarin O. J. (2016). Length-weight relationship and condition factor of Silver Catfish (*Chrysichthysnigrodigitatus*) from the Lower Reaches of the New Calabar River Niger Delta. *International Journal of Innovative Studies in Aquatic Biology and Fisheries (IJISABF*), 2(4), 1-7.

Akinsanya, B., Otubanjo, A. and Ibidapo, C. (2007). Helminth biload of *Chrysichthysnigrodigitatus* from Lekki Lagoon, Lagos, Nigeria. *Turkish Journal of Fisheries and Aquatic Sciences*, 4(7), 83–87.

UNDER PEER REVIEW

Atama, C., Okeke, O., Ekeh, F. and Ezenwaji, N. (2013). Length -Weight relationship and condition factor of six cichlid species of Anambra River. *Journal of Fisheries and Aquaculture*, 4(5), 82-86.

Bush, A.O., Lafferty, K.D., Lotz, J.M. and Shostak, A.W. (1997). Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology*, 83(4), 575–583.

Chukwu, K.O. and Ansa, E.J. (2017). Length-weight relationship and condition factor of *Bostrychus Africanus* from New Calabar River, Nigeria. *Nigerian Journal of Fisheries*, 16, 1584-1590.

Chukwu, K.O. and Princewill, K. (2019). Stomach Content of the Sleeper Fish, *Bostrychus africanus* from the New Calabar River, Nigeria. *Research & Reviews: Journal of Veterinary Science and Technology*, 8(3), 13-16.

Elele, K. and Aziaka, B.O. (2019). Evaluation of parasites associated with mudskipper *Periophthalmus* spp. sold in selected waterfronts market in Port Harcourt Metropolis, Rivers State, Nigeria. *Nigerian Journal of Parasitology*, 40(2), 169-174.

Ezenwaka, C. O. and Nweke, O. S. (2021). Prevalence of gastrointestinal helminthesassociated with *Synodontisschall* from Otuogori River, Ogbia Local Government, Bayelsa State. *IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS)*, 14(3), 06-10.

Ijeoma, F.V. and Joshua, O.I. (2019). Trace metals in water, sediment and biota from Ikpukulu-Ama Creek Port Harcourt, Rivers State, Nigeria. *Preprints.org*; DOI: 10.20944/preprints201904.0145.v1.

Kaur, P. (2014). Histopathological effect of Senga species (Cestode: Pseudophyllidae) in the intestine of Piscia host. *World Journal of Pharmaceutical Sci*ence, 3(2),1506-1513.

Kayode, O., Okafor, J., Alade, A. and Oronaye, O. (2012). Helminth parasites of *Sarotherodongalilaeus*and *Tilapia zilli*, (Pisces; Cichlidea) from River Oshun, South West Nigeria. *International Journal of Aquatic Science*, 3(3), 49-55.

Lagrue, C., Kelly, D., Hicks, A., Poulin, R. (2011). Factors influencing infection patterns of tropically transmitted parasites among a fish community: Host Diet, Host – Parasite Compatibility or Both. *Journal of Fish Biology*, 79(4), 466-485.

Moravec, F. (2019). Parasitic nematodes of freshwater fishes of Africa. 1st edition. Czech Academy of Sciences, Czech Republic.

Moslen, M. and Daka, E.R. (2017). Length-weight relationship and condition factor of *Periopthalmuspapilio* (Bloch & Schneider, 1801) obtained from a tidal creek in the Bonny Estuary, Nigeria. *Journal of Aquaculture and Fisheries Management*, 1(1), 1-4

Ogamba E.N., Abowei, J.F.N. and Onugu, A. (2014). Length-weight relationship and condition factor of selected finfish species from Odi River, Niger Delta, Nigeria. *Journal of Aquatic Science*, 29(1A), 1-12.

Okoseimiema, I.J. and Vincent-Akpu, I.F. (2020). Distribution of trace metals in water, sedminent and biota from Ikpukulu-Ama creek, Port Harcourt, Rivers State, Nigeria. *Nigerian Journal of Fisheries*, 17(1), 1912-1924.

Okoseimiema, I.J., Vincent-Akpu, I.F., Onyeagbodor, P. and Oriakpono, E.O. (2020). Composition of phytoplankton communities in Ikpukulu-Ama Creek, niger delta, *International Journal of Contemporary Applied Researches*, 7(2), 21-31.

Oribhabor J, Ogbeibu E, Okaka E. (2012). The gastrointestinal helminth parasites of the threadfin fish, *Polydactyhisquadrifilis* (Family: Polynemidae) in a Niger Delta mangrove creek, Nigeria. *International Journal of Animal and Veterinary Advances*, 4, 240 -243.

Oribhabor, J., Ogbeibu, E. and Okaka (2010). E. The parasites gastrointestinal Helminth of some Scianid Species (Croakers) in a Niger Delta Mangrove Creek, Nigeria. *Tropical Freshwater Biology*, 91(7),15–23.

Paperna, I. (1996). Parasite, Infestations and Disease of Fishes in Africa—An Update. CIFA Technical Paper, 31, 1-220.

Robert, B., Wodi, S., Walters, V., Numdi, F.C. and Amuzie, C.C. (2022). Prevalence of *Raphidascaroidesafricanus* in market-derived gobies (*Bostrychus africanus*), Port Harcourt, Nigeria. *International Journal of Biological Research Advancement*, 2(1), 1-8.

Schneider, W. (1990). FAO Species Identification Sheets for Fishery Purposes. Field Guide to the Commercial Marine Resources of the Gulf of Guinea, Food and Agricultural Organization of the United Nations, Rome, Italy.

Seiyaboh, E. I. (2018). Variation in Condition Factor of Some Freshwater Fishes from Epie Creek, Bayelsa State, Nigeria. *Annals of Ecology and Environmental Science* 2 (2), 47-51

Soham, P. and Gadahar, D. (2016). Isolation and identification of Helmith parasite eggs having Zoonotic Potency from Sewage-Fed Pond Sediment, West Bengal, India. *International Journal of Advanced Research and Biol Science*, 3(1), 283-291.

Ssentongo, G.W., Ukpe, E.T. and Ajayi, T.O. (1986). Marine fishery resources of Nigeria: a review of exploited fish stocks. Published by Publications Division, Food and Agriculture Organization of the United Nations, Via delle Terme di Caracalla, 00100 Rome, Italy.

Ugbomeh, A.P. and Okereke, A.B. (2015). Assessment of cadmium and lead in the barnacle (*Semibalanusbalanoides*) and oyster (*Crassostrea gasar*) in Port Harcourt, Nigeria: Risk assessment of oyster consumption. *Research and Reviews: Research Journal of Biology*, 3(4), 7-11.

Ugbomeh, A.P., Okere, S., Sokari, G.M., Aisien, M.S.O. and Wala, C. (2018). Helminth parasites of gobies from two creeklets of the New Calabar River, Rivers State, Nigeria. *Asian Journal of Biology*, 6(4), 1-10.

Worgu, M. and Okaka, E. (2012). A comparative of Study gastrointestinal Helminth parasites infection of fresh and brackish water fishes from Warri River, Southern Nigeria. *African Research and Reviews*, 6(2), 13–23.

UNDER PEER REVIEW

Zhelev, Z.M., Popgeorgiev, G.S. and Mehterov, N.H. (2015). Changes in the hepatosomatic index and condition factor in the population of *Pelophylax ridibundus* (Amphibia: ranidae) from anthropogenically polluted biotopes in southern Bulgaria. Part II. *Bulgarian Journal of Agricultural Science*, 21, 517-522.

Comment [JS35]: reference should follow theguide of the journal

UNDER PEER REVIEW