

## Original Research Article

# In vitro screening of natural *Metarhizium* spp. isolates for bioinsecticidal and biofungicidal activity

### ABSTRACT

**Aims:** to increase conidia production of local *Metarhizium* spp.—strains by selecting the low-cost media to test obtained bioformulation against the sucking pests and fungal plant pathogens.

**Study design:** The phytophages like black bean aphid (*Aphis fabae*), a scale insect (*Aspidiotus nerii*) and spider mite (*Tetranychus urticae*) are economically essential pests in Kyrgyzstan. The current requirement is to use environmentally friendly protection to reduce the number of these pests. The entomopathogenic fungus from the *Metarhizium* genus was used in reducing the number of harmful pests in vitro and in vivo experiments.

**Place and Duration of Study:** Department of Plant Protection, Kyrgyz-Turkish Manas University between May 2021 and May 2023.

**Methodology:** Natural *Metarhizium* spp. strains were isolated from dead insect bodies of the Lepidoptera and Apidae families. A laboratory bioassay was conducted to evaluate isolated *Metarhizium* spp. strains against nymphal stages and adults of sucking pests like *A. fabae* (Blackfly), *A. nerii* (Scale insect) and *T. urticae* (Spider mite). The inhibitory effect of *Metarhizium* spp. against *Fusarium oxysporum*, *Alternaria* spp. and *Rhizoctonia solani* was detected using a dual culture technique.

**Results:** The results of in vitro bioassay tests against sucking pests have revealed the LC50 and LT50 values of TLK-1 isolate. The LC50 of this isolate for *A. fabae* adults was  $1 \times 10^6$  conidia mL<sup>-1</sup>; for *A. nerii*, it was lower -  $7 \times 10^5$  conidia mL<sup>-1</sup>; while for adults *T. urticae* it was higher -  $13 \times 10^6$  conidia mL<sup>-1</sup>. Such mortality occurred after 49.63 hours in adults of *A. fabae* (LT50), after 46.30 hours in *A. nerii* (LT50), and after 75.87 hours in deutonymph and adults of *T. urticae* (LT50). Three isolates, TLK-1, TLA-2 and BZTK1, have exhibited biofungicide activity in pathogen fungi like *Fusarium oxysporum* and *Alternaria* spp.

**Conclusion:** The found isolates can be simultaneously used against pathogenic fungi, essential for developing inexpensive biological formulas based on these fungi.

Keywords: *entomopathogenic fungus, Metarhizium* spp., *bioinsecticide activity, biofungicide activity, bioformulation for plant protection*

## 1. INTRODUCTION

Entomopathogenic fungi in nature are essential in keeping the number of agriculture and forest pests at an economically low-damaged level. The main natural habitats for these fungi are the soil, plant parts, plant rhizosphere, and insect body [ 1,2,3].

Fungi in the genus *Metarhizium* (Hypocreales: Clavicipitaceae) are considered a relevant biological agent for reducing pests' population and mass death. This entomopathogenic fungus affects about 200 types of insects from orders like Lepidoptera, Homoptera, Hymenoptera, Coleoptera, and Diptera. Among them are plant aphids, mites, thrips, greenhouse whiteflies, tomato whiteflies, and larvae of the butter beetle [4,5,6].

The fungus can survive saprophytically in the soil and be isolated from the soil using the 'Galleria' bait method[7]. Some studies recorded that *Metarhizium* species are often found in European, Central and South American, African and Asian countries [5]. According to the authors, southeastern Asia is the probable origin of the evolution of *Metarhizium* spp.[8 ,9]. *Metarhizium* spp. have asexual, cylindrical conidia, which are produced by spores. The conidia stick together to form chains and make large masses during asexual development [9,10, 11, 12].

Abiotic factors like humidity, sunlight and temperature (25°C) can be affected by the growth and mycelium formation of these fungi to efficiency to harmful insects. There are various facts about the effect of water on the fungus; *Metarhizium* spp. has a hydrophobic nature and does not mix well with water, forming float suspensions on the water. *Metarhizium* spp. mixed with Tween 80 and stored at room temperature for six weeks. The influence of humidity and temperature on the medium-term (3-4 months) storage of *Metarhizium* species conidia was studied. Conidia harvested after a 24-day culture on rice showed greater resistance to long-term storage than conidia from 12-day cultures. Dried conidia stored as a powder maintained a germination rate of 95% at 10-14°C but only up to 27% at 28-32°C. In another experiment, dried conidia maintained over 90% germination for 128 days, with or without silica gel. at 10 - 14 °C or -15 - -18° C [13, 14, 15].

Like other entomopathogenic fungi, *Metarhizium* spp. produces conidia that penetrate the cuticle of host insects and mites, infecting them by adhesion, germination, invasion, colonization, and dissemination [ 10, 16, 17]. Hence, *Metarhizium* spp. may have an advantage over other biological agents (i.e. bacteria and viruses) for some applications as a contact biopesticide. Because of its ability to rapidly produce infectious conidia when incubated on inexpensive nutrient media, there is significant interest in the commercial mass production of highly effective bioproducts with broad ranges based on *Metarhizium* spp. [2,3,18]. In vitro tests have recorded the ability of *M. anisopliae* to infect mite eggs of the spider mite *Tetranychusurticae*. At the same time, the mortality of mature stages (deutonymphs and imago) of this mite depended on the highest conidial concentration of fungi [19]. *M. anisopliae* could infect other species, like the citrus rust mite *Phyllocoptrutaoleivora* [20].

The *M. anisopliae* was tested against second and third larval instars of tomato pinworm *T. absoluta* [21, 22] in vitro conditions, and LT50 results were of 1.93 and 2.8 days at high ( $10^8$  spore/ ml<sup>-1</sup>) spore concentration. Based on other research results of *M. anisopliae*, strain Egy-5 infected adult females of *Aulacaspistubercularis* (Diaspididae), mortality rates were significantly higher ( $P = 0.04$ ) [23, 24, 25, 26]. A pest considered a severe pest, such as aphids, *M. anisopliae* has been shown to be effective against this insect [27, 28, 30].

Over the last two decades, studies have recorded a multifunctional lifestyle of *Metarhizium* spp. species; their colonization of plants may exert growth promotion, and immunity modulation affects them. Except as insect pathogens, these fungi can benefit their host plant through growth promotion and protection against stresses [ 31, 32, 33]. Plant growth promotion has been observed for multiple species of *Metarhizium* spp. in tomato [33, 34], maize [ 32, 35, 36, 37], soybean [ 38], peanut [39], potato [40], cassava [41], sweet pepper [42], switchgrass and haricot beans [ 43].


The phytophages like black bean aphid (*Aphis fabae*), a scale insect (*Aspidiotusnerii*) and, spider mite (*Tetranychusurticae*) are economically essential pests in Kyrgyzstan. The current requirement is to use environmentally friendly protection to reduce the number of these pests. Therefore, the search for fungi biosources, assessment of fungi virulence and conidia productivity are relevant to formulate a bioproduct based on *Metarhizium* spp. This study aims to increase conidia production by local *Metarhizium* spp strains by selecting the low-cost media to test obtained bioformulation against the sucking pests and fungal plant pathogens.

## **2. MATERIAL AND METHODS**

**2.1. *Metarhizium anisopliae* strains:** In this study *Metarhizium* spp. strains newly isolated from insects' cadavers were used.

**2.2. Methods of isolation from dead insects:** To remove the external microflora, the dead insect body was carefully washed in running water, then rinsed again with sterile distilled water and placed in a sterile cup (in some cases, if the studied object is resistant to alcohol, it can be rinsed with a small amount of ethyl alcohol), then 9 ml of sterile physiological solution added to the body, placed in a sterile porcelain container and crushed. The resulting suspension is diluted. The optimal dilution for fungi is  $10^2$  or  $10^3$ . A dilution of  $10^2$  was used in this work, the prepared solution of 0.3 mL was poured into sterile Petri dishes using a pipette, and 500 ml pre-cooled nutrient medium (Czapek agar (CZA), PDA, etc.) was added, and thoroughly mixed. Inoculated dishes were incubated at 25°C for 7-10 days for colonies to grow (Figure 1).

**2.3. Cultivation and morphological study of *Metarhizium* spp.strains:** *Metarhizium* spp. can grow in two conditions: on solid and liquid media, showing different growth phases. Its chlamydospore predominates in the liquid medium, and conidia in the dry mass are common in the solid medium. The result can be provided using liquid and solid media to produce chlamydospores and conidia. One percent from the liquid or solid medium of *Metarhizium* spp. conidia production will be sufficient.



**Figure 1.** The scheme of primary isolation of natural *Metarhizium* spp. strains from insects' cadavers

The morphology of *Metarhizium* spp. depends on the compounds contained in the medium. During the research stages, three types of cultivation for *Metarhizium* spp. isolates were used:

1. Cultivation in a liquid medium
2. Cultivation in a solid medium
3. Grown on the grain surface.

The Morphology of *Metarhizium*spp. isolates were identified following the methods [44]. The growth characteristics of isolates on the culture medium (PDA) were inspected continuously for 20 days, and their colony size, mycelial color, colony reverse, and color of conidial mass were visualized. Pure fungal cultures were obtained and examined microscopically (400× and 1000×) for characterization of morphological structures using lactophenol blue staining. The stained slides were observed under an 400× with a binocular light microscope (Leica, USA). The fungal isolates were identified based on the shape of the conidia and the arrangement of spores on the mycelia. Photomicrographs were obtained using a camera (CMEX) on the microscope (Euromex Oelphi-X, Observer series). When used for bioassays,

the number of conidia spotted in the center of plates and conidial concentrations were determined using a haemocytometer and the selected concentration of cell suspensions obtained by diluting in 0.01%.

**2.4. Production of fungi biomass during cultivation in liquid medium:** CSYE liquid medium (g/l: Glucose -40; KNO<sub>3</sub> - 10; KH<sub>2</sub>PO<sub>4</sub>- 5; MgSO<sub>4</sub>- 2; CaCl<sub>2</sub>- 0.05; Yeast extract - 2.3; Distill. water 500ml, pH-5.5-6) was used. The prepared nutrient medium was autoclaved and poured into 100 ml flasks. Fungi culture was added and incubated with shaking for 48 hours. As a result, a green mass appears on the surface of the medium.

**2.5. Growth of fungal biomass in solid medium:** Several solid agar mediums were used for the cultivation of *Metarhizium spp.* natural isolates :

*Potato Dextrose Agar (PDA):* Potatoes, infusion from: 200.0 g; Dextrose: 20.0 g, Agar: 15.0 g, distilled water (D/W):1000mL. Final pH (at 25°C) 5.6±0.2.

*Czapek Agar(original),* the composition of Cane sugar: 30 gm, monopotassium phosphate: 1 g, magnesium sulfate: 0.5 g, potassium chloride: 0.5 g, Iron sulfate: 0.01 g, distilled water (D/W): 1000 mL.

*Wrot Agar:* Malt extract 15.0 g; peptones 0.75g; maltose 12.75g; dextrin 2.75g; glycerin 2.35g; potassium phosphate monosubstituted 0.4g; ammonium chloride 1.0g; agar-agar 20.0g.

*Oat Agar medium* ( Oat – 100g; agar- 15g; dis. water -1000 mL; pH- 6.2).


A suspension was prepared from growing fungi conidia in distilled water, and 0.01% Tween 80, vortexed and the concentration of conidia was counted using a Neubauer hemacytometer.

**2.6. Growth of fungal biomass on the grain surface:** Growing in a solid medium on a rigid substrate takes up much space for aeration and grows according to the quality of the medium. Mass production of conidia varies depending on the various compounds in the medium. Fungi production is challenging; most industries want to grow in low-cost mediums. Plant grain wastes are the primary source for the surface cultivation and mass production of

fungi biomass. Oat bran for surface cultivation of fungi was used. The oat bran was simmered at 212°F (100 °C) for 10 min, then it was placed in the polyethene packets (37.5 x 26.3 cm) and sterilised at 270 °F (132 °C) for 60 min. After autoclaving, the oat bran media was weighed on 180 mg, and the wet media was different, from 20 % to 70%. The ready oat bran was inoculated with the liquid fungi culture grown on CSYE liquid medium. The polyethene packets were incubated at 24°C for 7 d. After 48-54 h incubation, culture appeared with green mycelium, and during the incubation, was weighed (growing from 1d 180mg to 7d 255mg). The obtained bioproduct can be stored for eight months ( Figure 2).

### 2.7. Bioassay for entomopathogenic activity of the *Metarhizium* spp isolates:

*Metarhizium* spp isolates' conidia produced on the oat-bran medium, harvested after 7d cultivation and used for inoculum to infect the pests. Four concentrations of suspension:  $7,2 \times 10^1$ ,  $7,2 \times 10^2$ ,  $7,2 \times 10^3$  and  $7,2 \times 10^4$  conidia/mL with adding 0.01% Tween 80 were prepared and vortexed. The plant leaves were washed with sterile distilled water. The insects' deutonymphs and adults of *A. fabae*, *A.nerii* and *T.urticae* (Table 1) were released on the leaf surface of host plants and directly sprayed with different concentrations (above mentioned) *Metarhizium* spp. conidia suspensions.



**Figure 2.** The growth and mycelium formation of *Metarhizium* spp. on the oat bran during seven days after inoculation

In control, the leaves were sprayed with sterile water. Each treatment was replicated three times. Thirty insect deutonymphs and adults for each treatment were tested. Observations for infected insects have lasted within seven days at 25± 2°C.

**Table 1.** Insect population from the Chui region, Bishkek, was used in screening tests.

| Host                                     | Hosts class  | Stage of host       | Month collected | Host plant             | Region        |
|--|--|---------------------|-----------------|------------------------|---------------|
| <i>Aphis fabae</i> (Black bean aphid)    | <i>Insecta</i> ,<br><i>Aphidoidea</i><br>(arthropods)    | adults, deutonymphs | December 2021   | <i>Cucumis sativus</i> | Chui, Bishkek |
| <i>Aspidiotus nerii</i> (Scale insect)   | <i>Insecta</i> ,<br><i>Diaspididae</i><br>(arthropods)   | deutonymphs         | March 2022      | <i>Citrus limon</i>    | Chui, Bishkek |
| <i>Tetranychus urticae</i> (Spider mite) | <i>Arachnida</i><br><i>Tetranychidae</i><br>(arthropods) | adults, deutonymphs | December 2021   | <i>Citrus limon</i>    | Chui, Bishkek |

**2.7.1. Assessment of mortality:** Treatments were recorded daily and continued for seven days. The leaf was counted every 24h under a stereo microscope (Euromex Oelphi-X, Observer series) in the Petri dishes and compared with control treatments. The mortality was calculated with Probit analysis (Probit analysis software in SPSS version 25).

**2.8. Antagonistic tests:** The inhibitory effect of *Metarhizium* spp. against *Fusarium oxysporum*, *Alternaria* spp. and *Rhizoctonia solani* was detected using a dual culture technique [45] on the YPG plate. One mycelial plug (6 mm diameter) of *Metarhizium* spp. was placed on a new YPG plate 1 cm away from the border of every Petri dish, and this was incubated at 25°C for two days. Then, a mycelial plug (6 mm diameter) of *Fusarium oxysporum*, *Alternaria* spp. and *Rhizoctonia solani* colony was put on the other side of the plate. Plates were incubated for 4-6 days until a clear inhibition zone was quantified at 25 ± 2 °C, and three replicates were tested. Control tests were also carried out using *Fusarium oxysporum*, *Alternaria* spp. and *Rhizoctonia solani* alone. The percentage inhibition of growth of the tested pathogen in the presence of antagonist isolates was calculated over control. The percentage inhibition of radial growth (PIRG) was calculated by using the formula developed by [45]: Inhibition %: = [(R - r)/R] × 100], where R = diameter of the

fungal colony in control,  $r$  = diameter of the fungal colony in dual inoculation. Interactions were assayed by giving ranking according to Bell's ranking scale [46], where R1= complete overgrowth of the pathogen by antagonist; R2= 75 % overgrowth; R3= 50% overgrowth; R4= growth inhibition at the line of contact; R5= pathogen over the growing antagonist.

**2.9. Molecular characterisation of *Metarhizium* spp:**For DNA extraction, fungal isolates were grown on Yeast Extract Agar at 25°C. Approximately 100 mg of mycelium was transferred to a screw-capped 2 mL microcentrifuge tube containing Lysing matrix C (MP Biomedicals) and DNA was extracted using the DNeasy Plant kit (Qiagen). Purified DNA was eluted in 100  $\mu$ L elution buffer AE and stored at -20 °C. The following DNA sequences were amplified from fungal DNA samples using standard Taq DNA polymerase (GoTaq (Promega). The internal transcribed spacer (ITS) region of nuclear rDNA was PCR amplified from genomic DNA by using ITS5 (GGAAGTAAAAGTCGTAACAAGG) and ITS4 (TCC TCC GCT TAT TGA TAT GC) primers [47].The generalised PCR protocol consisted of one initial denaturation step of 95°C for 2 min, 35 cycles of 45 sec at 95°C, 45 sec at the primer-specific annealing temperature and an elongation-step at 68 °C for an amplicon-specific elongation time, followed by a 5 min final elongation step at 68 °C. PCR product size was controlled by agarose gel electrophoresis, and PCR products were purified using the Qiaquick PCR purification kit (Qiagen). A commercial provider, Macrogen (Seoul, Korea), performed Sanger sequencing of PCR products using PCR and additional sequencing primers. Raw sequence data were combined into a single consensus sequence for each marker using version 6 of the MEGA software package [48]. The search for homologous sequences was done using Basic Local Alignment Search Tools at the National Center for Biotechnology Information.

**2.10. Statistical analysis:**Spore germination, sporulation date, and screening test results were analysed using one-way variance (ANOVA) using SPSS software version 25 statistical programs. Multiple mean comparisons were made using the Tukey HSD test when statistical differences were found between data sets ( $P=0.05$ ). Lethal concentration ( $LC_{50}$  and  $LC_{90}$ ) was

analysed using Probit analysis software in SPSS version 25. The dose responses of each replicate were checked for estimation of lethal time to kill 50% (LT<sub>50</sub>) of exposed larvae. Analysis of variances and means were separated by using least significant differences (LSD).

### 3. RESULTS AND DISCUSSION

**3.1. Morphological and cultural features of natural *Metarhizium anisopliae* isolates:** Table 2 shows the source and site of isolation of *Metarhizium spp.* strains. As results have shown, natural fungi isolates were found in dead insect bodies from the Lepidoptera and Apidae families.


Morphological characteristics of *Metarhizium spp.* isolates were proved by detailed microscopy studies of colonies grown on different solid mediums. When observed under x40 magnification in Petri dishes, all of these isolates produced typical greenish *Metarhizium* conidial masses on the back and front sides of the culture dish, which is an essential preliminary distinguishing feature for identifying this putative entomopathogen from other fungal relatives [49]. In addition, microscopic examination confirmed the partial identification of the isolates, as the isolates produced spores of medium and small size, as well as elliptical spores (Fig. 3).

**Table 2.** The source and place of new *Metarhizium spp.* isolation


| Collection number of <i>Metarhizium spp.</i> natural isolates | Source and site of isolation  | Maintained medium types |
|---|---|-------------------------|
| 1 TLK-1   | Cadavers of (Lepidoptera) larvae found in Torken village, Jalal-Abad region. 41°50'26.7"N73°09'13.7"E | PDA                     |
| 2 A-3   | Cadavers of insect from (Apidae) , found in Sokuluk village, Chuy region<br>42°51'02.7"N74°17'13.6"E  | Czapek agar             |
| 3 BZTK1   | Cadavers of insect from (Hemiptera), found in Sokuluk village , Chuy region                           | CSYE                    |
| 4 TLA-2   | Cadavers of (Lepidoptera) larvae found in Torken village, Jalal-Abad                                  | PDA                     |

### 3.2. Mycelium growth intensity and conidia production at the solid medium, per/ml:

The intensity of mycelium growth of *Metarhizium spp.* strains was observed on solid nutrient media used in the experiment at 25 °C. The diameter of the radial circle of mycelium growth of fungal isolates was measured daily. All studied strains grew intensively on oatmeal agar since, on the seventh day, the diameter of mycelium growth in strain TLK-1 reached  $7.5 \pm 0.96$  cm, in BZTK1 and TLA-2 -  $7.0 \pm 0.95$  cm, and A -  $3 \pm 0.91$  cm. A significant growth rate was observed on the PDA medium, forming a wide circle of mycelium growth, the diameter in all strains reaching up to  $5.0 \pm 0.95$  cm. The remaining two media observed no high fungal isolates' growth intensity (Fig. 4).



**Figure 3. A-F.** Conidiogenous cells and conidia of *Metarhizium* species. **A.** Mature phialides with developing conidia (TLK-1), Black bar = 189,55 $\mu$ m (40x); **B.** Mature conidia (TLK-1). Black bar = 720, 6 $\mu$ m (100x); **C.** Mature phialides (TLA-2); **D.** Black bar = 2023,02 $\mu$ m (100x) of mature phialides(TLA-2); **E.** Mature phialides with developing conidia(TLK-1); **F.** Distinctive ellipsoidal conidia in regular chains.




**Figure 4.** Mycelium growth of TLK-1, A-3, BZTK1 and TLA-2 strains on different medium for seven days. The colony diameters were measured and subjected to statistical analysis. Error bars represent standard deviations. Differences were analyzed by analysis of variance (ANOVA),  $P=0.05$


Conidia production at different solid media was assessed using an improved Neubauer hemocytometer under a light microscope, then calculated using the formula  $M = x \cdot 10^4 / (0.1 \cdot 1/400)$ , per/ml. Figure 5 shows a spore concentration of tested *Metarhizium* spp. isolates on Czapek, oat, Potato Dextrose, and Wort Agar. The isolates used solid nutritional formulations differently. Almost all isolates preferred oatmeal agar, produced high concentrations of spores; for example, the TLA-2 isolate produced  $7.2 \times 10^7$  conidia,  $\text{mL}^{-1}$  of suspension, BZTK1 isolate -  $5.3 \times 10^7$ , TLA-2 -  $6.3 \times 10^7$ . Of all the isolates studied, A-3 produced few conidia on almost all media. All isolates produced low conidia on Czapek's media, Wort agar, and PDA. Thus, a cheap medium containing oatmeal was chosen to obtain the conidia production.

**3.3. Mycelium growth intensity at different temperatures:** Culture of tested new *Metarhizium* spp. isolates were incubated at  $5^\circ$ ,  $15^\circ$ ,  $25^\circ$  and  $36^\circ$  C on a PDA medium. The results have demonstrated that the growth rate of fungal isolates at  $5^\circ$  C was slow, so during eight days of incubation, the diameter of TLA-2 and TLK-1 colonies reached  $3.5\text{-}3.8 \pm 0.96$  mm; the other two isolates grew more slowly than the strains mentioned above. At  $15^\circ$  C, the growth intensity was higher than at  $5^\circ$  C; for example, the diameter of TLA-2 and TLK-1 colonies made  $10.2 - 13.0 \pm 0.96$  mm for eight days of incubation. The temperature optimum for all tested *Metarhizium* spp. isolates were  $25^\circ$  C, their colony diameter reached from 12,


2 ± 0.95 mm to 24,5 ± 0.96 mm for eight days. The low growth rate was reported at 36 °C on the PDA medium (Figures 6 and 7).



**Figure 5.** Conidia production by *Metarhizium spp* isolates in a solid medium for seven days. Error bars represent standard deviations. Differences were analyzed by analysis of variance (ANOVA,  $P=05$ )



**Figure 6.** The growth intensity of *Metarhizium spp* isolates on a PDA medium for 8 days, at 5°C and 15 °C. Error bars represent standard deviations. Differences were analyzed by analysis of variance (ANOVA,  $P= 05$ )



25°C

32°C

**Figure 7.** The growth intensity of *Metarhizium* spp. isolates on a PDA medium for 8 days, at 25°C and 36 °C. Error bars represent standard deviations. Differences were analyzed by analysis of variance (ANOVA,  $P=0.05$ )

### 3.4. Insecticidal activity of *Metarhizium* spp. isolates' conidia produced on the oat-bran medium:


Considering that new isolates have preferred the oat-containing medium, oat bran was used to develop a low-cost formulation based on newly obtained *Metarhizium* spp. isolates.

Dried formulation with high content of tested isolates conidia was used for assessment of their insecticidal activity against adults and deutonymphs of *A. fabae*, *A. nerii* and *T. urticae* and using the doses  $7.2 \times 10^{-1}$ ,  $7.2 \times 10^{-2}$ ,  $7.2 \times 10^{-3}$ ,  $7.2 \times 10^{-4}$  conidia ml.

Figure 8 shows the insecticidal activity of different concentrations of the tested *Metarhizium* spp. TLK isolate towards the nymphalid stage of the pests mentioned above.

The dose of conidia  $7.2 \times 10^{-1}$  showed low pathogenicity for the nymphalid stage of all three pests, and the mortality ranged from  $22.0 \pm 0.96$  % for *T. urticae*;  $29.0 \pm 0.95$  % for *A. fabae* and  $40.0 \pm 0.95$  % *A. nerii*. While the dose  $7.2 \times 10^{-2}$  of TLK-1 has  $50 \pm 0.95$  % mortality in *T. urticae*,  $60 \pm 0.95$  % in *A. nerii* and  $60 \pm 0.95$  % in the population of *A. fabae*. A dose of  $7.2 \times 10^{-3}$  conidia has already shown a more effective activity concerning the larval stage of

all three pests; death has reached  $78 \pm 0.95 \%$  in *T.urticae* nymphalid population,  $90, 0 \pm 0.95 \%$  in the population of *A. fabae*,  $97,0 \pm 0.95 \%$  in *A.nerii*. The most active insecticidal dose was  $7.2 \times 10^{-4}$ , at which the TLK isolate caused  $97.0\% \pm 0.95 \%$  death in *T.urticae* and  $98,9.0\% \pm 0.95 \%$  in *A. fabae*  $99,9 \pm 0.95 \%$  in *A.nerii* nymphalid population.



**Figure 8.** The insecticidal activity of natural *Metarhizium spp.* TLK -1 isolate to the larvae stage of pests for seven days after applying different conidia doses. Error bars represent standard deviations. Differences were analyzed by analysis of variance (ANOVA,  $P= 05$ ).

The results of in vitro bioassay tests against sucking pests have revealed the  $LC_{50}$  and  $LT_{50}$  values of TLK-1 isolate (Table 3). The dose that caused 50% mortality in the population of used pests was different. The  $LC_{50}$  of this isolate for *A. fabae* adults was  $1 \times 10^6$  conidia  $\text{mL}^{-1}$ ; for *A.nerii* it was lower -  $7 \times 10^5$  conidia  $\text{ml}^{-1}$ ; while for adults *T.urticae* it was higher -  $13 \times 10^6$  conidia  $\text{ml}^{-1}$ . Such mortality occurred after 49.63 hours in adults of *A. fabae* ( $LT_{50}$ ), after 46.30 hours in *A.nerii* ( $LT_{50}$ ), and after 75.87 hours in deutonymph and adults of *T. urticae* ( $LT_{50}$ ). These data indicate that sucking pests have shown susceptibility to infection caused by the *Metarhizium spp.* isolate TLK-1, while the smallest dose of  $1 \times 10^6$  conidia could cause death in 50% of the *Aphis fabae* adult population after 49.63 hours. The shortest time during which death occurred in 50% of the population was noted in *A.nerii* deutonymphs at a dose of  $7 \times 10^5$  conidia  $\text{ml}^{-1}$ . After 46.30 hours was on the adults green- white mycelium, the lower mortality the time was adults of *T.urticae*, was hatched after 75.87 hours, green mycelium grew out of the cuticle of the adults.

**Table 3.** Mortality (LC<sub>50</sub> values) and (LT<sub>50</sub> values) for adults and deutonymphs of *A. fabae*, *A.nerii* and *T. urticae* and after inoculation of *Metarhizium* spp.


| Target pests                                      | LC <sub>50</sub> values (conidia ml) |           |    |                   |                                | LT <sub>50</sub> values (h) |          |    |                   |                      |
|---|--------------------------------------|-----------|----|-------------------|--------------------------------|-----------------------------|----------|----|-------------------|----------------------|
|   | n                                    | $\chi^2$  | df | Slope±S<br>E      | LC <sub>50</sub> conidia<br>ml | n                           | $\chi^2$ | df | Slope±S<br>E      | LT <sub>50</sub> (h) |
| <i>Aphis fabae</i> adult, and deutonymphs         | 1<br>0                               | 0.60      | 1  | -<br>2,53±1.5     | 1× 10 <sup>6</sup>             | 10                          | 1.71     | 1  | -<br>6,9±5.5<br>2 | 49.63                |
| <i>Aspidiotus nerii</i> adult, and deutonymphs    | 1<br>0                               | 1.02<br>3 | 1  | -3,2±1.7          | 7× 10 <sup>5</sup>             | 10                          | 0.86     | 1  | -<br>9,5±5.6<br>8 | 46.30                |
| <i>Tetranychus urticae</i> adult, and deutonymphs | 1<br>0                               | 0.30      | 1  | -<br>6,4±4.4<br>2 | 13× 10 <sup>6</sup>            | 10                          | 0.6      | 1  | -<br>19±16.1      | 75.87                |

LC<sub>50</sub> and LT<sub>50</sub> values and 95% confidence limits (CL) expressed as conidia (×10<sup>4</sup>) (Proc Probit (SPSS). LT<sub>50</sub> values for mortality were estimated by analysis (Proc Probit (SPSS) data for >72 hours for *A. fabae*, *A.nerii* and *T. urticae*, corresponding to the last day of incubation in bioassay >10 days.

The optimal climatic conditions created during the experiment (humidity 60-65%, temperature 25<sup>0</sup> C) ensured the fungus's pathogenicity and its conidia could form hyphal bodies on the cuticle of infected pests. At the same time, hyphae produced different pigments on the cuticles of different insects. On the cuticle, deutonymphs of *A.nerii* were white-green hyphae (Fig.9). On the cuticle of *Aphis fabae* adults the fungus produced grey-green hyphae (Fig.10). At the adult stage of *T. urticae*, the fungal isolate had produced dark green hyphae. According to [50], insect tolerance to infection with *Metarhizium* spp. may change depending on environmental stresses. The production of fungal pigment depends on many factors.



**Figure 9.** Mycosis on the cuticle of *Aspidiotus nerii* produced by *Metarhizium* spp. TLK-1 isolate.




**Figure 10.** Mycosis on the cuticle of *Aphis fabae* adults produced by *Metarhizium* TLK-1 isolate.

**3.5. Biofungicide activity of new natural *Metarhizium* isolates:** Three isolates, TLK-1, TLA-2 and BZTK-1 have exhibited biofungicide activity in two pathogen fungi like, *Fusarium oxysporum* and *Alternaria spp.* According to Bell's ranking scale [46], interactions between *Metarhizium* isolates and plant pathogens were assayed by ranking R4= growth inhibition at the line of contact. The TLK-1 isolate has shown significant activity against the pathogenic fungus *Fusarium oxysporum*, strain Fus:85 (Fig.11D) and relatively strong activity against *Alternaria spp* strain1.3 (Fig. 11F). The zone of inhibition of these pathogens from the


antagonistic effect of the TLK-1 isolate is visible in the figures, which was 1.8-2.0 mm on the fifth day of the joint incubation of these two types of fungi.

Other strains: TLA-2 and BZTK-1 also showed antagonistic activity against pathogenic *Fusarium oxysporum* and *Alternaria* sp. strains; their zone of inhibition was 1.7-1.8 mm (Fig.11 G, I, J, L, M and O). At the same time, these three isolates of *Metarhizium* did not show antagonistic activity against the pathogen *Rhizoctonia solani* (Fig.11.E, H, K and N). Colonies of these two different species of fungi grew together when co-cultivated.

Selected fungal isolates were subjected to DNA sequencing for identification. The ITS1 and ITS4 DNA fragments of select fungal isolates were amplified by PCR using the primer pairs ITS1 (CTTGGTCATTTAGAGGAAGTAA) and ITS4 (CAGGAGACTTGTACACGGTCCAG) and genomic DNA as the template. Based on their morphological characteristics, including the production of ellipsoidal conidia, and molecular characteristics (ITS, partial 18S [SSU rDNA] and EF1- $\alpha$  sequences), the isolates were identified as *Metarhizium anisopliae* (belonging to Clade E from Asia (Figure 12)).



**Figure 11.** Dual plate assays is showing an antagonistic activity of natural new *Metarhizium* isolates. **A-** Control, *Fusarium oxysporum*( strain Fus.85). **B-** Control, *Rhizoctonia solani*(strain Rhiz.2.1.1). **C-** Control, *Alternaria*spp.(strain Alter.1.3).**D-** Antagonistic activity of TLK-1 on Fus.85. **E-** Antagonistic activity of TLK-1 on Rhiz.2.1.1.; **F-** Antagonistic activity of TLK-1 on Alter.1.3. **G.** Antagonistic activity of TLA-2 on FUS.85. **H.** Antagonistic activity of TLA-2 on RHIZ.2.1.1. **I.** Antagonistic activity of TLA-2 on ALTER.1.3. **J.** Antagonistic activity of BZTK-1 on FUS.85. **K.** Antagonistic activity of BZTK-1 on RHIZ.2.1.1. **L.** Antagonistic activity on ALTER.1.3. **M.** Antagonistic activity of A-3 on FUS.85. **N.** Antagonistic activity of A-3 on RHIZ.2.1.1. **O.** Antagonistic activity of A-3 on ALTER.1.3



**Figure 12.** Neighbors-joining phylogenetic tree based on 18SrRNA gene sequences showing the position of new local strains TLA-2 KTMU, BZTK-1 KTMU and TKL-1 KTMU among other closely related *Metarhizium anisopliae* species from the database.

Fungi in the genus *Metarhizium* (Hypocreales: Clavicipitaceae) are considered a relevant biological agent, that affect about 200 types of insects from orders like Lepidoptera, Homoptera, Hymenoptera, Coleoptera, and Diptera [ 5,6, 51]. *Metarhizium* fungi have an advantage in rapidly producing infectious conidia when incubated on inexpensive nutrient media to produce the commercial mass production of highly effective bioproducts with broad ranges of activities [ 2,3,18]. A multifunctional lifestyle of *Metarhizium.spp.* species is recorded by several studies, except as insect pathogens, these fungi can benefit in inhibiting plant pathogens, in growth promotion and protecting against stresses [31, 32]. The studies have noted that native fungal isolates may offer a better alternative than imported strains. They may be better adapted to kill local insects and survive local environmental conditions as vectors for biological control [52]. The level of this fungus virulence depends on environmental conditions, and biotic and abiotic factors can affect the effectiveness of their application. Therefore, it is necessary to consistently screen the local strains to select pathogenic pests to increase the chances of obtaining effective microbial biopesticides.

In this study, in vitro experimental screenings were carried out to identify and select local *Metarhizium* strains with several properties, namely, bioinsecticidal and biofungicidal activity.

To cause mycosis or infectious disease to harmful insects in natural conditions, in addition to relevant abiotic factors, these fungi must have a sufficient dose of their conidia. Studies have shown that the most pathogenic fungal isolates produced spores on the corpses of larvae with a minimum ( $3.2 \times 10^6$  spores ml<sup>-1</sup>) to a maximum ( $5.4 \times 10^7$  spores ml<sup>-1</sup>) concentration. Considered to be rapid sporulation, with a large amount of conidia on corpses, it can cause increased horizontal pest infestation due to rapid self-propagation [53]. These conidia can attach to insects' covering tissues in a high dose. And when they stand on the cuticle cover, they secrete enzymes that dissolve insects' cuticles. This study selected various compositions of cheap nutrient media to obtain a high titer of this entomopathogenic fungus, which provided the fungi with intensive conidia formation. According to studies, among *Metarhizium*, there are species with a narrow spectrum of activity; for example, *Metarhizium acridum* is a specific locust pathogen [54, 55]. In our studies, the local strains have shown an entomopathogenic activity against sucking pests from various taxonomic groups, for example, against *Aphis fabae* (Insecta, Aphidoidea), *Aspidiotusnerii* (Insecta, Diaspididae) and *Tetranychusurticae* (Arachnida, Tetranychidae), which can be recommended as a universal biological control for the regulation of the number of the above economically important greenhouse pests, as well as in open ground conditions (Figure 7, Table 3).

Other authors have also obtained data showing the efficacy of *Metarhizium* to flies (Insecta: Diptera: Tephritidae); for example, [56] have shown the effect of *Metarhizium anisopliae* inoculation even on the mating behaviour of three fruit fly species: *Ceratitis cosyra*, *C. fasciventris* and *C. capitata*. Mating duration was reduced in males treated with these fungi. Some research has shown sub-lethal and lethal effects of *M. anisopliae* on the cotton bug *Oxycarenushyalinipennis*, showing that this entomopathogenic fungus can have an effect not only on the parent generation but on the next generation (F1) [57].

Most entomopathogenic fungi used as biocontrol agents require high humidity to germinate their spores. However, high humidity can also promote the growth of phytopathogens, causing rot, powdery mildew and others. Therefore, in recent years, research has focused on identifying entomopathogens that have dual effects on both pests and pathogens to resolve this issue.

The dual control potential of *Metarhizium anisopliae* SD3 isolate for pest and plant pathogens was reported [58, 59]. This fungus has shown virulence against green peach aphids (*Myzus persicae*) and antifungal activity to grey mould (*Botrytis cinerea*). The most virulence was observed in the simultaneous treatment with blastospores and cultural filtrate.

Our studies also confirmed that the studied local strains showed not only insecticidal activity but also biofungicidal activity against plant pathogens, strong activity was shown against the phytopathogenic fungus *Alternaria*, then against *Fusarium oxysporum*. At the same time, these entomopathogenic fungi do not show any activity *Rhizoctonia solani* (Figure 10). Many researchers have reported that *Metarhizium* spp. having virulence against such plant pathogens as *B. cinerea*, *Fusarium* spp., *Phytophthora megasperma*, *Podosphaera fuliginea*, *Pythium* spp., *Rhizoctonia solani*, *Sphaerotheca fuliginea* and *Verticillium dahliae* [58,60].

Our study is the first report of the dual activity of native entomopathogenic fungi against dangerous sucking pests from different taxonomic groups and against essential pathogens of many crop species such as *Fusarium oxysporum* and *Alternaria* spp. These new local strains can be recommended for bio-formulation development to protect plants from diseases and as bioinsecticides and bioacaricides against harmful mites and insects.

#### **4. Conclusions**

Biological control agents (BCAs) as an alternative to synthetic products remain the focus of researchers and several commercial ventures worldwide. These alternative products are already integrated into conventional growing methods, many of which are available to farmers.

Our results are significant as they point the way for other Integrated Pest Management (IPM) methods to be combined with these fungi to improve the effectiveness of insect control. We have identified and begun preliminary characterization of local isolates of *Metarhizium spp.*, which have the potential to be used in complex protection systems against various important pests, in particular suckers. The found isolates can be simultaneously used against widespread pathogenic fungi, essential for developing and producing inexpensive biological formulas based on these fungi. The results obtained on the optimization of cheap nutritional compositions for producing fungal biomass with a high titer of conidia open up new opportunities for creating low-cost, high-efficiency biopreparations with a multifunctional spectrum of action.

#### **Declarations**

#### **Availability of data and materials**

A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal."

#### **Ethics approval and consent to participate**

Not applicable

#### **REFERENCES**

1. Roberts DW, Lege SRJ. *Metarhizium*spp cosmopolitan insect-pathogenic fungi: mycological aspects. *Microbiol.* 2004; 54:1–70. [https://doi.org/10.1016/S0065-2164\(04\)54001-7](https://doi.org/10.1016/S0065-2164(04)54001-7).
2. Faria MR, Wraight SP. Mycoinsecticides and mycoacaricides: A comprehensive list with worldwide coverage and international classification of formulation types. *Biol Control.* 2007; 43:237–256. <https://doi.org/10.1016/j.biocontrol.2007.08.001>.
3. Arthurs S, Dara SK. Microbial biopesticides for invertebrate pests and their markets in the United States. *J InvertebrPathol.* 2019; 165:13–21. <https://doi.org/10.1016/j.jip.2018.01.008>.
4. Jitendra Dhaker JK, Kavia A, Sen P, Kaushal N, Datta S. Biomass Production of Entomopathogenic Fungi using various Agro Products in Kota Region, India. *I. Res. J. Biological Sci.* 2012;<http://www.isca.me/IJBS/Archive/v1/i4/3.ISCA-JBS-2012-088>.
5. Driver F, Milner RJ, Trueman JW. A taxonomic revision of *Metarhizium* based on a phylogenetic analysis of rDNA sequence data. *Mycol Res.* 2000;104(2):134–150. <https://doi.org/10.1017/S0953756299001756>.
6. Khater HF. Prospects of botanical biopesticides in insect pest management. *Pharmacologia*; 2012.

7. Zimmermann G. The 'Galleria bait method' for detection of entomopathogenic fungi in soil. *Journal of Applied Entomology*. 1986; 102:213-215.
8. Iwasaki H, Tokiwa T, Shiina M, Asami Y, Shiomi K, Omura S. et al. *Metarhizium aciculare* sp. nov. for euvesperins A and B producing *Metarhizium* strains. *Mycoscience*.2019;313-318. <https://doi.org/10.1016/j.myc.2019.06.005>.
9. Stephen AR, Ryan MK. Species limits, phylogeography and reproductive mode in the *Metarhizium anisopliae* complex. *Invertebrate Pathology*.2017; 60-66. <https://doi.org/10.1016/j.jip.2017.05.008>.
10. Leger SRJ. Biology and Mechanisms of Insect-Cuticle Invasion by Deuteromycete Fungal Pathogens. In: Beckage, N.E., Thompson, S.N. and Federici, B.A., Eds., *Parasites and Pathogens of Insects*. New York. Academic Press Inc;1993.
11. Zhang LB, Feng MG. Antioxidant enzymes and their contributions to biological control potential of fungal insect pathogens. *Appl Microbiol Biotechnol*. 2018; 102:4995–5004. <https://doi.org/10.1007/s00253-018-9033-2>.
12. Tong SM, Feng MG. Insights into regulatory roles of MAPK-cascaded pathways in multiple stress responses and lifecycles of insect mycopathogens. *Appl Microbiol Biotechnol*. 2019;103:577–587. <https://doi.org/10.1007/s00253-018-9516-1>.
13. Rogerio BL, Marcos F. Influence of two formulation types and moisture levels on the storage stability and insecticidal activity of *Beauveria bassiana*. *Biocontrol Science and Technology*. 2019;29:437-450.
14. Hedgecock S, Moore D, Higgins P, Prior C. Influence of moisture content on temperature tolerance and storage of *Metarhizium favoviride* conidia in an oil formulation. *Biocontrol Sci Technol*.1995; 5:371–377. <https://doi.org/10.1080/09583159550039828>.
15. Kim JS, Lee SJ, Lee HB. Enhancing the thermotolerance of entomopathogenic *Isaria fumosorosea* SFP-198 conidial powder by controlling the moisture content using drying and adjuvants. *Mycobiology*.2014;59–65.
16. Moller-kogler Zimmermann G. Zur autbewahrungEntomogenerpilzkulturen. *Entomophaga*.1980;25:301-311.
17. Neves PM, Alves SB. External events related to the infection process of *Cornitermes cumulans* (Kollar) (Isoptera: Termitidae) by the entomopathogenic fungi *Beauveria bassiana* and *Metarhizium anisopliae*. *Neotrop Entomol*. 2004; 33:51–56. <https://doi.org/10.1590/S1519-566X2004000100010>
18. Mascarin GM, Jaronski ST. (The production and uses of *Beauveria bassiana* as a microbial insecticide. *World J Microbiol Biotechnol*.2016; 32:177. <https://doi.org/10.1007/s11274-016-2131-3>
19. David MB, Markus K, Hamadi IB, Sunday E, Nguya KM. Susceptibility of developmental stages of *Tetranychus urticae* (Acari: Tetranychidae) to infection by *Beauveria bassiana* and *Metarhizium anisopliae* (Hypocreales: Clavicipitaceae). *Tropical Insect Science*. 2014; 190 – 196. <https://doi.org/10.1017/S1742758414000381>.

20. Alves SB, Tamai MA, Rossi LS, Castiglioni E. *Beauveria bassiana* pathogenicity to the citrus rust mite *Phyllocoptrutaoleivora*. *Experimental and Applied Acarology*. 2005; 37:117–122.
21. Rodríguez M, Gerding M, France A. Entomopathogenic fungi isolates selection for egg control of tomato moth *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae). *Agric Tec Chile*. 2006; 66:151-158.  
<http://www.bioline.org.br/abstractat06016>.
22. SabbourMM, Singer SM. Evaluations of two *Metarhizium* varieties against *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Egypt. *IJSR*.2014;3:1983–1987.
23. Atef Sayed MM, Christopher AD. Virulence of Some Entomopathogenic Fungi Isolates of *Beauveria bassiana* (Hypocreales: Cordycipitaceae) and *Metarhizium anisopliae* (Hypocreales: Clavicipitaceae) to *Aulacaspistubercularis* (Hemiptera: Aspidiotusnerii ) and *Iceryaseychellarum* (Hemiptera: Monophlebidae) on Mango. *Crop Journal of Economic Entomology*. 2019; 1–13.<https://doi.org/10.1093/jee/toz187>
24. Ahmed SA, El-Saedi AA, Ahmed HM, Sayed AM. Biological aspects and life tables of the coccinellids predator *Rodolia cardinalis* and its prey *Iceryaseychellarum* (Homoptera: Margarodidae). *The Second International Conference of Egyptian Society of Economic Entomology*. Egypt: Egyptian National Agricultural Library.2007;59–77.
25. Sayed AMM. Influence of certain bio-agents and climatic changes on the population density of the white mango scale insect, *Aulacaspistubercularis*. *J Agric Res*.2012; 90: 607–624.
26. Sayed AMM, Behle RW. Evaluating a dual microbial agent biopesticide with *Bacillus thuringiensis* var. *kurstaki* and *Beauveria bassiana* blastospores. *Biocontrol Sci. Technol*. 2017;27: 461–474.
27. González-Mas N, Sánchez-Ortiz A, Valverde-García P, Quesada-Moraga E. Effects of endophytic entomopathogenic ascomycetes on the life-history traits of *Aphis gossypii* Glover and its interactions with Melon plants. *Insects*. 2019;10:165. <https://doi.org/10.3390/insects10060165>.
28. Hong SI, Kim K. Selection of entomopathogenic fungi for aphid control. *J BiosciBioeng*. 2007;104(6):498–505.<https://doi.org/10.1263/jbb.104.498>.
29. Mukherjee A, Debnath P, Ghosh SK, Medda PK. Biological control of papaya aphid (*Aphis gossypii* Glover) using entomopathogenic fungi. *Springer Link*. 2020;1-10.<https://doi.org/10.1007/s42535-019-00072-x>.
30. Milner RJ. Prospects for biopesticides for aphid control. *BioControl*.1997;42(1);227–239. <https://doi.org/10.1007/BF02769900>.
31. Behie SW, Zelisko PM, Bidochka MJ. Endophytic insect-parasitic fungi translocate nitrogen directly from insects to plants. *Science*. 2012;1576–1577.<https://doi.org/10.1126/science.1222289>.
32. Liao X, O'Brien TR, Fang W, Leger RJS. The plant beneficial effects of *Metarhizium* species correlate with their association with roots. *Appl Microbiol Biotechnol*.2014; 98:7089–7096.<https://doi.org/10.1007/s00253-014-5788-2>.
33. Enrique GP, María AOA, Elihú B. The entomopathogenic fungus *Metarhizium anisopliae* enhances *Arabidopsis*, tomato, and maize plant growth. *Plant Physiology and Biochemistry*. 2022; 34-43.<https://doi.org/10.1016/j.plaphy.2022.02.008>.

34. Krell V, Jakobs-Schoenwandt D, Vidal S, Patel AV. Encapsulation of *Metarhizium brunneum* enhances endophytism in tomato plants. *Biol Control*. 2018;116: 62–73. <https://doi.org/10.1016/j.biocontrol.2018.116.62-73>.
35. Ahmad I, Jimenez-Gasco MdM, Luthe DS, Barbercheck ME. Systemic colonization by *Metarhizium robertsii* enhances cover crop growth. *J Fungi*. 2020; 6:64. <https://doi.org/10.3390/jof6020064>.
36. Ahmad I, Jimenez-Gasco MdM, Luthe DS, Shakeel SN, Barbercheck ME. Endophytic *Metarhizium robertsii* promotes maize growth, suppresses insect growth, and alters plant defense gene expression. *Biological Control*. 2020; 144:104167. <https://doi.org/10.1016/j.biocontrol.2019.104167>.
37. Flonc B, Barbercheck M, Ahmad I. Observations on the relationships between endophytic *Metarhizium robertsii*, *Spodoptera frugiperda* (Lepidoptera: Noctuidae), and maize. *Pathogens*. 2021; 10:713. <https://doi.org/10.3390/pathogens10060713>.
38. Russo ML, Pelizza SA, Vianna MF, Allegrucci N, Cabello MN, Toledo AV. et al. Effect of endophytic entomopathogenic fungi on soybean *Glycine max* (L.) Merr. growth and yield. *J King Saud Univ—Sci*. 2018; 31:728–736. <https://doi.org/10.1016/j.jksus.2018.04.008>.
39. Liu SF, Wang GJ, Nong XQ, Liu B, Wang MM, Li SL. et al. Entomopathogen *Metarhizium anisopliae* promotes the early development of peanut root. *Plant Prot Sci*. 2017;53:101–107. <https://doi.org/10.17221/49/2016-PPS>.
40. Maksim T, Marsel RK, Natalia S, Oksana GT, Olga Y. et al. (Can Potato Plants Be Colonized with the Fungi *Metarhizium* and *Beauveria* under Their Natural Load in Agrosystems. *Microorganisms*. 2012;9(7):1373. <https://doi.org/10.3390/microorganisms9071373>
41. Greenfield M, Gomez-Jimenez MI, Ortiz V, Vega FE, Kramer M, Parsa S. *Beauveria bassiana* and *Metarhizium anisopliae* endophytically colonize cassava roots following soil drench inoculation. *Biol Control*. 2016;95:40–48. <https://doi.org/10.1016/j.biocontrol.2016.01.002>.
42. Jaber LR, Alananbeh KM. Fungal entomopathogens as endophytes reduce several species of *Fusarium* causing crown and root rot in sweet pepper (*Capsicum annuum* L.). *Biol Control*. 2018; 126: 117–126. <https://doi.org/10.1016/j.biocontrol.2018.08.007>.
43. Sasan RK, Bidochka MJ. The insect-pathogenic fungus *Metarhizium robertsii* (Clavicipitaceae) is also an endophyte that stimulates plant root development. *Am J Bot*. 2012; 99: 101–107. <https://doi.org/10.3732/ajb.1100136>
44. Fernandes ÉKK, Keyser CA, Chong JP, Rangel DEN, Miller MP, Roberts DW. Characterization of *Metarhizium* species and varieties based on molecular analysis, heat tolerance and cold activity. *Journal of Applied Microbiology*. 2010; 115–128. <https://doi.org/10.1111/j.1365-2672.2009.04422.x>.
45. Skidmore AM, Dickinson CH. Colony interactions and hyphal interference between *Septoria nodorum* and phylloplane fungi. *Transactions of the British Mycological Society*. 1976;57-64. [https://doi.org/10.1016/S0007-1536\(76\)80092-7](https://doi.org/10.1016/S0007-1536(76)80092-7).

46. Bell DK, Wells HD, Markham CR. In vitro antagonism of *Trichoderma* species against six fungal plant pathogens. *Phytopathology*. 1982; 379-382. <https://doi.org/10.1094/Phyto-72-379>.
47. White TJ. Amplification and Direct Sequencing of Fungal Ribosomal RNA Genes for Phylogenetics. In: *PCR Protocols, a Guide to Methods and Applications*; 315-322. 1990.
48. Tamura K, Stecher G, Peterson D. et al. MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution*; 2725–2729. 2013. <https://doi.org/10.1093/molbev/mst197>.
49. Du ZY, Krzysztof Z, Natalie VP, Nathaniel EO, Christoph B, Gregory MB. Algal-fungal symbiosis leads to photosynthetic mycelium. *eLife Science*. 2019; <https://doi.org/10.7554/eLife.47815>.
50. Wang C, Duan Z, Leger SRJ. MOS1 osmosensor of *Metarhizium anisopliae* is required for adaptation to insect host hemolymph. *Eukaryot Cell*. 2008; 7:302–309. <https://doi.org/10.1128/EC.00310-072008>.
51. Jitendra Dhaker JK, Kavia A, Sen P, Kaushal N, Datta S. Biomass Production of Entomopathogenic Fungi using various Agro Products in Kota Region, India. *I. Res. J. Biological Sci*. 2012; <http://www.isca.me/IJBS/Archive/v1/i4/3.ISCA-JBS-2012-088>
52. Bilgo E, Lovett B, Leger RJ, Sanon A, Dabiré RK, Diabaté A. Native entomopathogenic *Metarhizium* spp. from Burkina Faso and their virulence against the malaria vector *Anopheles coluzzii* and nontarget insects. *Parasites and Vectors*. 2018; 11:11–6.
53. Conceschi RM, Moral AR, D'Alessandro PC, Demetri BGC. Transmission potential of the entomopathogenic fungi *Isaria fumosorosea* and *Beauveria bassiana* from sporulated cadavers of *Diaphorinacitri* and *Toxopteracitricida* to uninfected *D. citri* adults. *Biol Control*. 2016; 1–11. <https://doi.org/10.1007/s10526-016-9733-4>.
54. Gao QA, Jin K, Ying SH, Zhang YJ, Xiao GH, Shang YF, Duan ZB, Hu XA, Xie XQ. et al. Genome sequencing and comparative transcriptomics of the model entomopathogenic fungi *Metarhizium anisopliae* and *M. acridum*. *PLoS Genet*. 2011; 7(1). <https://doi.org/10.1371/journal.pgen.1001264>.
55. Pattermore JA, Hane JK, Williams AH, Wilson BAL, Stodart BJ, Ash GJ. The genome sequence of the biocontrol fungus *Metarhizium anisopliae* and comparative genomics of *Metarhizium* species. *BMC Genomics*. 2014; <https://doi.org/10.1186/1471-2164-15-660>
56. Dimbi S, Nguya KM, Ekesi S. Effect of *Metarhizium anisopliae* inoculation on the mating behavior of three species of African Tephritid fruit flies, *Ceratitis capitata*, *Ceratitis cosyra* and *Ceratitis fasciventris*. *Biological Control*. 2009; 111-116.
57. Shaukat RF, Freed S, Ahmed R, Raza M, Naeem A. Virulence and trans-generational effects of *Metarhizium anisopliae* on *Oxycarenushyalinipennis*. *Pest Manag Sci*. 2023; <https://doi.org/10.1002/ps.7568>.
58. Shin TY, Bae SM, Kim DJ, Yun HG, Woo SD. Evaluation of virulence, tolerance to environmental factors and antimicrobial activities of entomopathogenic fungi against two-spotted spider mite, *Tetranychus urticae*. *Mycoscience*. 2017; 58:204-12.
59. Lozano-Tovar MD, Ortiz-Urquiza A, Garrido-Jurado I, Trapero Casas A, Quesada-Moraga E. Assessment of entomopathogenic fungi and their

extracts against a soil-dwelling pest and soilborne pathogens of olive. *Biol Control*. 2013;67:409-20.

60. VegaFE, Goettel MS, Blackwell M, Chandler D, Jackson MA, Keller S, Koike M, Maniania NK, Monzón A, Ownley BH. et al. Fungal entomopathogens: new insights on their ecology. *Fungal Ecol*. 2009; 2:149-59.

UNDER PEER REVIEW