

## Original Research Article

# Effect of Sulphur Application on Yield, Uptake and Quality of Chickpea (*Cicer arietinum* L.)

### ABSTRACT

A field experiment was conducted at Pulse Research Station in Dr.Panjabrao Deshmukh Krishi Vidyapeeth, Akola, Maharashtra from the period of *Rabi* 2020-21 to 2022-23 to study the effect of sulphur application on yield, uptake and quality of Chickpea (*Cicer arietinum* L.). The experiment was laid out in Randomized block design (RBD) with 9 treatments and 3 replications. The study revealed that the incorporation of sulphur application is significantly influenced the yield, uptake and quality. The highest seed yield (24.18 q ha<sup>-1</sup>) and straw yield (30.16 q ha<sup>-1</sup>) were recorded with the application of S @30 kg ha<sup>-1</sup> through Bentonite sulphur along with RDF and it was found significantly superior over all the treatments. Similarly maximum N, P, K, S and micronutrient uptake and improved quality were observed with application of @ 30 kg ha<sup>-1</sup> through Bentonite sulphur along with RDF followed by the treatment of application of S @ 30 kg ha<sup>-1</sup> through Gypsum + RDF.

**Key words:** Bentonite Sulphur, Gypsum, Yield, Uptake, Quality, Chickpea.

### 1. INTRODUCTION

Chickpea (*Cicer arietinum* L.) is the most important grain legume in the world after dry beans and dry peas. Its cultivation is mainly confined to Asia with 90 per cent of the global area and production. Besides Asia it is also grown in North and Central America, the Mediterranean region, the west Asia and North Africa (WANA) region and Eastern Africa. India is top pulse producing country in the world. Among pulses chickpea ranks third in the world. The total area under pulse in India has increased from 19 million hectares in 1950-51 to 29 million hectares in 2016-17. The total pulse production in India was 18.43 million tonnes (2014-15), 19.41 million tonnes (2015-16) and 22.95 million tonnes (2016-17) (Anonymous, 2017 (b)). India rank first in Chickpea production. The largest gram producing state in India with respect to area are Madhya Pradesh, Maharashtra Uttar Pradesh, Andhra Pradesh, Rajasthan, Haryana and Karnataka. Madhya Pradesh rank first in Chickpea production. Maharashtra second major Chickpea producing state after Madhya Pradesh. In Maharashtra area under Chickpea crop was 13.53 lakh hectares with the production of 11.80 lakh tonnes with an average productivity of 872 kg ha<sup>-1</sup>. Therefore, Maharashtra contributes 18.36 per cent share in area and 20.03 per cent share in production of chickpea in India.

Sulphur is the 4<sup>th</sup> major essential plant nutrient after N, P and K because of its role in the synthesis of proteins, vitamins, enzyme and flavoured compounds in plant. Its amount required by the plant is even higher than phosphorus and comes after N and K. About 90% of plant sulphur is present in amino acids viz., Methionine, Cystine, and Cysteine. These amino acids are the building blocks of

protein. Sulphur is associated with production of crops of superior nutritional and market quality. Sulphur deficiencies have been reported from over 70 countries worldwide including India. Sulphur is becoming deficient in soil due to use of high grade S free fertilizers, cultivation of high yielding varieties and lack of industrial activity (Scherer, 2009).

The sulphur status of Indian as well as Maharashtra soil is changing rapidly because of heavy removal of the nutrients from the soil under multiple cropping system with high yielding fertilizer responsive varieties. Use of sulphur containing fertilizers in soil will be helpful to the farmers in improving growth of plant, increasing protein content, yield of chickpea. Sulphur, in chickpea, mainly influences the protein content. Sulphur helps towards conversion of nitrogen into protein in pulse crops. Sulphur also improves the S containing amino acid in crop and thus enhances the protein content (Das *et al.*, 2016).

In Maharashtra state, isolated attempts were made to work out a critical level of sulphur in the soils and plants. In Indian soils sulphur deficiency has been noticed 32.9 % (Shukla *et al.*, 2016), while in Maharashtra sulphur deficiency recorded to the extent of 37.48 % while in Vidarbha it was noticed 25.76 % (Katkare *et al.*, 2017). So it is essential to evaluate the effect of sulphur application on soil fertility, yield, nutrient uptake and quality of chickpea in black soil. The information generated through this investigation will be helpful to apply the sulphur fertilizers to chickpea crop in Vertisols.

Keeping all the above facts in view, the present investigation was undertaken with the objectives to study the "Effect of sulphur application on yield, uptake and quality of chickpea (*Cicer arietinum* L.).

## 2. MATERIALS AND METHODS

The experiment was conducted to study the sulphur requirement of Chickpea crop. The soil was analysed for soil properties, plant and seed samples were analysed for nutrient uptake and quality of chickpea crop.

The field experiment was conducted at Pulse Research Unit, Dr.Panjabrao Deshmukh Krishi Vidyapeeth, Akola during rabi season 2020-21 to 2022-23. The site is situated at the subtropical region at 22° 42' North latitude and 77° 02' East longitude and at an altitude of 307.42 m above mean sea level. Initial composite soil sample was collected from the experimental site and analyzed for soil properties. The experimental site was slightly alkaline in reaction (7.96), non-saline (0.24 dS m<sup>-1</sup>), medium in organic carbon (5.28 g kg<sup>-1</sup>), calcareous in nature (6.87%), low in available N (188.16 kg ha<sup>-1</sup>), low in available P (13.65 kg ha<sup>-1</sup>), very high in available K (581.2 kg ha<sup>-1</sup>), deficient in available S (9.82 mg kg<sup>-1</sup>) and sufficient in DTPA - Zn, Fe, Cu and Mn (mg kg<sup>-1</sup>). Treatment details were as T<sub>1</sub> was Absolute Control, T<sub>2</sub> is S-free RDF (N, P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O through Urea, DAP, MOP), T<sub>3</sub> was RDF (N, P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O through Urea, SSP, MOP), T<sub>4</sub> was T<sub>2</sub> + S @ 10 kg ha<sup>-1</sup> through Bentonite-Sulphur, T<sub>5</sub> was T<sub>2</sub> + S @ 20 kg ha<sup>-1</sup> through Bentonite-Sulphur, T<sub>6</sub> was T<sub>2</sub> + S @ 30 kg ha<sup>-1</sup> through Bentonite-

Sulphur, T<sub>7</sub> was T<sub>2</sub> + S @ 10 kg ha<sup>-1</sup> through Gypsum, T<sub>8</sub> was T<sub>2</sub> + S @ 20 kg ha<sup>-1</sup> through Gypsum, T<sub>9</sub> was T<sub>2</sub> + S @ 30 kg ha<sup>-1</sup> through Gypsum. The recommended dose of NPK & S fertilizer used were 20:60:40:30. The plant samples were also collected at maturity stage. The samples were air dried and then oven dried at 105°C. The treatment wise samples were ground by using grinding mill and stored with proper labelling in brown paper bags. The powdered samples were used for the chemical analysis of N, P, K, S and micronutrient content which were estimated using modified Kjeldahl's method (AOAC, 1995), phosphorous by Ammonium molybdate vanadate (Chapman pratt 1962), potassium by using Flame Photometer (Hanway and Heidal 1952), sulphur was estimated from di-acid extract turbidimetrically using Spectrophotometer (Chesnin and Yien 1950) and micro nutrients by using AAS (Issac and Kerber 1971). The test of statistical significance of the experimental data was carried out as per procedure described by (Panse and Sukhatme 1985).

### 3. RESULTS AND DISCUSSION

**Yield of Chickpea:** The data pertaining to seed yield of chickpea was influenced significantly (Table 1). The significantly higher seed yield (24.18 q ha<sup>-1</sup>) of chickpea was recorded with the application of S @ 30 kg ha<sup>-1</sup> through Bentonite sulphur along with RDF (T<sub>6</sub>) and it was found to be on par with treatment S @ 30 kg ha<sup>-1</sup> through Gypsum + RDF (T<sub>9</sub>), S @ 20 kg ha<sup>-1</sup> through Bentonite sulphur + RDF (T<sub>5</sub>). The lowest seed yield of chickpea (14.88 q ha<sup>-1</sup>) was recorded in absolute control. Our results are in line with Srinivasulu *et al.* (2015), reported the effect of sulphur application in increasing the seed and straw yield of chickpea, while Das *et al.* (2016), reported increase in growth, plant yield and yield attributing characters of chickpea with increasing sulphur doses. The increasing in yield might be due to the fact that S perform many physiological function in cysteine, methionine and chlorophyll synthesis. Thus these bioactivity of sulphur might have played important role in improving yield attributing characters and yield of chickpea. Similar results were also given by Patel *et al.* (2014).

Data in (Table 2). The significantly higher straw yield (30.16 q ha<sup>-1</sup>) of chickpea was observed with the application of S @ 30 kg ha<sup>-1</sup> through Bentonite sulphur along with RDF (T<sub>6</sub>) and it was found to be on par with treatment S @ 30 kg ha<sup>-1</sup> through Gypsum + RDF (T<sub>9</sub>), S @ 20 kg ha<sup>-1</sup> through Bentonite sulphur + RDF (T<sub>5</sub>), S @ 20 kg ha<sup>-1</sup> through Gypsum + RDF (T<sub>8</sub>). The lowest straw yield of chickpea (18.25 q ha<sup>-1</sup>) was recorded in control treatment T<sub>1</sub>. Jadeja *et al.* (2016), also reported increased seed and straw yield of chickpea with sulphur application as compared to control.

The increase in seed and straw yield was might be due to increased sulphur availability and uptake as well as its active involvement in synthesis of amino acids, regulation of various metabolic and enzymatic processes along with enhanced nitrogen fixation and biomass accumulation which ultimately contributed to growth and yield. Similar finding of increased stover yield with sulphur application were also given by Srinivasulu *et al.* (2015).

## Nutrient Uptake:

**Nitrogen Uptake:** Data pertaining to Nitrogen uptake is given in (Table 3). There is significant improvement of nitrogen uptake with inoculation of sulphur treatment and presented in Table 3. The significantly highest total nitrogen uptake by chickpea ( $151.02 \text{ kg ha}^{-1}$ ) was observed with the application of S @  $30 \text{ kg ha}^{-1}$  through Bentonite sulphur + RDF ( $T_6$ ) and it was found to be at par with treatment S @  $30 \text{ kg ha}^{-1}$  through Gypsum + RDF ( $T_9$ ) and S @  $20 \text{ kg ha}^{-1}$  through Bentonite sulphur + RDF ( $T_5$ ). The application of sulphur @ 10, 20 and 30 kg sulphur per ha registered increasing trend in total nitrogen uptake ( $123.47$  to  $151.02 \text{ kg ha}^{-1}$ ) in case of Bentonite sulphur and ( $117.60$  to  $142.11 \text{ kg ha}^{-1}$ ) in case of Gypsum. The application of S @  $30 \text{ kg ha}^{-1}$  through Bentonite sulphur + RDF ( $T_6$ ) increased 44.39 per cent total nitrogen uptake as compared to S-free treatment.

The increase in nitrogen content in grain and straw might be due to the synergistic effect of both N and S which increased their availability in soil. The increased N uptake as results of S application might be due to an increment in protein synthesis and enhance photosynthesis (Yanwen Zhao *et al.*, 2013). Patel *et al.*, (2014) reported that application sulphur @  $40 \text{ kg ha}^{-1}$  increase the nitrogen content in seed and straw respectively. Srinivasulu *et al.* (2015) reported that increasing doses of sulphur significantly increase the uptake of nitrogen, phosphorus, potassium and sulphur. These findings are in accordance with Karprekar (2003), Singh *et al.* (2004), Sindagi (2014) and Kumar *et al.* (2006).

**Phosphorus Uptake:** Data pertaining to phosphorous uptake is given in (Table 4). Significantly higher total phosphorus uptake by chickpea ( $15.10 \text{ kg ha}^{-1}$ ) was observed with the application of S @  $30 \text{ kg ha}^{-1}$  through Bentonite sulphur + RDF ( $T_6$ ) and it was found to be at par with treatment S @  $30 \text{ kg ha}^{-1}$  through Gypsum + RDF ( $T_9$ ) and S @  $20 \text{ kg ha}^{-1}$  through Bentonite sulphur + RDF ( $T_5$ ). The phosphorus uptake by chickpea was recorded lowest in absolute control ( $5.76 \text{ kg ha}^{-1}$ ). The increase in phosphorous uptake with application S is might be due to Synergetic effect of sulphur application on phosphorous availability. These findings are in accordance with the results reported by Mir *et al.* (2013), Bahadur and Tiwari (2014) and Singh *et al.* (2016). Better root growth with proper nutrient supply may be the reason for extraction of available plant nutrient from soil depth and ultimately which improve the nutrient content in plant parts.

**Potassium Uptake:** Potassium uptake was also significantly improved with inoculation of sulphur treatment and presented in (Table 5). The significantly higher potassium uptake by chickpea ( $99.83 \text{ kg ha}^{-1}$ ) was observed with the application of S @  $30 \text{ kg ha}^{-1}$  through Bentonite sulphur + RDF ( $T_6$ ) and it was found at par with treatment S @  $30 \text{ kg ha}^{-1}$  through Gypsum + RDF ( $T_9$ ) and S @  $20 \text{ kg ha}^{-1}$  through Bentonite sulphur + RDF ( $T_5$ ). The lowest potassium uptake by chickpea ( $46.21 \text{ kg ha}^{-1}$ ) was recorded in control treatment  $T_1$ . Sulphur application might increase the availability of most of nutrient by reduction of pH of soil. This is may be reason of increased potassium uptake. The results content and uptake of potassium are in agreement with the findings reported by Singh *et al.* (2013), Das *et al.* (2016), Sindagi (2014), Mondal *et al.* (2005) and Solanki *et al.* (2017).

**Sulphur Uptake:** The data showed in (Table 6). indicated that the significantly highest sulphur uptake by chickpea seed ( $20.02 \text{ kg ha}^{-1}$ ) was recorded with the application of S @  $30 \text{ kg ha}^{-1}$  through Bentonite sulphur + RDF ( $T_6$ ) and it was found to be on par with treatment S @  $30 \text{ kg ha}^{-1}$  through

Gypsum + RDF (T<sub>9</sub>). The lowest sulphur total uptake by chickpea grain (8.35 kg ha<sup>-1</sup>) was recorded in absolute control. Increase in the sulphur content in seed and straw of chickpea might be due to application of increasing doses of sulphur. Similar results were also reported by Bahadur and Tiwari (2014); Singh *et al.* (2008); Patel *et al.* (2014); Kumar *et al.* (2006) and Chiaiese *et al.*, (2004) reported about the increment sulphur content in seed and stover of chickpea with the application of sulphur. Application of sulphur resulted in increased nitrogen fixation which might have promoted production of higher amount of above grounds dry matter that could have led to higher acquisition of nutrients ultimately resulted in higher nutrient content in seed and stover. Higher nutrient coupled with higher seed and stover yield lead to higher nutrient uptake. Similar data regarding increase in the uptake of sulphur in seed and straw of chickpea with increasing doses of sulphur was reported by Kala *et al.* (2017); Srinivasulu *et al.* (2015) and Islam and Ali (2009). When sulphur in bentonite-S comes into contact with soil moisture it breaks apart into fragments of fine dimension, which allows swift solubilization of S and gypsum has very low solubility. Hence availability and uptake sulphur is high with bentonite S as compared to sulphur with gypsum.

**Zinc Uptake:** Data pertaining to zinc uptake is given in (Table 7). Zinc Uptake was also significantly improved with application of sulphur treatment. The highest total zinc uptake (138.95 g ha<sup>-1</sup>) by chickpea was observed in treatment of application of S @ 30 kg ha<sup>-1</sup> through Bentonite sulphur + RDF (T<sub>6</sub>) and it was found to be at par with treatment S @ 30 kg ha<sup>-1</sup> through Gypsum + RDF (T<sub>9</sub>) and S @ 20 kg ha<sup>-1</sup> through Bentonite sulphur + RDF (T<sub>5</sub>). The lowest total zinc uptake by chickpea crop (76.45 g ha<sup>-1</sup>) was recorded in control treatment T<sub>1</sub>. Sulphur application increased plant Zn uptake as solubility of Zn increases with decrease in soil pH. The results are corroborated with the findings reported by Sindagi (2014); Yoo and James, (2003), and Cui Wang, (2005),

**Iron Uptake:** The data regarding Iron uptake is presented in (Table 8) indicated that it is significantly improved with sulphur application. The highest total iron uptake by chickpea (313.88 g ha<sup>-1</sup>) was observed with the application of S @ 30 kg ha<sup>-1</sup> through Bentonite sulphur + RDF (T<sub>6</sub>) which was followed by treatment S @ 30 kg ha<sup>-1</sup> through Gypsum + RDF (T<sub>9</sub>) and S @ 20 kg ha<sup>-1</sup> through Bentonite sulphur + RDF (T<sub>5</sub>). The lowest total iron uptake by chickpea crop (181.83 g ha<sup>-1</sup>) was recorded in treatment absolute control. Sulphur application resulted in significant increased in Fe uptake is mainly due to acidification effect produced as result of sulphur application. The results are in accordance with the findings reported by Sindagi (2014). Sulphur application resulted in an increased in Fe uptake as a recorded by Malewar and Ismail (1997).

**Manganese Uptake:** The data presented in (Table 9) indicated that there was significant improvement in total uptake of manganese (140.24 g ha<sup>-1</sup>) by chickpea with application of treatment S @ 30 kg ha<sup>-1</sup> through Bentonite sulphur + RDF (T<sub>6</sub>) which was followed by treatment S @ 30 kg ha<sup>-1</sup> through Gypsum + RDF (T<sub>9</sub>) and S @ 20 kg ha<sup>-1</sup> through Bentonite sulphur + RDF (T<sub>5</sub>). The lowest manganese uptake by chickpea (78.42 g ha<sup>-1</sup>) was recorded in control treatment T<sub>1</sub>. There was significant increase in Mn uptake due to S application which coincide with finding of Rahman *et al.* (2011) who observed that an increase in Mn uptake by corn plant with the application elemental S as result of soil acidification although temporary. The similar findings was reported by Sindagi (2014).

**Copper Uptake:** Data pertaining to Copper uptake is given in (Table 10). Significantly highest uptake of copper ( $45.48 \text{ g ha}^{-1}$ ) by chickpea was observed in treatment application of S @  $30 \text{ kg ha}^{-1}$  through Bentonite sulphur + RDF ( $T_6$ ) and it was found to be at par with treatment S @  $30 \text{ kg ha}^{-1}$  through Gypsum + RDF ( $T_9$ ) and S @  $20 \text{ kg ha}^{-1}$  through Bentonite sulphur + RDF ( $T_5$ ). Sulphur application resulted in a significant increase in copper uptake, which is in line with previous finding and mainly due to acidification effect produced as a result of S application. Ghosh *et al.*, (2000) and Rahman *et al.*, (2011). The results are in accordance with the findings reported by Sindagiet *al.* (2014).

**Quality of chickpea:** The test weight (Table 11) of chickpea seed was found to vary from 20.46 to 25.11 g. Test weight (100 seed) was found to increase with the application of increasing doses of sulphur. The significantly highest test weight in chickpea seed (25.11 g) was recorded in treatment S @  $30 \text{ kg ha}^{-1}$  through Bentonite sulphur + RDF ( $T_6$ ) and it was found to be at par with treatment S @  $30 \text{ kg ha}^{-1}$  through Gypsum + RDF ( $T_9$ ). The lowest test weight in chickpea seed (20.46) was recorded in control treatment  $T_1$ . The application of increasing dose of sulphur from 10 to  $30 \text{ kg S per ha}$  on sulphur deficient soils increased the test weight of chickpea linearly from 22.35 to 25.11 g with Bentonite sulphur and from 22.12 to 24.58 g with Gypsum, respectively.

Data pertaining to Protein Content (%) is given in (Table 12). The protein content of chickpea seed was found to vary from 18.98 to 21.13 % in all treatment. Data pertaining to protein yield ( $\text{kg ha}^{-1}$ ) is given in (Table 13). The protein yield of chickpea seed was found to vary from 282.0 to  $510.8 \text{ kg ha}^{-1}$  due to application of sulphur. Data revealed that the protein content (%) and protein yield ( $\text{kg ha}^{-1}$ ) in (Table 12 and 13) were affected due to application of sulphur application might be due to increased sulphur and nitrogen availability which help in synthesizing some sulphur containing amino acids like Homocysteine, Cysteine and methionine, thus resulting in increased synthesis of protein. Similar finding were also reported by Kaisheret *al.* (2010); Patel *et al.* (2010) and Singh (2008) reported that combined application of S ( $20 \text{ kg ha}^{-1}$  recorded significantly higher protein content over control. Ram and Katiyar (2013) revealed that increase in S levels from 0 to  $40 \text{ kg ha}^{-1}$  significantly increased the protein content (23.92 and 24.07%).

**Table 1: Seed yield of chickpea as influenced by sulphur application**

Treatments	Seedyield( $\text{q ha}^{-1}$ )				% Response over	% increase over S
	2020-21	2021-22	2022-23	Pooled Mean		

						control	free RDF
1	Absolute control	16.52	13.80	14.33	14.88		
2	S free RDF (NPK through Urea, DAP, MOP)	17.84	20.40	18.91	19.05	28.0	
3	RDF (NPK through Urea, SSP, MOP)	18.75	23.96	21.05	21.26	42.8	11.58
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	19.93	23.56	20.67	21.39	43.7	12.29
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	22.80	24.86	22.53	23.40	57.2	22.82
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	23.26	25.58	23.70	24.18	62.5	26.94
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	18.70	23.35	20.25	20.76	39.5	9.01
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	19.27	24.45	22.00	21.91	47.2	15.01
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	22.00	25.35	23.10	23.48	57.8	23.28
	SE (m) ±	1.11	0.66	1.00	0.72		
	CD at 5%	3.34	1.99	3.02	2.15		
	CV	9.70	5.04	8.43	5.86		

**Table 2: Straw yield of chickpea as influenced by sulphur application**

Treatments	Straw yield (q ha <sup>-1</sup> )			
	2020-21	2021-22	2022-23	Pooled Mean
1 Absolute control	20.26	17.25	17.23	18.25
2 S free RDF (NPK through Urea, DAP, MOP)	21.97	25.49	23.43	23.63
3 RDF (NPK through Urea, SSP, MOP)	22.51	29.95	26.14	26.20
4 T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	24.20	29.38	26.11	26.56
5 T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	26.72	31.09	28.51	28.77
6 T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	27.81	31.98	30.70	30.16
7 T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	22.87	29.19	25.29	25.79
8 T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	23.38	30.58	27.51	27.16
9 T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	25.70	31.72	28.87	28.76
SE (m) ±	1.11	0.86	1.21	0.94
CD at 5%	3.33	2.57	3.64	2.82

**Table 3: Total uptake of N (kg ha<sup>-1</sup>) of chickpea as influenced by sulphur application**

Treatments		Total uptake of N (kg ha <sup>-1</sup> )			
		2020-21	2021-22	2022-23	Pooled Mean
1	Absolute control	86.49	76.11	74.51	79.03
2	S free RDF (NPK through Urea, DAP, MOP)	96.39	115.30	102.07	104.59
3	RDF (NPK through Urea, SSP, MOP)	107.19	143.25	118.67	123.04
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	112.44	139.35	118.62	123.47
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	134.18	153.19	134.68	140.68
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	141.38	162.83	148.86	151.02
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	103.30	135.58	113.92	117.60
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	113.03	149.05	128.60	130.22
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	129.66	157.29	139.38	142.11
	SE (m) ±	6.35	5.38	5.46	4.16
	CD at 5%	19.03	16.13	16.36	12.47

**Table 4: Total uptake of P (kg ha<sup>-1</sup>) of chickpea as influenced by sulphur application**

Treatments		Total uptake of P (kg ha <sup>-1</sup> )			
		2020-21	2021-22	2022-23	Pooled Mean
1	Absolute control	6.32	6.08	4.87	5.76
2	S free RDF (NPK through Urea, DAP, MOP)	7.85	10.31	9.70	9.29
3	RDF (NPK through Urea, SSP, MOP)	9.16	13.23	12.55	11.65
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	9.45	12.51	11.86	11.27
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	11.95	14.50	14.32	13.59
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	13.38	15.50	16.41	15.10
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	8.86	11.92	10.14	10.31
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	9.38	13.73	13.28	12.13
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	12.31	14.87	15.12	14.10
	SE (m) ±	0.58	0.72	1.01	0.52
	CD at 5%	1.75	2.17	3.04	1.55

**Table 5: Total uptake of K (kg ha<sup>-1</sup>) of chickpea as influenced by sulphur application**

Treatments		Total uptake of K (kg ha <sup>-1</sup> )			
		2020-21	2021-22	2022-23	Pooled Mean
1	Absolute control	49.23	45.54	43.88	46.21
2	S free RDF (NPK through Urea, DAP, MOP)	55.40	69.48	64.28	63.05
3	RDF (NPK through Urea, SSP, MOP)	66.00	95.39	85.59	82.33
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	66.97	88.88	79.95	78.60
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	80.52	99.78	94.77	91.69
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	87.48	107.00	104.99	99.83
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	62.03	85.76	75.51	74.43
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	66.26	94.32	87.38	82.65
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	79.47	103.05	98.16	93.56
	SE (m) ±	3.40	2.63	4.34	3.10
	CD at 5%	10.18	7.88	13.01	9.26

**Table 6: Total uptake of S (kg ha<sup>-1</sup>) of chickpea as influenced by sulphur application**

Treatments		Total uptake of S (kg ha <sup>-1</sup> )			
		2020-21	2021-22	2022-23	Pooled Mean
1	Absolute control	8.65	7.83	8.57	8.35
2	S free RDF (NPK through Urea, DAP, MOP)	10.05	12.54	12.20	11.60
3	RDF (NPK through Urea, SSP, MOP)	11.79	17.08	15.29	14.72
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	11.61	15.85	15.21	14.22
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	15.65	18.73	18.37	17.59
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	17.29	20.91	21.85	20.02
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	10.70	15.29	14.04	13.34
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	12.55	17.92	17.33	15.93
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	14.99	20.11	19.39	18.17
	SE (m) ±	0.86	1.00	0.99	0.62
	CD at 5%	2.59	3.00	2.99	1.85

**Table 7: Total uptake of Zn (g ha<sup>-1</sup>) of chickpea as influenced by sulphur application**

Treatments		Total uptake of Zn (g ha <sup>-1</sup> )			
		2020-21	2021-22	2022-23	Pooled Mean
1	Absolute control	84.21	71.74	73.40	76.45
2	S free RDF (NPK through Urea, DAP, MOP)	92.86	107.84	99.68	100.12
3	RDF (NPK through Urea, SSP, MOP)	98.24	129.70	113.68	113.87
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	106.90	130.02	116.01	117.64
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	122.78	140.26	128.97	130.67
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	128.97	147.63	140.26	138.95
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	99.77	128.88	112.22	113.62
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	104.60	137.88	124.83	122.44
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	120.18	146.16	134.45	133.60
	SE (m) ±	5.49	3.83	6.31	4.08
	CD at 5%	16.47	11.47	18.90	12.22

**Table 8: Total uptake of Fe (g ha<sup>-1</sup>) of chickpea as influenced by sulphur application**

Treatments		Total uptake of Fe (g ha <sup>-1</sup> )			
		2020-21	2021-22	2022-23	Pooled Mean
1	Absolute control	200.83	172.52	172.15	181.83
2	S free RDF (NPK through Urea, DAP, MOP)	218.84	257.19	236.45	237.49
3	RDF (NPK through Urea, SSP, MOP)	229.49	305.08	265.96	266.84
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	246.74	301.92	265.86	271.51
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	280.25	321.87	293.27	298.46
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	292.56	334.05	315.02	313.88
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	232.05	299.70	258.45	263.40
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	240.23	316.36	283.70	280.10
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	272.83	330.97	300.59	301.46
	SE (m) ±	13.23	10.15	14.24	9.40
	CD at 5%	39.66	30.41	42.68	28.18

**Table 9: Total uptake of Mn (g ha<sup>-1</sup>) of chickpea as influenced by sulphur application**

Treatments		Total uptake of Mn (g ha <sup>-1</sup> )			
		2020-21	2021-22	2022-23	Pooled Mean
1	Absolute control	85.68	74.45	75.13	78.42
2	S free RDF (NPK through Urea, DAP, MOP)	93.68	111.19	101.06	101.98
3	RDF (NPK through Urea, SSP, MOP)	98.67	132.68	114.74	115.36
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	107.48	133.10	116.97	119.18
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	123.18	143.78	130.03	132.33
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	129.41	150.20	141.12	140.24
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	100.81	131.93	113.44	115.40
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	105.07	140.19	125.58	123.61
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	120.43	148.12	134.25	134.27
	SE (m) ±	5.14	5.03	6.39	4.13
	CD at 5%	15.40	15.09	19.15	12.39

**Table 10: Total uptake of Cu (g ha<sup>-1</sup>) of chickpea as influenced by sulphur application**

Treatments		Total uptake of Cu (g ha <sup>-1</sup> )			
		2020-21	2021-22	2022-23	Pooled Mean
1	Absolute control	21.48	18.48	19.00	19.65
2	S free RDF (NPK through Urea, DAP, MOP)	24.75	29.87	27.33	27.31
3	RDF (NPK through Urea, SSP, MOP)	27.62	37.49	33.09	32.73
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	31.56	39.93	34.79	35.43
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	38.71	45.31	41.29	41.77
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	41.93	49.01	45.49	45.48
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	29.44	39.27	33.51	34.07
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	32.44	44.14	39.01	38.53
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	38.70	48.02	43.28	43.33
	SE (m) ±	2.33	1.58	1.94	1.26
	CD at 5%	6.97	4.75	5.80	3.79

**Table 11: Test weight (g/100 seeds) of chickpea as influenced by sulphur application**

Treatments		Test weight (g/100 seeds)			
		2020-21	2021-22	2022-23	Pooled Mean
1	Absolute control	19.95	20.23	21.19	20.46
2	S free RDF (NPK through Urea, DAP, MOP)	20.47	22.92	21.87	21.75
3	RDF (NPK through Urea, SSP, MOP)	21.69	24.18	23.64	23.17
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	21.20	23.69	22.15	22.35
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	21.81	24.47	23.09	23.12
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	24.55	25.67	25.10	25.11
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	20.89	23.47	22.00	22.12
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	21.33	24.08	22.81	22.74
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	24.05	25.27	24.43	24.58
	SE (m) ±	0.54	0.59	0.57	0.30
	CD at 5%	1.64	1.76	1.70	0.90

**Table 12: Protein content (%) of chickpea as influenced by sulphur application**

Treatments		Protein content (%)			
		2020-21	2021-22	2022-23	Pooled Mean
1	Absolute control	18.58	19.46	18.91	18.98
2	S free RDF (NPK through Urea, DAP, MOP)	19.20	19.77	19.63	19.53
3	RDF (NPK through Urea, SSP, MOP)	19.59	20.71	19.92	20.07
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	20.03	19.94	20.17	20.04
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	20.41	20.76	20.80	20.66
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	20.95	21.09	21.34	21.13
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	19.92	19.90	20.02	19.95
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	20.22	20.55	20.68	20.48
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	20.68	20.87	20.93	20.83
	SE (m) ±	0.45	0.39	0.34	0.13
	CD at 5%	1.34	1.18	1.03	0.39

**Table 13: Protein yield (kg ha<sup>-1</sup>) of chickpea as influenced by sulphur application**

Treatments		Protein yield (kg ha <sup>-1</sup> )			
		2020-21	2021-22	2022-23	Pooled Mean
1	Absolute control	306.82	268.90	270.23	282.0
2	S free RDF (NPK through Urea, DAP, MOP)	341.82	403.20	370.42	371.8
3	RDF (NPK through Urea, SSP, MOP)	367.15	496.10	420.19	427.8
4	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Bentonite Sulphur	398.78	470.00	416.30	428.4
5	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Bentonite Sulphur	466.37	515.80	467.67	483.3
6	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Bentonite Sulphur	487.07	539.50	505.78	510.8
7	T <sub>2</sub> + S @ 10 kg ha <sup>-1</sup> through Gypsum	371.76	464.20	405.50	413.8
8	T <sub>2</sub> + S @ 20 kg ha <sup>-1</sup> through Gypsum	390.22	502.60	454.68	449.2
9	T <sub>2</sub> + S @ 30 kg ha <sup>-1</sup> through Gypsum	455.02	528.70	483.60	489.1
	SE (m) ±	21.15	16.27	20.33	14.11
	CD at 5%	13.39	48.78	60.94	42.31

#### 4. CONCLUSION

On the basis of the above findings, it may be concluded that among the various treatments combinations, the application of S @ 30 kg ha<sup>-1</sup> through Bentonite sulphur along with RDF exhibited better performance in chickpea crop thereby influencing the yield, uptake and quality were significantly at T<sub>6</sub>. The significantly highest grain yield (24.18 q ha<sup>-1</sup>) and straw yield (30.16 q ha<sup>-1</sup>) were recorded with the application of S @ 30 kg ha<sup>-1</sup> though Bentonite sulphur along with RDF and the lowest seed yield (14.88 q ha<sup>-1</sup>) and straw yield (18.25 q ha<sup>-1</sup>) was recorded in treatment absolute control. Similarly, highest N, P, K, S and micronutrient uptake and quality parameters like test weight, protein (%) and protein yield (kg ha<sup>-1</sup>) were observed with application of @ 30 kg ha<sup>-1</sup> through Bentonite sulphur along with RDF followed by the treatment of application of S @ 30 kg ha<sup>-1</sup> through Gypsum + RDF.

#### REFERENCES

- Anonymous, (2017) b. Agricultural Statistics at a Glance, (2016). Directorate of Economics and Statistics, Ministry of Agriculture Govt. of India (Website <http://www.dacnet.nic.in/ean>).
- AOAC. (1960). Official Methods of analysis (18<sup>th</sup> ed.) *Association of official agricultural chemists*, Washington.
- Bahadur, Lal and Tiwari, D.D. (2014). Nutrient management in mungbean (*Vigna radiata* L.) through sulphur and biofertilizers. *Legume Research*.37 (2): 180-187.
- Chapman, H.D. and Pratt, P.F. (1962). Methods of analysis for soils, plants and water, *University of California Agriculture Division*. Page number.
- Chesnin, L., & Yien, C. H. (1951). Turbidimetric determination of available sulfates. *Soil Science Society of America Journal*, 15(C), 149-151.

- Chiaiese, P., Ohkama-Ohtsu, N., Molvig, L., Godfree, R., Dove, H., Hocart, C., Fujiwara, T., Higgins, T. J. V. and Tabe, L. (2004). Sulphur and nitrogen nutrition influence the response of chickpea seeds to an added, transgenic sink for organic sulphur. *Journal of Experimental Botany*.55(404): 1889-1901.
- Cui Y. and Wang Q. (2005): Interactive effect of Zn and S on their uptake by spring wheat. *Journal of Plant Nutrition*, 28: 639–649.
- Das, S.K., B. Biswas. and K. Jana, (2016). Effect of farm yard manure, phosphorus and sulphur on yield parameters, yield, nodulation, nutrient uptake and quality of chickpea (*Cicer arietinum* L.). *Journal of Applied and Natural Sciences*.8 (2): 545-549.
- Das, S.K., Chopra, P., Chatterjee, S.R., Abrol, Y.P. and -Deb, D.L. (1975) Influence of sulphur fertilization on yield of maize and protein quality of cereals. *Fertilizers News*. 20(3): 30 - 32.
- Ghosh P.K., Hati K.M., Mandal K.G., Misra A.K., Chaudhry R.S., Bandyopadhyay K.K. (2000). Sulphur nutrition in oilseeds and oilseedbased cropping systems. *Fertilizer News*, 45: 27–40.
- Hanway, J. and Heidal, H. S. (1952). Soil testing laboratory procedures. *Jowa Agriculture*,57:1-31.
- Isaac, R. A. and Kerber, J. D. (1971). Atomic absorption and flame photometry: Techniques and uses in soil, plant, and water analysis. *Instrumental methods for analysis of soils and plant tissue*. 17-37.
- Islam, M. and S. Ali. (2009). Effect of integrated application of sulphur and phosphorus on nitrogen fixation and nutrient uptake by chickpea (*Cicer arietinum* L.). *Agrociencia Journal*. 43 (8): 815-826
- Jackson, M. T. and Faller, A. (1973). Structural analysis and dynamics of the plant communities of Wizard Island, Crater Lake National Park. *Ecological Monographs*, 43(4), 441-461.
- Jackson, W. A., Flesher, D., & Hageman, R. H. (1973). Nitrate uptake by dark-grown corn seedlings: some characteristics of apparent induction. *Plant Physiology*, 51(1), 120-127.
- Jadeja, A.S., A.V. Rajani, Foram, Chapdiya, S.C. Kaneriya, and N.R. Kavar, (2016). Soil application of potassium and sulphur and effect on growth and yield components of chickpea (*Cicer arietinum* L.) under south Saurashtra region of Gujarat. *International Journal of Science Environment and Technology*.5 (5): 3172-3176.
- Kaisher, M. S., Rahman, M. A., Amin, M. H. A., Amanullah, A. S. M. and Ahsanullah, A. S. M. (2010). Effects of sulphur and boron on the seed yield and protein content of mung bean. *Bangladesh Research Publications Journal*. 3(4), 1181-1186.
- Kala, D.C., R.N. Dixit, S.S. Meena, G. Nanda and R. Kumar. (2017). Effect of graded doses of sulphur and boron on yield attributes and nutrient uptake by chickpea. *International Journal of Current Microbiology and Applied Science*. 6 (6), 55-60.
- Kaprekar, N., D.S. Sasode, and A. Patil. (2003). Yield, nutrient uptake and economics of gram (*Cicer arietinum* L.) as influenced by phosphorus and sulphur levels and PSB inoculation under irrigated conditions. *Legume Research*.26: 125-127.

- Katkar, R.N., V.K. Kharche, S.R. Lakhe, P.R. Deshmukh, A.K. Shukla, Pankaj Tiwari, A.B. Aage and A.D. Kadlag. (2017). Geographical Information based Micro and Secondary nutrients in soils of Maharashtra. Dr. P.D.K.V., Akola. Bulletin No. 491: 10.
- Kumar, Ashok, Prasad, Shambhu. and S.B. Kumar. (2006). Effect of boron and sulphur on performance of gram (*Cicer arietinum* L.). *Indian Journal Agronomy*.51 (1): 57-59.
- Malewar G.U., Ismail S. (1997): Sulphur in balanced fertilization in western India. In: Proceedings of the TSI/FAI/IFA Symposium on Sulphur in Balanced Fertilization, New Delhi, 14.
- Mir, A.H., S.B. Lal, M. Salmani, M. Abid and I. Khan. (2013). Growth, yield and nutrient content of black gram (*Vigna mungo*) as influenced by levels of phosphorus, sulphur and phosphorus solubilizing bacteria. *SAARC Journal of Agriculture*.11 (1): 1-6.
- Mondal, S.S., P. Mandal, M. Saha, A. Bag, S. Nayak and G. Sounda. (2005). Effect of potassium and sulphur on the productivity, nutrient uptake and quality improvement of chickpea. *Journal of Crop and Weed*. 2 (1): 64-66.
- Panse, V. G. And P. V. Sukhantme. (1985). Statistical method of Agricultural. *Indian council of Agricultural research*, New Delhi.
- Patel, H. K., Patel, P. M., Suthar, J. V. and Patel, M. R. (2014). Yield, Quality and Post Harvest Nutrient Status of Chickpea as Influence by Application of Sulphur and Phosphorus Fertilizer Management. *International Journal of Scientific and Research Publications*, 4 (7), 1-4.
- Rahman M.M., Soaud A.A., Al Darwish F.H. and Sofian-Azirun M. (2011). Responses of sulfur, nitrogen and irrigation water on Zea mays growth and nutrients uptake. *Australian Journal of Crop Science*.5: 350–360.
- Ram, S. and Katiyar, T. P. S. (2013). Effect of sulphur and zinc on the seed yield and protein content of summer mung bean under arid climate. *International Journal of Natural Sciences*. 4(3), 563-566.
- Scherer, H.W. (2009). Sulfur in soils. *Journal Plant Nutrition Soil Science*.,172: 326–335
- Shukla, A.K., P. Tiwari, S. Siddiqui, A.K. Patra and S.K. Chaudhary. (2016). Micro and Secondary nutrients in Indian soils, condition of deficiency, prevention and recommendations. *Indian institute of soil science*. Bhopal.3: 25.
- Sindagi, A.S. (2014). Studies on levels of phosphorus and sulphur on yield, quality and uptake of nutrients by chickpea in a Vertisol under irrigation. M.Sc Thesis. *University of Agricultural Sciences*, Dharwad.
- Singh A.K., M.K. Meena, R.C. Bharati and R.M. Gade. (2013). Effect of sulphur and zinc management on yield, nutrient uptake, changes in soil fertility and economics in rice (*Oryza sativa*)-lentil (*Lens culinaris*) cropping system. *Indian Journal of Agricultural Sciences*.83 (3): 344–348.
- Singh, A. K., Meena, M. K. and Bharati, R. C. (2008). Sulphur and Zinc Nutrient Management in rice-lentil cropping system. *International Conference on "Life Science Research for Rural and Agricultural Development"* (pp 66-67). 27-29 December, 2011, CPRS Patna (Bihar).
- Singh, S., Bawa, S. S., Singh, S., Sharma, S. C. and Sheoran, P. (2008). Productivity, profitability and sustainability of rain-fed chickpea under inorganic and biofertilization in foothills of north-west Himalayas. *Archives of Agronomy and Soil Science*, 61(8), 1151-1163.

- Singh, S., S.S. Saini and B.P. Singh. (2004). Effect of irrigation, sulphur and seed inoculation on growth, yield and sulphur uptake of chickpea (*Cicer arietinum*) under late sown conditions. *Indian Journal Agronomy*.49 (1): 57-59.
- Solanki, S., Pamu and M.S. Solanki. (2017). Effect of sulphur and zinc on content and uptake of nutrients of summer green gram (*Vigna radiata* L. Wilezeck) under medium black calcareous soils. *International Journal of Agricultural Sciences and Research*.7: 657-662.
- Srinivasulu, D. V., Solanki, R. M., Kumar, N. N., Bhanuprakash, M. and Vema raju, A. (2016). Effect of irrigation based on IW/CPE ratio and sulphur levels on yield and quality of gram (*Cicer arietinum* L.). *Legume Research*. 39 (4), 601-604.
- Srinivasulu, D.V., R.M. Solanki, C.R. Kumari, and M.V. Babu. (2015). Nutrient uptake, yield and protein content of chickpea (*Cicer arietinum* L.) as influenced by irrigation and sulphur levels in medium black soils. *International Journal of Agricultural Sciences*.11: 54-58.
- Tiwari, B. K and Tripathi, P. N. (2014). Yield gap analysis of chickpea (*Cicer arietinum*) through front line demonstration on farmer's fields. *The Journal of Rural and Agricultural Research*, 14(1), 5-8.
- Yan wen., Zhao, Y. and Wang, J. (2013). Suitable water and nitrogen treatment improves soil microbial biomass carbon and nitrogen and enzyme activities of paddy field. *Transactions of the Chinese Society of Agricultural Engineering*, 29 (21), 91-98.
- Yoo M.S. and James B.R. (2003): Zinc exchangeability as a function of pH in citric acid-amended soils. *Soil Science*, 168: 356–367.