

Original Research Article

Affirmation of Experimental Results on Groundnut-Based Millet Intercropping System at Farmers Field under Rainfed Condition

ABSTRACT

Aims: The research aims to tackle the issue of low groundnut productivity in rainfed areas and low-fertility marginal lands, which is below the national average, leading to economic constraints for farmers. Millets, known for their resilience, are considered suitable for cultivation in challenging environments. Intercropping, growing multiple crops simultaneously, is recognized as an essential system to mitigate erratic climatic conditions and act as insurance against uncertainties. The experiment evaluates the productivity and profitability of an intercropping system combining groundnut and millets in a rainfed ecosystem. The goal is to provide a sustainable and economically viable option for farmers in such areas by leveraging the robustness of millets and improving agricultural outcomes and resilience to climate-related risks.

Study design: Complete randomized block design (RCBD)

Place and duration of study: A field experiment was carried out during *Kharif* 2017-18, 2018-19 and 2019-20 at Zonal Agricultural and Horticultural Research Station, Babbur farm, Hiriyyur.

Methodology: An experiment was laid out with complete randomized block design (RCBD) with ten treatments and was replicated thrice. Treatments comprises of testing of sole groundnut, finger millet and foxtail millet against intercropping of groundnut with finger millet, little millet and foxtail millet at 5:2 and 6:1 row proportion.

Results: The results of pooled data indicated that significantly higher groundnut pod equivalent yield (2123 kg ha^{-1}) was recorded with Groundnut + finger millet (5:2) as compared to other treatments. However, significantly higher net returns (Rs. 58532 ha^{-1}) and B:C ratio (2.55) were recorded with Groundnut + foxtail millet (6:1) as compared to other treatments.

Conclusion: The study confirms that cultivating crops as sole crops under rainfed conditions is risky due to the adverse effects of low and erratic rainfall, leading to reduced productivity. To achieve higher productivity in such challenging conditions, it is essential to adopt intensification and diversification of crops. Intercropping of groundnut + foxtail millet (6:1) was found profitable and efficient intercropping system.

Key words: Intercropping systems, groundnut + foxtail millet, groundnut + millets, groundnut pod equivalent yield.

1. INTRODUCTION

Groundnut is referred to as the "king of oilseeds" crops and is one of the most significant oilseed crops grown in tropical and subtropical areas of the world. Groundnut is a member of the Leguminosae family. Another name for groundnut is "poor man's almond" and it is one of food and economic crop of the country. India has 6.09 million hectares of groundnut cultivation, producing 10.21 million tonnes at productivity of 1676 kg ha⁻¹. The greatest producer in India, Gujarat produces 40.42 per cent of all groundnuts, followed by Rajasthan (18.91 %), Tamil Nadu (9.25 %), Andhra Pradesh (7.62 %), and Karnataka (6.62 %) [1]. In Karnataka, it is grown over an area of 0.70 million hectare with a production of 0.68 million tonnes having a productivity of 966 kg ha⁻¹. The productivity of groundnut is very low than national average productivity because primarily it is grown under rain fed situation and low fertile marginal lands which are subjected to the vagaries of monsoon. Low yield and in the worst circumstances, complete crop failure are the results of insufficient and unpredictable rainfall combined with insect pest damage. Millets are tough and robust plants that may be produced in a variety of unfavourable agro-climatic conditions. They play a significant part in the food security of communities that depend on livestock for their way of life. Since intercropping is one of the important cropping systems that is recommended to mitigate the aberrant climatic conditions [2], it was undertaken to study the effect of intercropping system on growth and yield of groundnut under rain fed condition. Intercropping aims to increase total productivity per unit area through equitable and judicious use of land resource and farming inputs, including labour.

2. MATERIAL AND METHODS

2.1 Experimental site and soil

The field experiment was conducted at Zonal Agricultural and Horticultural Research station, Babbur farm, Hiriyyur during *kharif* 2017-18, 2018-19 and 2019-20 under rainfed situation which is

comes under central dry zone of Karnataka. The experimental site is situated at 13° 57' 32" North latitude and 70° 37' 38" East longitude and at an altitude of 606 meters above MSL. The soil of the experimental site is belonged to order vertisol with slightly alkaline pH (8.10), low in organic carbon (1.90 g kg⁻¹), available nitrogen (258 kg ha⁻¹), medium in available phosphorus (35 kg ha⁻¹) and potassium (315 kg ha⁻¹).

2.2 Design of experiment and treatment details

The experiment was laid out in complete randomized block design (RCBD) concept consist of ten treatments with three replications. The treatment comprises of T₁: Sole groundnut, T₂: Sole finger millet, T₃: Sole little millet, T₄: Sole foxtail millet, T₅: Groundnut + finger millet (5:2), T₆: Groundnut + little millet (5:2), T₇: Groundnut + foxtail millet (5:2), T₈: Groundnut + finger millet (6:1), T₉: Groundnut + little millet (6:1) and T₁₀: Groundnut + foxtail millet (6:1). The cultivar used are for groundnut - G-2-52, finger millet - ML-365, little millet - Sukshema and foxtail millet - HMT 100-1. Best performed intercropping system *i.e.*, Groundnut + foxtail millet (6:1) was taken for farm trial at different locations *viz.*, Hiiyur, Kathalagere and Chitradurga during *kharif* 2020 and 2021.

2.3 Data collection for analysis

The crops were harvested separately from the net plot at physiological maturity and were threshed manually and pod and grain yield were weighed from the net plot and converted into kg ha⁻¹. Intercrop yields were computed as groundnut pod equivalent yields (GPEY). GPEY is a simple expression in an intercropping to compare the economics of intercrops by converting grain/seed/economic part *etc.* in terms of gross returns/net returns for valid comparison. The economics was worked out from prevailing market prices of inputs and outputs for different treatments.

2.4 Statistical analysis

The data recorded during the investigation were compiled and analysed for statistical significance as per the analysis of variance for the complete randomized block design (RCBD). Fisher's method of analysis of variance (ANOVA) as described by Gomez and Gomez (1984) [3] was adopted for the purpose. Standard error of mean and coefficient of variability have been worked out for a set of observations under each character at $P=0.05$ to interpret the significance.

3. RESULTS AND DISCUSSION

3.1 Productivity of groundnut + millets intercropping systems

Growing of crops as a sole crop found to be risky under rainfed conditions due to low and erratic rainfall, which ultimately results in low productivity. Under such conditions in order to achieve higher productivity intensification and diversification of crops is essential. In the present investigation, the result revealed that higher groundnut pod yield (2383 kg/ha) and gross returns (Rs. 111838/ha) were obtained in sole groundnut, this could be due to optimum plant population in sole groundnut and compared to intercropping groundnut (Table 1). Unlike observed in sole millets. Among the intercropping systems, groundnut + finger millet (5:2) recorded significantly higher groundnut pod equivalent yield (GPEY) of 2124 kg/ha than other treatments. However, it was statistically on par with Groundnut + little millet (5:2), Groundnut + foxtail millet (5:2), Groundnut + finger millet (6:1), Groundnut + little millet (6:1) and Groundnut + foxtail millet (6:1). It was mainly due to more space available between two millet rows thus there is a better availability of light lead to higher yield of groundnut in the intercropping system and thereby envisages effective utilization of the resources along with millets. Similar findings have been reported by Shwethanjali *et al.* (2018) [4]. Yield of any crop depended on its yield parameters. Significantly higher number of pods and pod weight plant-1 were recorded under groundnut intercropped with foxtail millet at ratio of 6:1 followed by little millet and finger millet. This mainly due to variation in translocation of photosynthates from source to sink as there is a greater availability of light due to differential growth habit and its efficient use, less competition for resources by component crops and efficient utilization of available resources. Similar findings have been reported by Maitra *et al.* (2000) [5] and Bassi and Dugje (2015) [6].

When comparing returns, it is found that higher net returns (Rs. 58532/ha) and B:C ratio (2.55) was recorded with Groundnut + foxtail millet (6:1) as compared to other treatments (Table 1). It was mainly due to higher groundnut pod equivalent yield and lower cost of cultivation under intercropping systems of Groundnut + foxtail millet (6:1) than other intercropping systems. Intercropping of groundnut + foxtail millet (6:1) recorded higher B:C ratio (2.55) to the tune of 23 % as compared to sole groundnut (1.96). This was mainly due to low COC especially seed price of groundnut and higher yield of millets. LER all the intercropping treatments in the present study recorded more than one value indicating the yield advantage in all intercropping systems. This might be due to higher yield of groundnut in the intercropping system and thereby envisages effective utilization of the resources along with millets. Higher yield levels under intercropping systems were mainly due to variation in translocation of photosynthates from source to sink as there is a greater

availability of light due to differential growth habit and its efficient use, less competition for resources by component crops and efficient utilization of available resources.

3.2 Results of farm trial

The study conducted on-farm trials for two consecutive years (2020 and 2021) to investigate the impact of different cropping systems under rainfed conditions. It revealed that growing crops as a sole crop in such conditions poses risks due to low and erratic rainfall, resulting in low productivity. To achieve higher productivity, the study suggests intensification and diversification of crops. The study specifically focused on evaluating the performance of a groundnut-based intercropping system with foxtail millet crops at different locations. The intercropping system involved planting groundnut and foxtail millet together, which proved to be beneficial.

The results indicated that the intercropping system produced approximately 10.35 % higher groundnut pod equivalent yield, reaching 2066 kg/ha, compared to the sole groundnut yield of 1852 kg/ha (Table 2). Additionally, the net returns from the intercropping system were significantly increased, reaching Rs. 69159/ha, compared to the normal sole groundnut which generated net returns of Rs. 49795/ha. This represents a considerable increase of 29% in net returns when adopting the intercropping method compared to growing groundnut alone.

Furthermore, the study showed that the intercropping treatment with groundnut and foxtail millet (6:1) recorded a higher benefit-cost ratio (B:C ratio) of 2.72, which was 25 % higher compared to the B:C ratio of 2.05 observed in the case of sole groundnut. The B:C ratio is a key indicator of economic profitability in agriculture, and the higher value achieved through intercropping demonstrates its economic viability and potential benefits.

One of the major advantages of the intercropping system in rainfed conditions is its ability to act as a natural insurance against total crop failure. The intercropped crops provide mutual support and resilience, reducing the risk of complete failure in case of adverse weather events or other challenges. This aspect enhances production sustainability and helps farmers cope with the uncertainties associated with rainfed agriculture. In conclusion, the study provides valuable evidence supporting the adoption of groundnut-based intercropping with foxtail millet as an effective strategy to enhance productivity, increase net returns, and improve production sustainability under rainfed conditions. By diversifying and intensifying their cropping systems, farmers can mitigate risks and

achieve more stable and profitable outcomes in challenging environments characterized by low and erratic rainfall.

UNDER PEER REVIEW

Table 1: Groundnut pod yield, millet grain yield, groundnut pot equivalent yield and economics of groundnut and millet based intercropping systems under rainfed conditions

Treatments	Groundnut Pod Yield (kg/ha)			Mean	Millet Grain Yield (kg/ha)	GPEY (kg/ha)	LER	Gross returns (Rs/ha)	Net returns (Rs/ha)	B:C ratio
	2017	2018	2019							
T ₁ : Sole groundnut	1862	2565	2721	2383	-	2383	1.00	111838	54766	1.96
T ₂ : Sole finger millet	-	-	-	-	2257	1174	1.00	51316	24103	1.89
T ₃ : Sole little millet	-	-	-	-	1531	757	1.00	35889	13314	1.59
T ₄ : Sole foxtail millet	-	-	-	-	1870	758	1.00	35454	13195	1.59
T ₅ : Groundnut + finger millet (5:2)	1260	1517	1620	1475	1158	2124	1.20	97371	56935	2.41
T ₆ : Groundnut + little millet (5:2)	1334	1634	1745	1571	754	1989	1.22	90992	53431	2.42
T ₇ : Groundnut + foxtail millet (5:2)	1486	1656	1768	1636	900	2060	1.24	91709	53726	2.41
T ₈ : Groundnut + finger millet (6:1)	1590	1723	1840	1717	671	1971	0.99	93693	53257	2.32
T ₉ : Groundnut + little millet (6:1)	1683	1850	1975	1836	482	1972	1.04	93805	56244	2.50
T ₁₀ : Groundnut + foxtail millet (6:1)	1744	1910	2039	1897	575	2015	1.05	96237	58532	2.55
S.Em \pm	95	156	6.86	86	76	124	0.05			
C.D. ($P=0.05$)	286	479	21.14	260	228	372	0.17			

Table 2: Yield and economics farm trial at different locations

Technology	Groundnut + Foxtail millet (6:1)			Sole groundnut		
	2020	2021	Pooled	2020	2021	Pooled
Groundnut Pod yield (kg/ha)	1675	1663	1669	1901	1803	1852
Foxtail millet yield (kg/ha)	884	1011	948	-	-	-
GPEY (kg/ha)	2050	2082	2066			
Gross returns (Rs. /ha)	107994	111217	109606	99953	95034	97494
Net returns (Rs. /ha)	67691	70626	69159	51456	48134	49795
B:C ratio	2.67	2.77	2.72	2.07	2.03	2.05

4. CONCLUSION

In conclusion, the study confirms that cultivating crops as sole crops under rainfed conditions is risky due to the adverse effects of low and erratic rainfall, leading to reduced productivity. To achieve higher productivity in such challenging conditions, it is essential to adopt intensification and diversification of crops. Intercropping of groundnut + foxtail millet (6:1) was found profitable and efficient intercropping system.

The farm trial reveals that adopting an intercropping system with groundnut and foxtail millet resulted in a 10.35 % increase in groundnut pod equivalent yield as compared to sole groundnut. This intercropping approach also led to significantly higher net returns representing a substantial 29 % increase compared to the net returns from normal sole groundnut cultivation. Moreover, the intercropping treatment with groundnut and foxtail millet (6:1) showed a higher benefit-cost ratio (B:C ratio) of 2.72, indicating greater economic profitability compared to the B:C ratio of 2.05 observed in sole groundnut cultivation. This suggests that intercropping has economic viability and potential benefits, making it a promising approach to enhance agricultural productivity and financial returns.

REFERENCES

1. Anonymous. Agricultural Statistics at a Glance 2021. Government of India Ministry of Agriculture & Farmers Welfare Department of Agriculture & Farmers Welfare Directorate of Economics & Statistics. 2022; pp. 82-83.
2. Willey RW. Intercropping its importance and research needs 1. Competition and yield advantage and 2. Agronomy and Research Approaches. Field Crop Abstracts, 1979;32: 73-85.
3. Gomez KA and Gomez AA. Statistical Procedure for Agriculture Research, 2nd Ed., John Willey and Sons, New York, 1984; p. 68.
4. Shwethanjali KV, Kumar Naik AH, Basavaraj Naik T and Dinesh Kumar M. Effect of groundnut-based millets intercropping system on growth and yield of groundnut (*Arachis hypogaea* L.) under rainfed condition. International Journal of Agriculture Sciences. 2018;10(17): 7033-7034.
5. Maitra S, Ghosh DC, Sounda G, Jana PK and Roy DK. Productivity, competition and economics of intercropping legumes in finger millet (*Eleusine coracana*) at different fertility levels. Indian Journal of Agriculture Sciences. 2002;70(12): 824-828.
6. Bassi JA and Dugje IY. Benefits of intercropping selected grain legumes with pearl millet in Nigerian Sudan Savannah. International Journal of Advance Agricultural Research. 2016;4: 65-77.