

Acute Oral Toxicity and Antispasmodic Effects of Two Extracts of *Diospyros mespiliformis* Hochst. ex A. DC. (Ebenaceae) Immature Fruit on the Isolated Rat Duodenum

ABSTRACT

Introduction : *Diospyros mespiliformis* Hochst. ex A. DC. (Ebenaceae) is an artisanal, nutritional, and medicinal plant. It is used in Burkina Faso for the treatment of gastrointestinal disorders.

Aims: Our study aimed to evaluate the chemical profile, the antioxidant and anti-inflammatory activities, the spasmolytic effects, and the safety of use of hydroethanolic extract (HE_FDM) and aqueous macerate extract (AM_FDM) of *Diospyros mespiliformis* immature fruits.

Place and Duration of Study: Immature fruits of *Diospyros mespiliformis* were collected in August 2022 in Goundi, in the Sanguié Province of Burkina Faso. The experiments were conducted in September - December 2022 at Laboratoire de Recherche-Développement de Phytomédicaments et Médicaments/IRSS/CNRST, Ouagadougou.

Methodology: Phytochemical screening by HPTLC and the determination of total polyphenols and flavonoids were carried out. The antioxidant activity was evaluated according to DPPH, ABTS, FRAP, and LPO methods. The inhibitory activity of 15-lipoxygenase and Phospholipase A2 was evaluated. The *ex vivo* antispasmodic effect of the extracts was tested on isolated rat duodenum using ACh and BaCl₂ as contracting agents. Finally, acute oral toxicity was carried out on female mice (NMRI).

Results: At the end of these tests, the extracts contain tannins, flavonoids, sterols, triterpenes, and saponosides. HE_FDM gave the highest levels of total phenolics (71.32±2.26 mg TAE/g and flavonoids (16.37±0.24 mg QE/g). Of the two extracts, HE_FDM gave the best antioxidant activity. However, these activities were lower than the reference substances (p<0.05). The extracts have a moderate inhibitory effect on phospholipase A2 and 15-Lipoxygenase. In addition, the most active HE_FDM caused a spasmolytic effect with E_{max} of 98.87±5.14% and 81.61±7.56%, respectively, during contractions induced by BaCl₂ and ACh. Finally, the extracts are practically non-toxic.

Conclusion: Finally, this work provided scientific data and could justify the use of *D. mespiliformis* immature fruits in the treatment of gastrointestinal disorders.

Keywords: *Diospyros mespiliformis*, Immature fruits, Antioxidant, Antispasmodic, Safety of use

1. INTRODUCTION

Gastrointestinal disorders are one of the most common illnesses in humans. These most common conditions affect the greatest number of children and adolescents[1]. To deal with these gastrointestinal disorders, people regularly turn to alternative medicine. In fact, traditional medicine remains the first line of health care for 80% of the population in developing countries [2]. Plants are a widely exploited source of bioactive compounds in alternative and modern medicine. They constitute a useful matrix for identifying phytochemical compounds, which are then optimized by the pharmaceutical industry [3]. Burkina Faso offers a great botanical treasure trove, given its geographical and climatic diversity, and is a major source of potentially therapeutic plants for the treatment of diseases with gastrointestinal spasmodic components[4]. Antispasmodics or spasmolytics are drugs used to treat muscle spasms, which are intense and sudden pathological contractions of involuntary or smooth muscles [5]. On a pre-clinical level, ethnobotanical, biochemical, pharmacological, and toxicological studies and clinical trials have been carried out to reinforce this cultural heritage and provide scientific evidence[6, 7]. In view of this cultural dimension, it is vital to enhance the value of medicinal plants by stepping up research into plant species widely used for their medicinal and especially spasmolytic properties. Among these potentially antispasmodic plants is *Diospyros mespiliformis*Hochst. ex A. DC (Ebenaceae)[8]. The literature tells us that extracts of *Diospyros mespiliformis* trunk bark have antiproliferative properties [9]. Leaf extracts have antioxidant and antimicrobial properties [8, 10]. The anti-plasmodial activity of methanolic fractions and extracts of trunk bark and leaves has also been demonstrated [8, 11]. In addition, the anti-inflammatory and spasmolytic properties of leaves have recently been demonstrated [4]. However, given the traditional use of the fruits of this plant, the present study was carried out to examine the safety and possible antispasmodic activity of the aqueous and hydroethanolic extracts of the immature fruits on acetylcholine (Ach)- and BaCl₂-induced contractions of the isolated rat duodenum.

2. MATERIAL AND METHODS

2.1 Plant Material

Immature fruits of *Diospyros mespiliformis*Hochst. ex A. DC (Ebenaceae) were collected in May 2022 in Goundi, in the Sanguié Province of Burkina Faso. One sample (4267 OUA) was identified and authenticated by a botanist from the Plant Biology and Ecology Laboratory at Joseph KI-ZERBO University, Burkina Faso. Fruits cut into small pieces were dried in the shade, away from direct sunlight and with ventilation. After drying, the fruits were pulverized using a mechanical grinder (Gladiator Est. 1931 Type BN 1 Mach. 404611083) to obtain a dry extractive powder.

2.2 Preparation of Fruits Extracts

Extraction was carried out using the maceration exhaustion technique[12]. One hundred (100) g of immature fruit powder was macerated in 500 mL of distilled water or 80% ethanol (absolute ethanol/water; 80/20; v/v) for 24 h at room temperature. After 24 h, each residual pomace was repeated twice under the same conditions with distilled water or 80% ethanol. The two supernatants were filtered through fine mesh nylon cloth and centrifuged at 2000 rpm for 5 min. The hydroethanolic filtrate was concentrated in a rotavapor (ventilated Electronic Microprocessor Controller CPS) under reduced pressure at a constant temperature of 60°C, then frozen and freeze-dried. The filtrate of the aqueous macerate was frozen and freeze-dried. The

hydroethanolic extract (HE_FDM) and the aqueous extract (AM_FDM) were then recovered and stored in two hermetically sealed bottles for the various tests.

2.3 Phytochemical Investigation: High-Performance Thin-Layer Chromatography

High-performance thin-layer chromatography (HPTLC) was used to detect flavonoids and tannins in the two extracts (HE_FDM and AM_FDM). It was carried out on chromatoplates (60 F₂₅₄, 10 x 5 cm, glass support 10 x 20 cm, Macherey-Nagel, GermanyMerck) following the literature[3]. Approximately 20 µL of each extract was streaked with a semi-automatic sample dispenser (CAMAG, Linomat 5, Switzerland) along the baseline 0.8 cm from the bottom edge of the plate. After deposition and drying, the plates were placed in a tank containing eluent previously saturated (20× 10cm, saturation time: 30 min). The solvent system used depended on the metabolite to be identified: ethyl acetate/formic acid/H₂O (18/2/4/2/1 v/v/v/v) for tannins, ethyl acetate/formic acid/H₂O, (8/2/1 v/v/v/) for flavonoids, ethyl acetate/hexane (8/2 v/v) for sterol-triterpenes and hexane/ethyl acetate/methanol (10/5/5 v/v/v) for saponosides. After migration over 0.8 dm in length, the plates were dried, and sulphuric anisaldehyde reagent for saponosides, Neu reagent for flavonoids, Liebermann and Burchard reagent for Sterol-triterpenes and 5% FeCl₃ for tannins revealed the chromatographic profiles. The profiles were then observed under visible light (tannins).

2.4 Determination of Total Phenolic Compounds

The total phenolic content of HE_FDM and AM_FDM was determined using the Folin-Ciocalteu Reagent (FCR) [12]. The reference compound (Gallic acid) was used as to produce the standard curve. Twenty-five(25) µL of sample at a concentration of 1 mg/mL was mixed with 125 µL of FCR. 100 µL of 7.5% w/v sodium carbonate solution was added to the mixture. After one hour, absorbance at 760 nm was measured using a microplate reader (Spectro UV, Epoch Biotek, USA). Results were expressed as mg Gallic acid equivalent (GAE)/g dry extract.

2.5 Determination of Flavonoid Compound

Flavonoid content was assessed using an aluminum chloride reagent [12]. A standard calibration curve was plotted with Quercetin. one hundred(100) µL of HE_FDM or AM_FDM(1 mg/mL) were mixed with 100 µL of a 2% w/v Aluminium trichloride solution. After 10 min, absorbance was measured using a mass spectrophotometer. Absorbance at 415 nm was measured using a spectrophotometer (Epoch Biotek, USA) after 10 min. Results were expressed as mg Quercetin equivalent (QE)/g dry extract.

2.5 Assessment of Antioxidant Properties

2.5.1 DPPH• Essay

Free radical scavenging activity of HE_FDM, AM_FDM, and Trolox was performed using 2,2-diphenyl-1-picrylhydrazyl (DPPH) as previously described[13]. The absorbance of 10 µL of samples and Trolox added to 200 µL of DPPH (0.04 mg/mL) was measured at 490 nm after 30 min of incubation in the dark at room temperature using a Bio-Rad spectrophotometer (model 680, Japan). The result was expressed as antioxidant capacity equivalent to Trolox. As a function of sample concentration, a DPPH•-inhibition percentage curve was plotted. The percentage inhibition of the DPPH radical was calculated using the following formula:

Inhibition (%) = $[(A_c - A_e)/A_c] \times 100$; A_e and A_c represent the absorbances of the extract/ascorbic acid and the control (DPPH solution without sample). The concentration required to inhibit 50% of DPPH (IC_{50}) was determined on the curve.

Anti-radical power (ARP) was determined by the formula: $ARP = 1/IC_{50}$; ARP: Anti Radical Power; IC_{50} : 50% inhibitory concentration expressed in $\mu\text{g/mL}$

2.5.2 Ferric Reducing Antioxidant Power (FRAP) Assay

The FRAP assay was performed on HE_FDM, AM_FDM and Trolox as previously described [13]. The mixture of 0.5 mL samples with 1.25 mL phosphate buffer and 1.25 mL potassium hexacyanoferrate aqueous solution (1%) was incubated for 30 min at 50°C . Next, 1.25 mL of trichloroacetic acid (10%) was added and centrifuged at $3000 \times g$ for 10 min. FeCl_3 solution (0.125 mL, 0.1%), and distilled water (0.625 mL) were added to the supernatant (0.625 mL). The absorbance of the reaction medium was measured at 700 nm using a spectrophotometer (Agilent, Santa Clara, CA) equipped with ChemStation UV-visible software. Trolox was used to plot the calibration curve. The FRAP activity of HE_FDM and AM_FDM was expressed as mol Trolox equivalent/gram dry extract.

2.5.3 ABTS \cdot^+ Assay

The ABTS free radical scavenging activity of HE_FDM, AM_FDM and Trolox was assessed using the procedure described previously [12]. In a volume of 5 mL of distilled water, 19.2 mg of ABTS were dissolved. Potassium persulphate (3.312 mg) was added to the ABTS solution (3.84 mg/mL). After adding the potassium persulphate, the solution was left for 16 h in the dark at room temperature before use. On the day of the experiment, 4.5 mL of the mixture was diluted in 220 mL of absolute ethanol. The range of 8 dilutions to be tested was prepared from the parent concentration of the samples (1 mg/mL). On a 96-well microplate, 200 μL of ABTS solution mixed with 20 μL of HE_FDM or AM_FDM or Trolox were added to each well. After incubation for 30 min at 25°C , absorbances were read against a blank at 415 nm using an Agilent 8453 spectrophotometer with ChemStation UV-visible software. Measurements were performed in triplicate. The percentage inhibition of absorbance at 415 nm was calculated according to the formula:

$$\% \text{ Inhibition} = [(A_0 - A) / A_0] \times 100$$

A_0 is the absorbance of the control; A is the absorbance of the sample.

The absorbance inhibition curve as a function of the concentration of the extract or reference substance (Trolox) was constructed to determine the 50% inhibitory concentration (IC_{50}). Anti-radical power (ARP) was determined using the formula: $ARP = (1/CI_{50})$; ARP: Anti-radical power; IC_{50} : 50% inhibitory concentration expressed in $\mu\text{g/mL}$

2.5.4 Lipid Peroxidation Inhibition (LPO) Assay

The lipid peroxidation activity of rat liver was determined using 2-thiobarbituric acid (Sigma Aldrich) [12]. $\text{FeCl}_2 \cdot \text{H}_2\text{O}_2$ was used to induce peroxidation of liver homogenate. A 0.2 mL volume of HE_FDM or AM_FDM (1.5 mg/mL) was mixed with 1 mL of 1% rat liver homogenate, then 50 μL FeCl_2 (0.5 mM) and 50 μL H_2O_2 (0.5 mM) were added. The mixture was incubated for 60 min at 37°C , then 1 mL trichloroacetic acid (15%) and 1 mL 2-thiobarbituric acid (0.67%) were added. The mixture was heated in boiling water for 15 min. The experiment was performed in triplicate and the absorbance was read at 532 nm. Trolox was used as the reference product. The percentage inhibition was calculated using the following formula: Percentage inhibition (%) = $[1 - (A_1 - A_2)/A_0] \times 100$

A_1 is the absorbance of the control (without sample); A_2 is the absorbance with the sample; A_0 is the absorbance without liver homogenate.

UNDER PEER REVIEW

2.5 Anti-Inflammatory Activity

2.6.1 Phospholipase A2 (sPLA2) Inhibition Assay

The sPLA2 activity of bee venom was determined according to the instructions of the manufacturer Abcam (Japan) described in catalog no. ab133089[12]. A 96-well microplate was used to perform the sPLA2 inhibition assay. A final concentration of 100 µg/mL of HE_FDM, AM_FDM and Betamethasone (reference compound) was used. Absorbances were read spectrophotometrically (Agilent 8453) at 415 nm against a blank that had not received the enzyme. The experiment was performed in triplicate and the percentage inhibition of sPLA2 at 100 µg/mL was calculated using the following formula: Inhibition (%) = [(AE - AI)/AE]x100.

AE: Absorbance of enzyme assay - Absorbance of blank; AI: Absorbance of inhibition assay - Absorbance of blank.

2.6.2 Lipoxygenase Inhibition Assay

Lipoxygenase inhibition was determined using linoleic acid (1.25 mM) as substrate [12]. Inhibitors (HE_FDM or AM_FDM/ reference substance: Zileuton) were prepared to a final concentration of 100 µg/mL. 146.25 µL of 15-lipoxygenase solution (820.51U/mL) was added to 3.75 µL of each inhibitor. Next, 150 µL of linoleic acid was added. A spectrophotometer (Epoch Biotek Instruments, USA) was used to measure absorbances at 234 nm against enzyme-free blanks. The tests were carried out in triplicate and the percentage of lipoxygenase inhibition was calculated using the formula:

Inhibition(%) = [(AE - AI)/AE]x100

AE: Absorbance enzyme test - Absorbance blank; AI: Absorbance inhibition test - Absorbance blank.

2.7 Experimental Animals

Female NMRI (Naval Medicinal Research Institute) mice and Wistar male rats with average weights of 28±3 g and 205±15 g, respectively from the animal house of the “Institut de Recherche en Sciences de la Santé/Centre National de la Recherche Scientifique et Technologique (IRSS/CNRST), Burkina Faso” were used. The animals were placed in an enclosure at a temperature of 21-23 °C with relative humidity between 50 and 60%, and subjected to the light/dark cycle of 12 h/12 h according to the rearing conditions of these species. Standard laboratory pellets enriched with protein (29%) and free access to water in bottles were provided to the animals [12, 14].

2.8 Spasmolytic Effect of Two Extracts of *D.mespiliformis* Fruitson Isolated Rat Duodenum

The protocol used has been described by Nitiéma et al.[4]. Wistar rat is fasted for 24 h before the start of the experiment and then sacrificed. Part of the duodenum is removed and immediately placed in Tyrode's oxygenated physiological solution [KCl (0.2g), NaCl (8g), MgCl₂ (0.01g), NaH₂PO₄ (0.05g), NaHCO₃ (1g), CaCl₂ (0.2g), and Glucose (1g) in 1000mL of distilled water]. A fragment (1.5 cm) was freed of adhesions and mounted in the isolated organ bath thermostated at 37°C, with a pneumatic bubbler for organ oxygenation. One end of the isolated intestine fragment is attached to the hook of the support rod, and the other end to the isometric transducer, which in turn is connected to the recorder via an amplifier. This device visualizes the contractions of the isolated rat duodenum. The Tyrode solution is renewed every 15 min during the 45 min stabilization period. KCl (80 mM) is administered into the organ vessel to stimulate the duodenum after observing the regularity of its contractile activity, followed by rinsing. Solutions

of HE_FDM and AM_FDM are administered after precontraction with Acetylcholine (ACh, 10^{-6} M) or Barium chloride (BaCl_2 , 160 $\mu\text{g/mL}$). This makes it possible to assess, respectively, the extract's effect on normal contractile activity in the isolated duodenum and its interaction with the cholinergic system and potassium fluxes in the cells. The percentage inhibition of contraction (PI) is calculated using the following formula:

$$\text{PI} = (h_1 - h_2 / h_1) \times 100$$

h_1 : height of peaks due to contractor alone; h_2 : height of peaks due to contractor in the presence of extract.

2.9 Acute Oral Toxicity

The acute oral toxicity test for HE_FDM and AM_FDM was performed on female NMRI (Naval Medicinal Research Institute) mice in accordance with OECD guideline 423 [15, 16]. Two batches, each consisting of three mice, were made up and placed separately in polypropylene cages, a control batch of 03 mice and a test batch of 03 mice. HE_FDM or AM_FDM was administered by gavage using an esophageal tube in a single dose to the test mice after fasting for 3 h. A dose of 2000 mg/kg body weight of HE_FDM and AM_FDM was chosen as the starting dose. Control batch received the solvent for dissolving the extract (distilled water, 10 mL/kg). The mice were observed individually for 2 h after administration, at the end of which food was restored. They were then observed twice daily for a period of 14 days to monitor for mortality and signs of toxicity such as bleeding, excitement, convulsions, vomiting, salivation, hyperventilation, diarrhea, lack of appetite, sleep, and coma. The weight of each mouse and the quantities of water and food consumed were measured every 2 days during 14 days of experimentation. On day 15, the mice were sacrificed and a necropsy was performed on the organs (liver, kidneys, lungs, spleen, and heart) then weighed. The relative weight of each organ was calculated [(Organ weight (g) / Fasting mouse weight on the day of sacrifice (g)) $\times 100$]. This test was repeated after the mice were sacrificed under the same conditions as the 2 other batches (control batch and extract batch).

2.10 Statistical Analysis

Values are given as arithmetic means \pm SEM, and means \pm SD. The concentration-response curves were constructed using GraphPad Prism 8.4.3 Software, San Diego, CA. Two pharmacological parameters were obtained: the maximal effect generated by the agonist (E_{max}) and a concentration of agonist producing 50% of the maximum response (EC_{50}). Statistical comparisons were performed using Student's T-test, one- and two-way ANOVA. Post hoc test was performed using Bonferroni's test analysis to compare all the groups. A p -value < 0.05 was considered as statistically significant.

3 RESULTS

3.1 HPTLC Phytochemical Investigation

The phytochemical analysis of HE_FDM and AM_FDM highlighted the presence of saponosides, tannins, flavonoids, and sterol-triterpenes (Table 1).

Table 1: Phytochemical profile of hydroethanolic extract (HE_FDM) and aqueous macerate extract (AM_FDM) of *D. mespiliformis* fruits

Extracts	Saponosides	Tannins	Flavonoids	Sterol-triterpenes
HE_FDM	+	+	+	+
AM_FDM	+	+	+	-

+: presence of compound; -: compound not detected

3.2 Total Phenolic and Flavonoids Contents in Extracts of *D. mespiliformis* Immature Fruits

The total phenolics and flavonoid contents in extracts of *D. mespiliformis* immature fruits are shown in Table 2. These results are expressed in milligrams of quercetin equivalent per gram of dry extract (mg QE/g) for flavonoid, and in milligrams of tannic acid equivalent per gram of dry extract for total phenolics (mg TAE/g). HE_FDM and AM_FDM showed the similar value of Total phenolics and flavonoids compound content.

Table 2 : Total phenolics and flavonoid contents in extracts of *D. mespiliformis* immature fruits

Extracts	Total phenolics(mg TAE/g)	Flavonoids (mg QE/g)
HE_FDM	71.32±2.26	16.37±0.24
AM_FDM	61.23±0.67	15.83±0.21

QE: quercetin equivalent; TAE: tannic acid equivalent

3.3 Antioxidant Activity

The antioxidant activity of *D. mespiliformis* immature fruits extracts are shown in Table 3. HE_FDM and AM_FDM showed lower antioxidant activity with DPPH and ABTS than Trolox (*p<0.05). The IC₅₀ for Trolox were 6.34±0.04 µg/mL and 3.78±0.21 µg/mL respectively for DPPH and ABTS. The ferric ion reduction capacity (FRAP) showed similar values for both extracts. The lipid peroxidation inhibitory power (LPO), expressed as a percentage (at 100 µg/mL) of the two extracts was comparable to that of Trolox, 48.11±3.88% (p>0.05).

Table 3: *In vitro* antioxidant activity of *D. mespiliformis* immature fruits extracts

Extracts	ABTS		DPPH		FRAP	LPO
	IC ₅₀ (µg/mL)	AR P	IC ₅₀ (µg/mL)	ARP	mol EAA/g	Inhibition (%) (at 100 µg/mL)
HE_FDM	6.73±0.46*	0.15	13.82±1.04**	0.07	1149.35±13.43	41.86±5.12
AM_FDM	33.87±0.86***	0.03	83.07±6.40***	0.01	1091.95±5.12	60.75±5.09
Trolox	3.78±0.21	0.26	6.34±0.04	0.16	-----	48.11±3.88

HE_FDM: hydroethanolic extract; AM_FDM: aqueous macerate extract; IC₅₀: inhibition concentration 50%; ARP: anti-free radical power; n = 3; *p<0.05; **p<0.01; ***p<0.001 vs Trolox; EAA: Ascorbic acid equivalent.

3.3 Anti-inflammatory Activity *in vitro*

The evaluation of the *in vitro* anti-inflammatory activity of the two extracts by inhibiting 15-lipoxygenase and Phospholipase A₂ is recorded in Table 4. HE_FDM, and AM_FDM have similar effects in terms of inhibition on 15-lipoxygenase. However, the Zileuton presented a better IC₅₀, 2.92±0.32 µg/mL (p<0.001). Phospholipase A₂ activity of HE_FDM and AM_FDM showed that the extracts had a low inhibitory power compared to Betamethasone, 35.39±3.31% (p<0.001).

Table 4: 15-Lipoxygenase and Phospholipase A₂ inhibitory activity of *D. mespiliformis* immature fruits extracts

Fractions	15-Lipoxygenase IC ₅₀ (µg/mL)	Phospholipase A ₂ Inhibition (%)
HE_FDM	15.11±1.39***	5.77±0.63###
AM_FDM	19.12±2.01***	5.92±0.59###
Zileuton	2.92±0.32	----
Betamethasone	----	35.39±3.31

HE_FDM: Hydroethanolic extract; AM_FDM: aqueous macerate extract; ***p<0.001 vs Zileuton; ###p<0.001 vs Betamethasone; n = 3

3.4 Antispasmodic Effect of two extracts of *D. mespiliformis* Immature Fruits on Rat Duodenum

Relaxation of isolated rat duodenum by HE_FDM and AM_FDM is shown in Figure 1. Figure 1A shows the relaxation curves for both extracts and control on the isolated duodenum precontracted with ACh. HE_FDM curve was more deviated to the left compared to AM_FDM and control. Figure 1B shows the E_{max} of HE_FDM (98.87±5.14%), AM_FDM (64.28±10.75%) and control (7.08±4.68%) (p<0.05). Figure 2A shows similar relaxation results obtained with the extracts and the vehicle (control) on the isolated rat duodenum pre-contracted with BaCl₂. With a statistically significant difference, the best efficacy of 81.61±7.56% was obtained with the HE_FDM (Figure 2B).




Figure 1: Relaxation curves of Hydroethanolic extract (HE_FDM) and aqueous macerate extract (AM_FDM) of *D. mespiliformis* fruits on isolated ACh-precontracted rat duodenum (A) and histogram of maximum relaxation effect (B) ; ***p<0.001 vs Control; φφφp<0.001 vs AM_FDM; n = 5




Figure 2: Relaxation curves of Hydroethanolic extract (HE_FDM) and aqueous macerate extract (AM_FDM) of *D. mespiliformis* fruitson isolated BaCl₂-precontracted rat duodenum (A) and histogram of maximum relaxation effect (B) ; n = 5 ; ***p < 0.001 vs Control; φφφp<0.001 vs AM_FDM

3.5 Acute Oral Toxicity

3.5.1 Animal Behaviour

The dose of 2000 mg/kg body weight showed no signs of mortality or remarkable behavioral changes in female mice at the first and second stages of administration of HE_FDM, and AM_FDM (Table 5).

Table 5: Mortality of female mice administered a single dose of *D. mespiliformis* immature fruits extracts

Extract administered	Mortality	
	1 st test	2 nd test
Control (distilled water)	0/3	0/3
HE_FDM(2000 mg/kg)	0/3	0/3
AM_FDM(2000 mg/kg)	0/3	0/3
Excitement	--	--
Sleepiness	--	--
Hair standing up	--	--
Lack of appetite	--	--
Diarrhea	--	--
Hyperventilation	--	--
Convulsions	--	--
Bleeding	--	--
Coma	--	--

HE_FDM: hydroethanolic extract; AM_FDM: aqueous macerate extract

3.5.2 Evolution in body weight, feed, and water consumption of mice after 14 days of monitoring

Figure 3 shows the mean weight gain, feed consumption, and water consumption for 2 weeks in female mice given vehicle (distilled water, 10 mL/kg) or the single dose (2000 mg/kg) of HE_FDM, and AM_FDM. There was no statistically significant difference between the parameters of the 03 study batches.




Figure 3: Changes in weight (A), water (B) and feed (C) consumption of female mice from control and test batches with extracts of *D. mespiliformis* immature fruits during 2 weeks of follow-up; HE_FDM: hydroethanolic extract; AM_FDM: aqueousmacerate extract; n = 6

3.5.3 Macroscopic Observation and Relative Organ Weights of Mice

Fresh macroscopic examination of vital organs such as the lungs, liver, heart, kidneys and spleen of control, HE_FDM-treated and AM_FDM-treated mice showed that there were no visually apparent changes in color or appearance or lesions of the various organs. In addition, Table 6 shows the relative organ weights of batches of control mice and mice treated with the two extracts. No statistically significant variation was observed between the relative organ weights.

Table6: Relative weights of female mice from control and test batches with *D. mespiliformis* immature fruits extracts during 2 weeks of follow-up; n = 6

Substances	Relative organ weight (mean±SD)				
	Heart	Kidneys	Lungs	Liver	Spleen
Control (Distilled water)	0.52±0.03	1.17±0.03	0.80±0.04	5.08±0.3	0.42±0.03
HE_FDM (2000 mg/kg)	0.51±0.05	1.22±0.04	0.79±0.06	5.15±0.16	0.45±0.03
AM_FDM (2000 mg/kg)	0.48±0.09	1.30±0.08	0.83±0.07	4.96±0.38	0.43±0.06

HE_FDM: hydroethanolic extract; AM_FDM: aqueousmacerate extract

4. DISCUSSION

Plants contain a multitude of bioactive compounds and as a result have been used for thousands of years in the treatment of numerous pathologies. The treatment of gastrointestinal disorders is no exception [17]. The leaves of *Diospyros mespiliformis* are widely used as an antispasmodic [18]. The phytochemical screening of the two extracts from the immature fruits of the plant is comparable to that reported in studies showing that *D. mespiliformis* immature fruits contain flavonoids, tannins, sterols, terpenes and saponosides [4, 19]. Indeed, these phytochemicals are endowed with antioxidant properties that can trap free radicals and block the implications of oxidative stress [12]. The antioxidant properties of methanolic fruit extract have been demonstrated and our results corroborate these data [20]. In addition, according to the literature, flavonoids have anti-inflammatory and antispasmodic properties. Recent studies have shown that, like fruit extracts, the ethyl acetate fraction (84.15±1.73 mg TAE/g) and the residual aqueous fraction (89.67±2.35 mg TAE/g) of plant leaves contain total phenolics. These compounds are known for their spasmolytic properties [4, 21, 22]. Furthermore, the phenolic compounds, in particular tannins, flavonoids and triterpenes present in the two extracts in the study are inhibitors of pro-inflammatory enzymes involved in the production of free radicals and the regulation or protection of the antioxidant defense system [23, 24]. In addition, increased spasmodic activity and diarrhea activate inflammatory pathways. Interestingly, the fruit extracts have anti-inflammatory activity albeit moderated by inhibition of phospholipase A2 and 15-lipoxygenase. These results confirm the anti-inflammatory properties of the plant, especially the leaves [4, 25]. Indeed, the flavonoids and sterols/triterpenes present in HE_FDM and AM_FDM are known for their ability to inhibit pro-inflammatory enzymes [12, 26]. Pharmacological investigations showed that HE_FDM and AM_FDM have relaxing properties on isolated rat duodenum by inhibiting the stimulatory action of ACh on muscarinic receptors. Of the two extracts, HE_FDM was the more effective. ACh induces contraction by activating G protein-coupled smooth muscle M3 receptors, leading via Inositol Triphosphate (IP₃) to the release of intracellular Ca²⁺. Its action causes a significant positive inotropic effect marked by contractile activity reflecting the increased peristalsis of the gastrointestinal tract [27, 28]. This curative effect shows that HE_FDM and AM_FDM have anticholinergic properties. In fact, the tannins, flavonoids, saponosides and sterol-triterpenes in HE_FDM and AM_FDM, known for their anti-diarrheal and spasmolytic properties, are thought to inhibit the action of ACh [29]. Work on *D. mespiliformis* leaves fractions also showed spasmolytic effects on an isolated rat duodenum, with an efficacy for the ethyl acetate fraction (E_{max} = 90.40±7.84%) compared with the residual aqueous fraction [4]. In addition, HE_FDM and AM_FDM have voltage-dependent calcium channel antagonist activity through inhibition of BaCl₂-induced smooth muscle contraction. Indeed, both extracts at concentrations of 0.03-10 mg/mL induced a relaxing effect on the rat duodenum by significantly reducing contractions in a concentration-dependent manner. These extracts are thought to have a similar action to papaverine with a musculotropic effect, by inhibiting phosphodiesterase [30]. In addition, the elevated cytoplasmic Ca²⁺ concentration in smooth muscle cells is the main stimulus for duodenal contraction. This results from influx of extracellular Ca²⁺ and/or intracellular release of stored Ca²⁺ [31]. Interestingly, the concentration-dependent spasmolytic effect of HE_FDM and AM_FDM on the contractile activity of the isolated rat duodenum may be the result of Ca²⁺ uptake by phosphorylated proteins under the influence of protein kinase activated by increased levels of adenosine 3,5-cyclic monophosphate (cAMP). It may also be due to an inhibition of calcium influx or an increase in calcium efflux without altering influx [4, 22, 30]. For their proven spasmolytic properties, HE_FDM is the most

effective even compared to the plant leaf fractions. The safety of a pharmaceutical product is a very important criterion in therapeutics. For this reason, HE_FDM and AM_FDM were evaluated for acute oral toxicity. The results showed that their LD₅₀ is estimated at 5000 mg/kgbw according to the United Nations Globally Harmonised System with demonstrated adverse effects [16]. These results show that the immature fruits of the plant are practically non-toxic and can therefore be used appropriately. In line with the literature, studies have shown that fractions and extracts from the leaves and bark of the trunk of *Diospyros mespiliformis* can be consumed without risk [4, 32]. However, the mechanism of action of these fractions requires further investigation, in particular the bio-guided evaluation of their *in vivo* anti-diarrheal properties and their medium- and long-term oral toxicity.

5. CONCLUSION

This study provides evidence that *Diospyros mespiliformis* immature fruits can be used safely to treat intestinal spasms and diarrhea. This finding is the first to our knowledge to pharmacologically demonstrate the spasmolytic effects of the immature fruits of this plant. The results showed that hydroethanolic and aqueous extracts of *D. mespiliformis* immature fruits inhibit ACh- and BaCl₂-induced contractions of the intestinal muscle. Tannins, saponosides, flavonoids, sterols and terpenoids are thought to be responsible for the antioxidant, anti-inflammatory and spasmolytic properties. The present study would therefore justify the traditional use of this plant for gastrointestinal symptoms. However, there is still a need to understand the mechanisms involved and to ensure that extracts are safe in the medium and long term.

CONSENT

Not applicable

ETHICAL APPROVAL

Experiments were conducted in accordance with the procedures of the Helsinki Declaration Guide to Good Practice in Animal Experimentation. In addition, experimental procedures on mice and rats were carried out in accordance with the Guide for the Care and Use of Laboratory Animals of the US National Institutes of Health and with Directive 2010/63/EU on animal experimentation.

REFERENCES

1. Akinmurele OJ, Sonibare MA, Elujoba AA, Ogunlakin AD, Yeye OE, Gyebi GA, Ojo OA, Alanzi AR: Antispasmodic Effect of *Alstonia boonei* De Wild. and Its Constituents: Ex Vivo and In Silico Approaches. *Molecules* 2023, 28(20):7069.
2. Khan MSA, Ahmad I: Herbal medicine: current trends and future prospects. In: *New look to phytomedicine*. edn.: Elsevier; 2019: 3-13.
3. Ouedraogo WRC, Belemnaba L, Nitiéma M, Kaboré B, Koala M, Ouedraogo S, Semde R, Ouedraogo S: Phytochemical study, antioxidant and vasodilatation activities of leafy stem extracts of *Flemingia faginea* Guill. & Perr.(Barker), a medicinal plant used for the

- traditional treatment of arterial hypertension. *Pharmacological Research-Modern Chinese Medicine* 2023, 7:100231.
4. Nitiéma M, Ouédraogo WRC, Rainatou B, Ouédraogo PE, Noura OM, Kaboré B, Bélem-Kabré WLME, Traoré TK, Compaoré S, Koala M: Phytochemical profile, acute oral toxicity, antioxidant, and antispasmodic effects of ethyl acetate and aqueous residual fractions of *Diospyros mespiliformis* Hochst. ex A. DC (Ebenaceae) leaves on isolated duodenum of rat. *Journal of Drug Delivery and Therapeutics* 2023, 13(12):148-154.
 5. Heghes SC, Vostinaru O, Rus LM, Mogosan C, Iuga CA, Filip L: Antispasmodic effect of essential oils and their constituents: A review. *Molecules* 2019, 24(9):1675.
 6. Bencheikh N, Radi FZ, Fakchich J, Elbouzidi A, Ouahhoud S, Ouasti M, Bouhrim M, Ouasti I, Hano C, Elachouri M: Ethnobotanical, Phytochemical, Toxicological, and Pharmacological Properties of *Ziziphus lotus* (L.) Lam.: A Comprehensive Review. *Pharmaceuticals* 2023, 16(4):575.
 7. Bencheikh N, Elbouzidi A, Kharchoufa L, Ouassou H, Alami Merrouni I, Mechchate H, Es-Safi I, Hano C, Addi M, Bouhrim M: Inventory of medicinal plants used traditionally to manage kidney diseases in north-eastern Morocco: Ethnobotanical fieldwork and pharmacological evidence. *Plants* 2021, 10(9):1966.
 8. Ribeiro A, Serrano R, da Silva IBM, Gomes ET, Pinto JF, Silva O: The Genus *Diospyros*: A Review of Novel Insights into the Biological Activity and Species of Mozambican Flora. *Plants* 2023, 12(15):2833.
 9. Abba A, Agunu A, Abubakar A, Abubakar U, Jajere M: Phytochemical screening and antiproliferative effects of methanol extract of stem bark of *Diospyros mespiliformis* Hochst (Ebenaceae) against guinea corn (*Sorghum bicolor*) seeds radicles length. *Bayero Journal of Pure and Applied Sciences* 2016, 9(1):1-5.
 10. Shagal M, Kubmarawa D, Alim H: Preliminary phytochemical investigation and antimicrobial evaluation of roots, stem-bark and leaves extracts of *Diospyros mespiliformis*. *International Research Journal of Biochemistry and Bioinformatics* 2012, 2(1):11-15.
 11. Oguche M, Nzelibe H: In-vivo antiplasmodial activity of aqueous, n-butanol and ethylacetate fractions of leaf and stem bark methanol extracts of *Diospyros mespiliformis* on *Plasmodium berghei berghei* (Nk65) infected mice. *International Journal of Biochemistry Research & Review* 2016, 12(4):1-9.
 12. Belem-Kabré WLME, Nitiéma M, Odjo SB, Noura MO, Yaro B, Kaboré B, Somda DG, Traoré TK, Koala M, Ilboudo Sylvain: Phytochemical Analysis and Contractile Effects of Aqueous and Hydroethanolic Extracts of *Anastatica hierochuntica* L.(Brassicaceae) on the Isolated Uterus of Mice. *Pharmacology & Pharmacy* 2023, 14(8):252-270.
 13. Nitiéma M, Soleti R, Koffi C, Belemnaba L, Mallegol P, Ouédraogo N, Kini FB, Ouédraogo S, Guissou IP, Andriantsitohaina R: Ethyl Acetate Fraction of *Lannea microcarpa* Engl. and K. Krause (Anacardiaceae) Trunk Barks Corrects Angiotensin II-Induced Hypertension and Endothelial Dysfunction in Mice. *Oxidative Medicine and Cellular Longevity* 2019, 2019:13.
 14. Nitiéma M, Ilboudo S, Belemnaba L, Ouédraogo GG, Ouédraogo S, Ouédraogo N, Sylvain O, Guissou IP: Acute and sub-acute toxicity studies of aqueous decoction of the trunk barks from *Lannea microcarpa* Engl. and K. Krause (Anacardiaceae) in rodents. *World journal of pharmacy and pharmaceutical sciences* 2018, 7(9):30-42.
 15. OCDE: Toxicité orale aiguë - Méthode par classe de toxicité aiguë. Lignes directrices 423 de l'OCDE pour les essais de produits chimiques. In., vol. ; 2001, 1-14

16. Ouédraogo WRC, Belemnaba L, Nitiéma M, Kaboré B, Compaoré S, Koala M, Semdé R, Ouédraogo S: Evaluation of acute, subacute toxicity and in vivo impact of aqueous decoction of *Flemingia faginea* Guill. & Perr.(Barker) leafy stems on NMRI mice and normotensive Wistar rats. *Journal of Drug Delivery and Therapeutics* 2023, 13(1):1-10.
17. Rauf A, Akram M, Semwal P, Mujawah AA, Muhammad N, Riaz Z, Munir N, Piotrovsky D, Vdovina I, Bouyahya A: Antispasmodic potential of medicinal plants: a comprehensive review. *Oxidative Medicine and Cellular Longevity* 2021, 2021.
18. Wink M: Modes of action of herbal medicines and plant secondary metabolites. *Medicines* 2015, 2(3):251-286.
19. Ebbo A, Mammam M, Suleiman M, Ahmed A, Bello A: Preliminary phytochemical screening of *Diospyros mespiliformis*. *Research Journal of Agriculture and Environmental Management* 2015, 4:23-26.
20. Ndhlala AR, Chitindingu K, Mupure C, Murenje T, Ndhlala F, Benhura MA, Muchuweti M: Antioxidant properties of methanolic extracts from *Diospyros mespiliformis* (jackal berry), *Flacourtia indica* (Batoka plum), *Uapaca kirkiana* (wild loquat) and *Ziziphus mauritiana* (yellow berry) fruits. *International journal of food science & technology* 2008, 43(2):284-288.
21. Nafuka SN, Mumbengegwi DR: Phytochemical analysis and in vitro anti-plasmodial activity of selected ethnomedicinal plants used to treat malaria associated symptoms in Northern Namibia. *International Science and Technology Journal of Namibia* 2014:078-093.
22. Joseph N, Esther NLT, Benoît NT, Désiré DDP, Bibi-Farouck AO, Théophile D, Pierre K: Effects of the aqueous extract of *Pittosporum mannii* Hook. f.(Pittosporaceae) stem barks on spontaneous and spasmogen-induced contractile activity of isolated rat duodenum. *Journal of ethnopharmacology* 2015, 172:1-9.
23. Shahidi F, Ambigaipalan P: Phenolics and polyphenolics in foods, beverages and spices: Antioxidant activity and health effects—A review. *Journal of functional foods* 2015, 18:820-897.
24. Singla RK, Dubey AK, Garg A, Sharma RK, Fiorino M, Ameen SM, Haddad MA, Al-Hiary M: Natural polyphenols: Chemical classification, definition of classes, subcategories, and structures. In., vol. 102: Oxford University Press; 2019: 1397-1400.
25. Ebbo A, Sani D, Suleiman M, Ahmad A, Hassan A: Assessment of antioxidant and wound healing activity of the crude methanolic extract of *Diospyros mespiliformis* Hochst ex a. Dc (Ebenaceae) and its fractions in Wistar rats. *South African Journal of Botany* 2022, 150:305-312.
26. Bai M, Yao G-D, Ren Q, Li Q, Liu Q-B, Zhang Y, Wang X-B, Huang X-X, Song S-J: Triterpenoid saponins and flavonoids from licorice residues with anti-inflammatory activity. *Industrial Crops and Products* 2018, 125:50-58.
27. Tanahashi Y, Komori S, Matsuyama H, Kitazawa T, Unno T: Functions of muscarinic receptor subtypes in gastrointestinal smooth muscle: A review of studies with receptor-knockout mice. *International Journal of Molecular Sciences* 2021, 22(2):926.
28. Ward SM: Postjunctional M2 muscarinic receptors augment neurally mediated cholinergic contractions of Murine airway smooth muscle. *Function* 2022, 3(1):zqab067.
29. Russo M, Coppola V, Giannetti E, Buonavolontà R, Piscitelli A, Staiano A: Oral administration of tannins and flavonoids in children with acute diarrhea: a pilot, randomized, control-case study. *Italian journal of pediatrics* 2018, 44(1):1-6.

30. Abdoul GLB, Traore A, Ouedraogo M, Belemlilga M, Traore TK, Belemnaba L, Ouedraogo N, Lupu A, Ouedraogo S, Guissou IP: Pharmacological study of trunk bark of *Acacia nilotica* var *adansonii* (Guill et Perr). o Ktze (Mimosaceae): Assays, antioxidant and antispasmodic activities. *Journal of Drug Delivery and Therapeutics* 2019, 9(3-s):524-530.
31. Kuo IY, Ehrlich BE: Signaling in muscle contraction. *Cold Spring Harbor perspectives in biology* 2015, 7(2):a006023.
32. Ebbo A, Sani D, Suleiman M, Ahmad A, Hassan A: Acute and sub-chronic toxicity evaluation of the crude methanolic extract of *Diospyros mespiliformis* hochst ex a. Dc (ebenaceae) and its fractions. *Toxicology reports* 2020, 7:1138-1144.

UNDER PEER REVIEW