

Original Research Article
**Effect of different organic sources of nitrogen
on growth, yield, quality, and economics of
vegetable cluster bean (*Cyamopsis
tetragonoloba* L.)**

Comment [T1]: The title is not smart because there are many types of source of nitrogen

ABSTRACT

An experiment was conducted on vegetable cluster bean variety Pusa Navbharat to find out the effect of different organic sources of nitrogen on growth, yield and economics of vegetable cluster bean. This experiment was conducted in certified organic plot for the three years (2020, 2021, 2022) during semi *rabi* season at Agricultural and Horticultural Research Station, AAU, Khambholaj. The experiment was laid out in randomized block design with three replications. There were eight treatments viz., T₁ : No manures (Control) , T₂: N equivalent of 20 kg/ha through FYM, T₃: N equivalent of 20 kg/ha through Vermi compost, T₄: N equivalent of 20 kg/ha through Castor cake, T₅: Bio NP (*Rhizobium* and PSB), T₆: N equivalent of 10 kg/ha through FYM + Bio NP (*Rhizobium* and PSB), T₇: N equivalent of 10 kg/ha through Vermi compost + Bio NP (*Rhizobium* and PSB) and T₈: N equivalent of 10 kg/ha through Castor cake + Bio NP (*Rhizobium* and PSB). Plant height, green pod yield, its related traits and crude protein content in pods were recorded every year and pooled analysis was carried out. There was significant difference observed for green pod yield and its related traits. Whereas, plant height and crude protein content in pods was remain non-significant for different organic sources of nitrogen. Application of N equivalent of 20kg/ha through FYM recorded maximum yield and its related traits with higher net return.

Keywords: Cluster bean, Organic , Nitrogen, FYM, Vermicompost, Castor cake

1. INTRODUCTION

Cluster bean (*Cyamopsis tetragonoloba* L.) is an important legume vegetable crops popularly known as a guar in India. It is short duration crop mainly cultivated in arid and semi-arid regions of tropical India. In India, it is mainly cultivated in Rajasthan, Gujarat, Punjab, Haryana, Uttar Pradesh, and Maharashtra. In Gujarat during 2022-23 it was cultivated in 45,568 ha with the production of 452,527 MT and productivity of 9.93 t/ha. [1]. In Gujarat Kutch, Banaskantha, Mehsana, Sabarkantha, Vadodara and Ahmedabad are the major cluster bean producing districts.

Comment [T2]: Needs reference

Cluster bean is erect growing plant having deep root system with angled stem, leaves trifoliolate, ovate and serrate while pods borne on auxiliary raceme. Green and tender pods of cluster bean are used as vegetables. It is also grown as a forage crop and being a legume crop it has capacity to fix atmospheric nitrogen through symbiosis and improve the fertility of soil. It may also act as cover crop, which reduce soil erosion and weed problem in mono and intercropping system. Due to more remunerative market prices and low input cost, the farmers of Gujarat have shown more interest in this crop. Organic production of cluster bean contributes the enhancement of production with holistic sense, which includes ecological, economic, and social aspects. Use of organic sources of nutrients in combination with biofertilizers found effective components in organic farming for reliable and cheap supply of nutrients. These combinations are eco-friendly, safe and improve soil fertility with improving physical, chemical, and biological conditions of soil. The beneficial effects of FYM/vermicompost /poultry manure addition are also related to improvement in soil physical properties [2]. There is no systematic information on manure requirement for cluster bean in organic farming hence, the experiment was planned with objective to study the effect of organic sources of nitrogen on the growth, yield, and quality of cluster bean.

Comment [T3]: Not plant, rather plant

Comment [T4]: If so what are the importance of applying nitrogen to the cluster bean?

Comment [T5]: All needs references

Comment [T6]: The statements of the problem was not well defined as a reason.

Comment [T7]: What about how the data were collected
2. about the soil type, method of application
.methods of data record methods and soon

2. MATERIAL AND METHODS

An experiment was conducted to evaluate the performance of vegetable cluster bean for different agronomic sources of nitrogen alone and with biofertilizers in certified organic plot at Agricultural and Horticultural Research Station, AAU, Khambholaj. This experiment was carried out for three years during the semi rabi 2019 to 2022. Randomized block design (RBD) with three replications was used for the experiment. Seeds were sown with a spacing of 30 cm x 15 cm. Recommended doses of 20 kg nitrogen in different treatments was applied through FYM/vermicompost/castor cake as alone and in combination with Bio NP (Rhizobium and PSB). Normal crop management practices were followed for the successful raising of crop. Observation viz., plant height at last harvest (cm), number of clusters per plant, pods per cluster, pods per plant, pod length (cm), green pod yield and crude protein content (%) were recorded for three years and the data were statistically analysis by adopting the standard procedures described by Panse and Sukhatme [3].

Comment [T8]: In the introduction part it said from 2020-2022, but in this section it says from 2019-2022, which one is correct?

Comment [T9]: Who recommend it? If the recommended was known, it must be used as a control for other treatments.

Comment [T10]: 1. At what stage the data were recorded?
2. Why the harvesting index and maturity index were not recorded ?
3. 1000 seed weight, growth parameters and yield parameters in detail
4. where are the quality parameters of the crop ?

3. RESULTS AND DISCUSSION

Inquisition of the data in Table 1 showed that plant height at last harvest and crude protein content of pods was not influenced significantly by different organic sources of nitrogen alone and with Bio NP. Effects of different organic sources of nitrogen alone and with Bio NP (Rhizobium and PSB) were found significant for yield attributes like no. of clusters per plant, no. of pods/clusters, no. of pods/plant and pod length. Significantly higher number of clusters per plant (16.16) recorded with treatment T₂ (N equivalent of 20 kg/ha through FYM) which remain at par with treatment T₃, T₄, and T₈ while the lowest number of clusters per

Comment [T11]: Discussion must separate growth parameters and yield parameters very well separately

Comment [T12]: It is not the right form of writing.

plant (11.75) was recorded in T₁. These might be due to application of FYM/vermicompost/Castor cake alone and in combination with Bio NP increased the availability of nutrients to the plant from different organic sources and biofertilizers may fix atmospheric nitrogen, convert nutrients from insoluble to soluble form, scavenge phosphorus from soil, help in mineralization process and improve yield [4]. Maximum number of pods/cluster (10.49) recorded with treatment T₂ (N equivalent of 20 kg/ha through FYM), which remain at par with treatment T₃, T₄, T₇ and T₈. Whereas minimum pods per cluster noted in treatment T₁ (8.88). This might be due to the application of organic manures alone and with bio fertilizer made nutrients easily available to plants which influenced the role of photosynthesis, protein synthesis and absorbance capacity of nutrients from root. were helpful to convert flowers to pods.

Table 1. Effect of different organic sources of nitrogen on growth parameters, green pod yield, crude protein content of vegetable cluster bean

Treatments	Plant height at harvest (cm)	No of clusters/plant	No. of pods/cluster	No of pods/plant	Pod Length (cm)	Green pod yield (q/ha)	Crude protein content (%)
T ₁ No manures (Control)	111.36	11.75	8.88	121.76	10.74	122	2.88
T ₂ N equivalent of 20 kg/ha through FYM	110.50	16.16	10.49	182.39	15.06	169	2.91
T ₃ N equivalent of 20 kg/ha through Vermi compost	116.78	14.99	9.99	167.76	13.91	150	3.12
T ₄ N equivalent of 20 kg/ha through Castor cake	112.81	16.13	9.98	174.75	14.27	157	2.93
T ₅ Bio NP (Rhizobium and PSB)	111.00	13.79	9.15	142.42	11.81	142	2.99
T ₆ N equivalent of 10 kg/ha through FYM + Bio NP (Rhizobium and PSB)	109.53	14.27	9.42	154.08	13.68	144	2.99
T ₇ N equivalent of 10 kg/ha through Vermi compost + Bio NP (Rhizobium and PSB)	123.11	13.81	9.97	160.92	13.30	153	2.70
T ₈ N equivalent of 10 kg/ha through Castor cake + Bio NP (Rhizobium and PSB)	113.22	15.18	9.96	168.17	13.89	154	2.82
S.Em. ±	3.71	0.56	0.21	5.42	0.33	4.44	0.09
CD at 5%	NS	1.70	0.61	16.43	0.93	13	NS
CV %	10.34	8.63	7.65	8.35	8.55	10.35	6.52

Maximum number of pods per plant (182.39) was recorded in T₁ but it was at par with T₃, T₄ and T₈. These might be due to increasing the availability of nutrients in the root zone which is essentially for the formation and development of the pods ultimately more number of clusters and pods per plant. These results are in accordance with the findings of [5], [6], [7], and [8] reported that use of FYM, castor cake and Rhizobium + PSB recorded higher number of pods per plant. Maximum pod length was observed in T₂ (15.06 cm) which was at par with T₄. This might be due to organic sources viz., FYM and castor cake release nutrients slowly which were helpful to increase pod length. The highest green pod yield (169 q/ha) was recorded in treatment T₂ (20kg/ N from FYM) but it was at par with T₄. This might be due to cumulative effect of all the yield contributing character viz., number of clusters per plant, pods per cluster, pod length and number of pods per plant. It is fact that farm yard manure enhance the adsorptive capacity of soil for cation and anion and these adsorbed

Comment [T13]: It must be discussed very well. How organic manures alone and with bio fertilizer influenced the role of photosynthesis, protein synthesis and absorbance capacity of nutrients from root?

Comment [T14]: Not T1 but T2

Comment [T15]: Is not the reason of itself

Comment [T16]: 1.Release nutrients slowly may increase yield for the next season crop production. Quick release of nutrients may good for the curret crops. Therofre why release nutrients slowly which helpful to increase pod length? It must be discused again
2.release nutrients slowly is the property of all organic fertilizer. Therefore why FYM and castor cake release nutrients slowly which were helpful to increase pod length, can be reason?

ions are released slowly for the entire crop growth period resulted an increased nutrient availability at crop growth period ultimately an increase yield of plant. Organic matter also functions as source of energy for soil micro flora which brings about the transformation of inorganic nutrients held in soil in a form that is readily utilized by growing plants. The beneficial effects of FYM addition are also related to improvement in soil physical properties which in turn helps in better nutrient absorption by plants. The results are partially in agreement with the findings of [9], [10], [11], [12]. These results are in partial agreement with the findings of [13] in groundnut,

The economics in terms of net return and benefit-cost ratio of crop cultivation technology will have a greater impact on practical utility and acceptance by farmers. The economics of different organic sources of nitrogen application in vegetable cluster bean had been computed and exhibited in Table 2. Data on economics revealed that treatment T₂ (N equivalent of 20 kg/ha through FYM) registered maximum net realization of ₹ 427153/ha with higher value of BCR (6.40). Lower net return (Rs. 169775/ha) and benefit cost ratio (3.27) were obtained with control treatment [(Detailed instruction about this section is given below.

Table 2. Effect of different organic sources of nitrogen on cost economics of Cluster bean

Treatments	Green pod yield (kg/ha)	Gross Income (₹/ha)	Comm on Cost (₹/ha)	Treatm ent cost (₹/ha)	Total Cost of Cultiv ation (₹/ha) (4+5)	Net Realiz ation (₹/ha)	BCR
1	2	3	4	5	6	5	6
T ₁ No manures (Control)	12226	244520	68479	0	74745	169775	3.27
T ₂ N equivalent of 20 kg/ha through FYM	16876	337520	68479	2174	78414	259106	4.30
T ₃ N equivalent of 20 kg/ha through Vermi compost	15029	300580	68479	12502	89878	210702	3.34
T ₄ N equivalent of 20 kg/ha through Castor cake	15665	313300	68479	8008	84889	228411	3.69
T ₅ Bio NP (<i>Rhizobium</i> and PSB)	14189	283780	68479	20	76023	207757	3.73
T ₆ N equivalent of 10 kg/ha through FYM + Bio NP (<i>Rhizobium</i> and PSB)	14422	288440	68479	1107	77229	211211	3.73
T ₇ N equivalent of 10 kg/ha through Vermi compost + Bio NP (<i>Rhizobium</i> and PSB)	15254	305080	68479	6271	82961	222119	3.68
T ₈ N equivalent of 10 kg/ha through Castor cake + Bio NP (<i>Rhizobium</i> and PSB)	15449	308980	68479	4024	80467	228513	3.84

Comment [T17]: There must be correlation analysis for each parametrs

4. CONCLUSION

From the results of three years experimentation, it can be concluded that organic cultivation of vegetable cluster bean through different organic sources of nitrogen significantly influenced green pod yield and its traits. Application of required 20 kg N through FYM gave higher green pod yield and net return

Comment [T18]: Not enough rewrite again

REFERENCES

1. Anonymous. Area, production and productivity of horticultural crops in Gujarat 2022-23. <https://doh.gujarat.gov.in/horticulture-census.htm>.
2. Kofoed, AD. The significance of FYM. *Kodemines Tidskrift*. 1987; 19:37-63.
3. Panse, VG and Sukhatme, PV. *Statistical Methods for Agricultural Workers*. Indian Council of Agricultural Research Publication. 1985; 87-89.
4. Roychowdhury, D, Paul, M and Banerjee, S. A. Review on the effects of biofertilizers and biopesticides on rice and tea cultivation and productivity. *International Journal of Science, Engineering and Technology*. 2014; 2(8): 96-106.
5. Singh, B and Kumar, R. Effect of integrated nutrient management on growth, yield and nutrient uptake of cluster bean (*Cyamopsis tetragonoloba* L.) under irrigated conditions. *Agriculture Science Digest*. 2016; 36 (1) : 35-39.
6. Patel, H, Parmar, V, Patel, P and Mavdiya, V. Effect of organic fertilizers on yield and yield attributes of cluster bean (*Cyamopsis tetragonoloba* L.) Cv. Pusa Navbahar. *International Journal of Chemical Studies*. 2018; 6(4): 1797-1799.
7. Sharma P, Meena, RS, Kumar, S, Gurjarm DS, Yadav, GS, Kumar, S. Growth, yield and quality of cluster bean (*Cyamopsis tetragonoloba* L.) as influenced by integrated nutrient management under alley cropping system. *Indian Journal of Agricultural Sciences*. 2019; 89(11):1876-1880.
8. Solanki, KR, Patel, DK, Desai, NA and Vala, YB. Effect of organic sources on growth, yield attributes and yield of summer cluster bean (*Cyamopsis tetragonoloba* L.) under organic farming. *The Pharma Innovation Journal*. 2022; 11(5): 1156-1160.
9. Patel, CS, Patel, JB, Suthar, JV and Patel, PM. Effect of integrated nutrient management on cluster bean [*Cyamopsis tetragonoloba* (L.) Taub] seed production cv. Pusa Navbahar. *International Journal of Agricultural Sciences* 2010; 6(1): 206-208.
10. Patel, HS and Patel KD. Effect of organic fertilizers on growth and economics of cluster bean (*Cyamopsis tetragonoloba* L.) cv. Pusa Navbahar. *International Journal of Chemical Studies*. 2018; 6(5): 861-864.
11. Patel, AS, Kacha, RP, Nayee, PJ and Patel, RM. Nutrient management through organic source in grain amaranthus (*Amaranthus hypochondriacus* L.) under middle Gujarat condition. *International Journal of Chemical Studies*. 2020; 8(4): 1115-1116.
12. Brahmabhatt, JH, Acharya, SK, Patel, GS, Thakar, C. and Solanki, NR. Effect of Organic Nutrient Management on Growth and Yield of Cluster Bean (*Cyamopsis tetragonoloba* L.). *Legume Research*. 2021; DOI: 10.18805/LR-4556.
13. Patel, R, Patel, PM, Patel, HK, Patel, H and Patel T. Nitrogen management through organic sources and biofertilizers in summer groundnut (*Arachis hypogaea* L.). *The Pharma Innovation Journal*. 2022; 11(9): 1342-1347.