

Isolation, Identification and Antibigram of Bacteria ~~isolated from~~ isolated from Raw Cow Milk in Dutsin-Ma, Katsina State

Style Definition: Normal

Style Definition: Heading 3: Font color:
Accent 1

Style Definition: Balloon Text

Style Definition: Hyperlink

Style Definition: Normal (Web)

Style Definition: Default

Abstract

The study was conducted to isolate, identify the bacteria, to know the sources of contamination of milk and antibiotic sensitivity of bacteria obtained from Dutsin-Ma, Katsina State. A total of 45 samples were collected from different locations in Dutsin-Ma such as Wednesday market, opposite FUDMA ~~take off~~ take-off site and Hospital road. All these samples were ~~analyzed~~ analysed by culturing in different media such as *Salmonella-Shigella* agar, Eosin Methylene Blue agar, Mannitol Salt Agar, Nutrient agar, Cetrimide agar, MacConkey agar. Biochemical tests were performed to identify the organism. Among 45 samples, 20 (37.7%) were *Staphylococcus* ~~spp~~ species. Similarly, 11 (20.8%), 6 (11.3%), 4(7.5%) and 12(22.6%) were found positive for *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella* and ~~*Salmonella spp.*~~ *Salmonella species* respectively. Results of antibiotic sensitivity test represent that, out of ten antibiotics *Staphylococcus* ~~sp~~ species were very sensitive against Gentamicin (95%), Ciprofloxacin (90%), Streptomycin (70%), and highly resistant against Zinnacef (60%), Ampiclox (70%), Amoxicillin (50%). *Salmonella* ~~sp~~ species were highly sensitive to Pefloxacin (83.3%), Sparfloxacin (83.3%), Ciprofloxacin (95%), but resistant against Augmentin (83.3%), Streptomycin (75%), Sulfamethoxazole (66.6%). *Klebsiella* ~~spp~~ species were highly sensitive Pefloxacin (50%), Sparfloxacin, Chloramphenicol (75%), Ciprofloxacin (75%), but resistant ~~to Gentamicin~~ to Gentamicin(100%), Streptomycin (100%), Sulfamethoxazole (75%), Augmentin (75%). *Escherichia coli* were highly sensitive to Gentamicin (72.7%), Ciprofloxacin (100%), Ofloxacin (90.9%), Sparfloxacin (72.7%), but highly resistant to Sulphamethoxazole (72.2%), Whereas, *Pseudomonas* ~~aeruginosa~~ aeruginosa were highly sensitive Ciprofloxacin (100%), Chloramphenicol (66.6%), Ofloxacin (66.6%), but highly resistant against Streptomycin (100%), Augmentin 83.35%), Pefloxacin (83.3%). Data of this study suggested that raw milk contaminated with drug resistant bacteria may cause public health hazard.

Key words: Raw, ~~Milk~~ cow milk, Antibiotics, Resistant, Antibiogram, Isolation.

Introduction

Raw milk is obtained from cows at homes in the Fulani hamlets ~~and~~, villages where ~~as~~ shelf-life and ~~safety~~safeties of the products are not considered. Milk is a major component in the human diet of vegetarian class, but it also serves as very good medium for the growth of many Microorganisms including pathogenic bacteria [Ruegg, 2003]. Milk is a highly valuable Food, but raw milk ~~contains and favors~~favours growth of many ~~microorganisms. Foodborne~~micro-organisms. Food borne illnesses are an important challenge to public health and cause significant economic problem in many countries [WHO. 2015]. The crucial goal of all food safety programs is to prevent food products contaminated by potential pathogens from reaching the consumer.

Milk is an excellent medium for bacterial growth, ~~which not only spoils~~whichspoils the milk and associated products but also can cause infections in consumers. Because of the specific production, it is not possible to fully avoid contamination of milk with ~~microorganisms~~micro-organisms; therefore, the microbial contamination of milk is an important tool in determining its quality [Srinu *et al.*, 2012]. During the normal milking operation however, milk is subjected to contamination from many different sources including: (1) the udder and body of cows, (2) dust from the air, (3) litter and floor (4) flies, insects and rodents (5) water supply (6) hands and clothes of the milkier (7) utensils, bottles (8) atmosphere, etc. [Ensminger *et al.*, 1994]. Thus, milk and the dairy products prepared from milk could be an important source of food borne pathogens (Oliver *et al.*, 2019). Milk can be contaminated by various types of ~~microorganisms~~micro-organisms such as *Streptococcus*, *Staphylococcus*, *Micrococcus*,

Formatted: English (United Kingdom)

Escherichia, *Bacillus*, *Salmonella* and *Pseudomonas sp* (Mubarack *et al.*, 2010). Huge numbers of microbes can get access to milk and various milk products including *Escherichia coli*, which is an indicator of milk contamination, constituting a public health hazard [Virpari *et al.*, 2013].

~~The diseases~~ Consumption of spoiled milk cause transmissible ~~diseases~~ to humans ~~through the consumption of spoiled milk~~ like brucellosis, tuberculosis, salmonellosis, listeriosis, *Escherichia coli* infections and many others were described extensively in 1962 by [Kapla *et al.*, 1962].

Antimicrobial development and eventual clinical adoption is one of the most significant issues in medical history, with engineered medicines having saved millions of lives against diseases that would have been lethal [Gajdács, *et al.*, 2012]. Nonetheless, due to the development of ~~multi drug~~ multi drug resistance (MDR) in these pathogens, treating infectious diseases is becoming ~~increasingly more~~ difficult. The present study was undertaken aiming to isolate and identify the bacteria, ~~and to determine the~~ antibiotics resistant of bacteria from raw cow in Dustin-Ma, Katsina State, Nigeria.

Bacterial contamination in raw milk has been a great threat to the economic and human health which ~~cause causes~~ mild disease to life threatening illness. Raw milk can carry dangerous bacteria such as *Salmonella*, *E. coli*, *Listeria*, *Campylobacter*, and others that cause ~~food borne~~ food borne illness, often called “food poisoning.” These bacteria can seriously injure the health of anyone who drinks raw milk or eats products made from raw milk. However, the bacteria in raw milk can be especially dangerous to people with weakened immune systems (such as transplant patients and individuals with HIV/AIDS, cancer, and diabetes), children, older adults, and pregnant women. In fact, CDC finds that ~~food borne~~ food borne illness from raw milk especially affects children and teenagers. Hence, ~~the need for this~~ present study ~~is carried out~~.

Bacterial contamination of raw milk is of grave public health concern, especially when they are Antibiotics Resistant bacteria (ARB). This is because in addition to being a human pathogen, ARB ~~are~~ is of health concern that can be disseminated from contaminated milk via mobile genetic elements like plasmids and ~~transposon~~ transposase. Therefore, ~~resulting to~~ result too complicated, untreatable, and prolonged infections in humans, leading to higher healthcare costs and sometimes death [Ezekiel *et al.*, 2019]. The aim of this research is to isolate, identify and determine antibiotics resistant of bacteria from raw cow in Dustin-Ma, Katsina State, Nigeria.

Material and Methods

Sample Collection

A total of 45 cow milk samples were collected randomly from 3 different locations in Dutsin-ma local government area of Katsina state, Nigeria. Nine samples were taken per week from each location for five (5) weeks. The samples were collected in sterile universal bottle and transported to the Microbiology Laboratory of the Department of Microbiology, Federal University, Dutsin-Ma for microbiological analysis.

Isolation of Bacteria

Serial dilution of 10^{-1} , 10^{-2} , 10^{-3} , 10^{-4} and 10^{-5} were prepared. Aliquots of 1ml from each dilution was inoculated into sterile nutrient agar and MacConkey agar plates using pour plate method and incubated at 37°C for 24 hours. Plate counts were recorded in cfu/ml. Colonial appearance such as size, shape, consistency, ~~eol~~ colour, elevation and differential characteristics such as pigmentation and the isolate was further sub-cultured in eosin methylene blue, ~~SalmonellaShigella~~ SalmonellaShigella and mannitol salt agar and Gram Staining were done to further identify the isolates [Cheesbrough,2003].

Gram Staining Technique

A smear of the suspected colony from overnight culture plates were made on clean, grease – free slide. The smear was heat-fixed on slide by passing the slide briefly over Bunsen burner flame. Smear was then covered with crystal violet stain for 1 minute. Stain was removed, rinsed with tap water. Afterwards, Lugol’s iodine was added for 1 minute while decolonization was carried out by addition of acetone for few seconds. Slide was quickly washed with distilled water and counter stained with Safranin for 1 minute. It was then finally flooded with water and blot dried and examined under the microscope using the oil-immersion objective lens. Suspected *Staphylococcus aureus* isolates were Gram-positive cocci (appearing purples) and arranged in clusters, *Klebsiella* spp., ~~*Escherichia coli*~~, *Escherichia coli*, *Pseudomonas aerogenosa* and *Salmonella* spp. isolate were Gram-negative rod (appearing pink) (Cheesbrough,2003)

Biochemical Characterization and Identification

Catalase Test

A drop of 3% hydrogen peroxide solution was placed on a clean, grease-free glass slide. A loopful of overnight colony of the test organism was thereafter emulsified on the hydrogen peroxide. Observed bubble formation was regarded positive and no bubble formation was regarded negative (Cheesbrough, 2003).

Coagulase Test

One loopful of the colony was emulsified on a clean grease free glass slide. Ten microliter of citrated human plasma was added and observed for the presence of agglutination which indicates a positive reaction (Cheesbrough, 2003).

Urease Test

The media was prepared based on the manufacturer’s instruction ~~then~~, 40% of urea solution was added and mixed well, then poured in a tube and slant, then the isolate was inoculated and

incubated for 18-24hrs. Positive test is demonstrated by an intense magenta to bright pink ~~color~~colour in 15-24hrs, negative test show no ~~color~~colour (Cheesbrough, 2003).

Triple Sugar Iron Test

The media was prepared based on the manufacturers instruction poured in a test tube sterilized at 115⁰C for 30minutes then it was allowed to set in ~~the~~sloped form with a butt inoculation was made by stabbing trough the ~~center~~centre of the medium to the bottom of the tube with a straight inoculation needle then the surface of the agar slant was streaked with the isolated colony. Positive result was indicated by ~~color~~colour change, gas production and production of H₂S (Cheesbrough, 2003).

Indole Test

The media was prepared based on the manufacturers instruction and the isolated colony was inoculated for 37°C for 24-28 hours then Kovac's reagent was added to the broth culture ~~formation of~~Change in colour from pink to red ~~color~~ (“cherry-red ring) in the reagent layer on top of the medium ~~within second of adding the reagent~~indicate positive and no ~~color~~colour change indicated negative (Cheesbrough, 2003)

Methyl Red Test

The broth (MRVP) was prepared according to manufacturer's instruction and the organism was inoculated then incubated for 18-24hrs ~~then add later~~ methyl red reagent~~color added, colour~~ change to red indicate positive and no ~~color~~colour change indicate negative (Cheesbrough, 2003).

Voge's-Proskauer Test

The broth (MRVP) was prepared according to manufacturer's instruction and the organism was inoculated then incubated for 18-24hrs ~~then add~~24hrs later VPI and VPII

reagent ~~color~~added, colour change and ring formation indicate positive lack of ~~color~~colour change indicate negative (Cheesbrough, 2003).

Citrate Utilization Test

The media Simmons agar was prepared based on the manufactures instruction poured in tube slanted then the organism was inoculated and incubate for 18-24hrs at 37°C positive result is indicated by ~~color~~colour change green to blue (Cheesbrough, 2003).

Antimicrobial Susceptibility Testing

Isolates were screened for phenotypic resistance and susceptibility to gram positive and gram negative disc. The procedure includes inoculation of stock cultures stored at 4°C on nutrient agar slants into 10ml of nutrient broth which was then incubated overnight at 37°C. Thereafter, serial dilution of 10¹ into sterile distilled water was carried out. Afterwards, 1 ml of the culture solution was transferred into sterile ~~petri~~Petri dishes. Thereafter, sterile Mueller Hinton agar that has been cooled to 55°C in water bath ~~was poured into each~~ and allowed to solidify. Antibiotic sensitivity disc was later placed on one of the solidified plates sterilely and both plates were incubated at 37°C in an incubator overnight. Zone of inhibition seen round the antibiotic disc the following day were measured while length was categorized as resistant, intermediate and sensitive after comparing with CLSI standard for each bacteria isolate (CLSI, 2021). Multidrug resistant isolates were selected based on their resistant to ≥ 3 classes of antibiotics.

Results

Formatted: Font: Bold

Formatted: Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Table 1: Demonstrated the mean bacterial counts of raw cow [milk](#) collected from the sample locations, in which Wednesday market of Dutsin-Ma is the highest with (8.5×10⁶ cfu/ml), while the least mean count was shown by Hospital Road with (1.04×10⁶ cfu/ml).

Table 2: [Shows](#) the morphological characteristics of all the bacterial isolates of [raw cow milk](#) using both the all-purpose and selective media used for the study.

Table 3: Presents the distributions of [all the gram staining and biochemical test result of all](#) bacterial species isolated from the study sites, which include Staphylococcus aureus, Salmonella species, Escherichia coli, Pseudomonas aeruginosa, and Klebsiella species.

Formatted: Line spacing: 1.5 lines

Formatted: Font: Bold

Table 4: [Shows](#) the prevalence of bacteria isolated from the study sites, in which Staphylococcus was found to be the most prevalent (37.8%), while Klebsiella spp. demonstrated the least prevalent (7.5%).

Table 5: [Shows](#) the percentage of occurrence of bacterial species isolated from the three sample locations. Staphylococcus aureus showed the highest percentage of occurrence (37.2%) while Klebsiella spp. showed the lowest prevalence (7.5%).

Table 6: Demonstrated the microscopic and biochemical characteristics of five bacterial species associated with the [raw cow milk](#).

Table 7: [Shows](#) the antibiotic susceptibility profiles of gram-positive bacteria [isolated from raw cow milk](#) on different antibiotics.

Table 8: [Shows](#) the antibiotic susceptibility profiles of gram-negative bacteria isolated from [raw cow milk](#) on different antibiotics

Table 1: Mean ~~Value~~ bacterial counts of ~~Bacteria Load~~ raw cow milk at 3 different location

S/No	Locations	Mean bacterial count	Percentage %
1.	Wednesday market	8.5 X 10 ⁶	48.5
2.	Front of school FUDMA take-off site	7.99 X 10 ⁶	45.6
3.	Hospital road	1.04 X 10 ⁶	5.9
Total		1.753 X 10⁷	100

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Normal

Table 2: Morphological Characteristics of Bacteria Isolated from Cow Milk in Dutsin-Ma

Media	Morphology	Bacteria
SSA	Pink with black dot	<i>Salmonella</i>
EMB	Green with metallic sheen	<i>Escherichia coli</i>
	Pink no sheen mucoid	<i>Klebseilla spp</i>
MSA	Round transparent	<i>Staphylococcus aureus</i>
Cetrimide	Greenish-blue	<i>Pseudomonas aeruginosa</i>

Key; SSA – ~~Salmonellashigella~~ *Salmonellashigella* Agar, EMB – Eosin methylene blue, MSA – Mannitol salt agar

Formatted: Font: Not Bold

Formatted: Line spacing: single

Table 3: Gram Staining and Biochemical Test Result

Bacteria	Gram Stain	Catalase	Oxidase	Urease	Citrate	Tsi	Indole	Mr	Vp	Coagulase
<i>Escherichia Coli</i>	-				-	+	+	+	-	
<i>S. Aureus</i>	+	+		+			-	+	+	+
<i>Salmonella Spp.</i>	-			-	-	+	-	+	-	
<i>Klebsiella Spp.</i>	-	+			+		-	-	+	+

|


<i>P. aeruginosa</i>	-	+	+	+	-	-	-
----------------------	---	---	---	---	---	---	---

- = negative + =positive, VP = Voge's Proskauer, MR = Methyl Red, TSI = Triple sugar ion test


|

Table 4. Distribution of Isolates of Raw Cow Milk from Different Locations

S/N	Isolate	Locations			Total
		Wednesday Market	Front of School FUDMA take-off site	Hospital Road	
1.	<i>Staphylococcus aureus</i>	7	8	5	20
2.	<i>Escherichia coli</i>	4	3	4	11
3.	<i>Pseudomonas aeruginosa</i>	2	1	3	6
4.	<i>Klebsiella Spp.</i>	2	-	2	4
5.	<i>Salmonella Spp.</i>	5	4	3	12
	Total	20	16	17	53



Formatted: Font: (Default) Times New Roman




Formatted: Font: (Default) Times New Roman

Figure 1: Distribution of Isolates of Raw Cow Milk from Different Locations


Table 5: Percentage of occurrence of Bacteria Species

S/N	Isolate	Wednesday Market	Hospital Road	Front of Fudma Take FUDMA take-offsite	Percentage (%)
1.	<i>Staphylococcus Aureus</i>	7	5	8	37.7
2.	<i>Escherichia Coli</i>	4	4	3	20.8
3.	<i>Pseudomonas</i>	2	3	1	11.3
4.	<i>Klebsiella Spp.</i>	2	2	-	7.5
5.	<i>Salmonella Spp.</i>	5	3	4	22.6
	Total	20	17	16	100

Formatted: Font color: Black, English (United Kingdom)



Formatted: Font: (Default) Times New Roman




Formatted: Font: (Default) Times New Roman

Figure 2: Percentage of Occurrence of Bacteria Species

Table 6: Antibiotics Susceptibility Pattern of Gram Positive Bacteria


Antibiotics profile (%)	Isolates		
	Staphylococcus aureus (n=20)		
	Susceptible	Intermediate	Resistance
Zinnacef	4(20)	4(20)	12(60)
Rocephin	10(20)	4(20)	5(25)
Streptomycin	14(70)	0(00)	4(20)
Erythromycin	12(60)	0(00)	8(40)
Gentamicin	19(95)	0(00)	1(5)
Ampiclox	4(20)	2(10)	14(70)
Sulfamethoxazole	12(60)	0(00)	8(40)
Amoxicillin	8(40)	2(10)	10(50)
Pefloxacin	10(50)	6(30)	4(20)
Ciprofloxacin	18(90)	2(10)	0(00)

Fig 3. Antibiotics resistance profile of *Staphylococcus aureus*



Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Fig 3. Antibiotics resistance profile of *Staphylococcus aureus*




Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Table 7: Antibiotics Susceptibility Pattern of Gram-Negative Bacteria from Raw Cow Milk

Antibiotics profile (%)	Isolates											
	<i>Salmonella</i> (n=12)			<i>Klebsiella</i> (n=4)			<i>Escherichia coli</i> (n=11)			<i>Pseudomonas aeruginosa</i> (n=6)		
	S	I	R	S	I	R	S	I	R	S	I	R
AM	4(33.3)	3(25)	5(41.6)	1(25)	1(25)	2(50)	4(36.3)	2(18.2)	5(45.5)	2(33.3)	0	4(66.6)
PEF	10(83.3)	0	2(16.6)	2(50)	1(25)	1(25)	4(36.3)	5(45.5)	3(27.3)	1(16.6)	0	5(83.3)
SP	10(83.3)	1(8.3)	1(8.3)	1(25)	3(75)	0	8(72.7)	2(18.2)	1(9.09)	2(33.3)	3(50)	1(16.6)
CH	7(58.3)	3(25)	2(16.6)	3(75)	1(25)	0	6(54.5)	0	5(45.5)	4(66.6)	0	3(33.3)
GEN	6(50)	2(16.6)	4(33.3)	0	0	4(100)	8(72.7)	1(9.09)	3(27.3)	2(33.3)	1(16.6)	3(33.3)
AU	2(16.6)	0	10(83.3)	1(25)	0	3(75)	4(36.3)	2(18.2)	5(45.5)	1(16.6)	0	5(83.3)
OFX	7(58.3)	2(16.6)	3(25)	1(25)	3(75)	0	10(90.9)	1(9.09)	1(9.09)	4(66.6)	2(33.3)	0
S	2(16.6)	1(8.3)	9(75)	0	0	4(100)	2(18.2)	4(36.3)	5(45.5)	0	0	6(100)
SXT	1(8.3)	3(25)	8(66.6)	0	1(25)	3(75)	2(18.2)	1(9.09)	8(72.7)	1(16.6)	2(33.3)	3(33.3)
CPX	11(95)	0	0	3(75)	1(25)	0	11(100)	0	0	6(100)	0	0


KEY S=Sensitive, I= Intermediate, R=Resistant, AM=Amoxicillin, PEF=Pefloxacin, SP=Sparfloxacin, CH=Chloramphenicol, CN=Gentamicin, AU=Augmentin, OFX=Ofloxacin, S=Streptomycin, SXT= Sulfamethoxazole, CPX=Ciprofloxacin

Fig 4: Antibiotics resistance profile of *Salmonella*




Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Fig 5: Antibiotics Resistance profile of *Klebsiella spp.*




Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Fig 6: Antibiotics resistance profile of *Escherichia coli*




Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Fig 7: Antibiotics resistance profile of *Pseudomonas aeruginosa*




Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Fig 4: Antibiotics resistance profile of *Salmonella*




Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Fig 5: Antibiotics Resistance profile of *Klebsiella spp.*




Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Fig 6: Antibiotics resistance profile of *Escherichia coli*



Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Bold



Discussion

The highest mean value of bacteria load of raw cow milk sample from Wednesday market as was 8.5×10^6 CFU/ml (48.5%), where the moderate mean values of bacteria load from FUDMA take off campus as was 7.99×10^6 CFU/ml (45.6%) ~~While the and~~ lowest mean value was ~~obtained~~ from Wednesday Market with Hospital road 1.04×10^6 CFU/ml (5.9%).

Morphological characteristics of the bacteria on Eosin Methylene Blue, Salmonella *Shigella*, Cetrimide and mannitol salt agar was done. On Eosin Methylene Blue agar ~~Escherichiacoli~~ Escherichiacoli and *Klebsiella* were identified, ~~Escherichiacoli~~ Escherichiacoli appeared green with metallic sheen, *Klebsiella* pink no sheen mucoid. On Cetrimide *Pseudomonas aeruginosa* appeared green. On *Salmonella-Shigella* agar *Salmonella* was identified which appear colorless with black dot, and on Mannitol salt agar ~~Staphylococcus aureus~~ Staphylococcus aureus was identified as round and transparent.

The distribution of the bacteria in different location, Wednesday market 20 were isolated *Staphylococcus aureus* (7), *Salmonella* spp. (5), *Klebsiella* spp. (2), *Pseudomonas aeruginosa* (2) and *Escherichia coli* (4), in hospital road 17 were isolated with *Staphylococcus aureus* (5), *Salmonella* spp. (3), *Klebsiella* spp. (2), *Pseudomonas aeruginosa* (3) and *Escherichia coli* (4), in front of FUDMA take off campus 16 were isolated with *Staphylococcus aureus* (8), *Salmonella* (4), *Klebsiella* (0), *Pseudomonas aeruginosa* (1) and *Escherichia coli* (3). *Klebsiella* spp. were not present in front of FUDMA take off campus.

~~The percentage of bacteria isolate.~~ This study revealed that *Staphylococcus aureus* has the highest occurrence of bacteria isolate percentage with 37.8% followed by *Salmonella* spp. with (22.6%), *Pseudomonas aeruginosa* (11.3%), *Escherichia coli* (20.8%) and *Klebsiella* spp. (7.5%). This finding does not agree with the findings of (Nwosu *et al.*, 2017) who reported that *E. coli* (86.7%) and *Salmonella* spp. (86.7%), *Staphylococcus aureus* (80.0%), *Klebsiella* spp. (73.3%) and *Pseudomonas aeruginosa* (66.7%). Reta *et al.*, (2016) reported a prevalence rate of (24.2%) for *Staphylococcus aureus* in cow milk consumed at Jigjiga city, Ethiopia and (20.8%) prevalence rate for *Escherichia coli* reported by Makat *et al.*, (2014) isolated from locally processed Cow milk products in Nassarawa state.

The antibiotics resistant to gram positive *Staphylococcus aureus*, Gentamicin, and Ciprofloxacin were sensitive with 95% and 90% respectively this does not correlate with the study of (Rokeya *et al.*, 2019) which show sensitivity to ciprofloxacin 64% and gentamicin 93%. The organism was also found to be resistant to Ampiclox and Zinnacef with 70% and 60% respectively.

The antibiogram of gram negative bacteria all the isolates were highly sensitive to ciprofloxacin *salmonella* was found to be also sensitive to Perfloxacin and Sparfloxacin with 83.3% respectively, also resistance to Augmentin (83%) and streptomycin (83.3%). In a study by Makat *et al.*, (2014), *Salmonella* spp. was also sensitive to both Perfloxacin (43%) and Sparfloxacin (85.7%) and also resistant to Augmentin (14.4%) and Streptomycin (14.2%) (Makat *et al.*, 2014). *Klebsiella* was sensitive to chloramphenicol (75%) and resistance to gentamicin and streptomycin with 100%. *Escherichia coli* was sensitive to Ofloxacin (90.9%) (but resistance to Sulfamethoxazole (72.7%). *Pseudomonas aeruginosa* was resistance to streptomycin (100%) and Augmentin (83.3%). This study does not correlate with the study of (Nwosu *et al.*, 2017)

Conclusion

The highest mean value of bacteria load of raw cow milk sample from Wednesday market as 8.5×10^6 CFU/ml (48.5%), where the moderate mean values of bacteria load from FUDMA take off campus as 7.99×10^6 CFU/ml (45.6%) While the lowest mean value was obtained from Wednesday Market with 1.04×10^6 CFU/ml (5.9%).

The distribution of the bacteria in different location, Wednesday market 20 were isolated ~~*Staphylococcus aureus*~~ *Staphylococcus aureus* (7), *Salmonella* spp. (5), *Klebsiella* spp. (2), ~~*Pseudomonas aeruginosa*~~ *Pseudomonas aeruginosa* (2) and ~~*Escherichia coli*~~ *Escherichia coli* (4), in hospital road 17 were isolated with staphylococcus aureus (5), *Salmonella* spp. (3), *Klebsiella* spp. (2), ~~*pseudomonas aeruginosa*~~ *pseudomonas aeruginosa* (3) and ~~*Escherichia coli*~~ *Escherichia coli* (4), in front of FUDMA take off campus 16 were isolated with ~~*Staphylococcus aureus*~~ *Staphylococcus aureus* (8), *Salmonella* (4), *Klebsiella* (0), *Pseudomonas aeruginosa* (1) and *Escherichia coli* (3). *Klebsiella* spp. were not present in front of FUDMA take off campus.

The percentage of bacteria isolate. This study revealed that *Staphylococcus aureus* has the highest occurrence with 37.8% followed by *Salmonella* spp. with (22.6%), *Pseudomonas aerogenosa* (11.3%), *Escherichia coli* (20.8%) and *Klebsiella* spp. (7.5%).

The antibiotics resistant to gram positive *staphylococcus aureus*, Gentamicin, and Ciprofloxacin were sensitive with 95% and 90% respectively. The antibiogram of gram negative bacteria all the isolates were highly sensitive to ciprofloxacin, *salmonella* was found to be also sensitive to Perfloxacin and Sparfloxacin with 83.3% respectively, also resistance to Augmentin (83%) and streptomycin(83.3%).

The results obtained shows that there is presence of pathogenic microorganism that may be potential source of food borne infection that may result to food borne diseases in the consumers of these sampled products. The total viable bacteria count in all samples were above the standard, according to Nigerian Agency for Food, Drugs Administration and Control (NAFDAC (2009), the microbial load limited for total liable colony count is 1.0×10^2 cfu/ml.

This study also showed high levels of resistance to commonly prescribed antibiotics augmentin, streptomycin, gentamicin and sulfamethoxazole in the bacterial isolates. This calls for strengthening of regulations that cover the sale, distribution, dispensing, and prescription of veterinary antibiotics. This is because antibiotic resistant bacteria may cause complicated, untreatable, and prolonged infections in humans, leading to higher healthcare cost and sometimes death.

Recommendation

It is recommended that good sanitary measures should be taken by the people handling the cows, these measures should include proper handling of the cow, personal hygiene, treatment of udder

infection of the cow, use of hygienic milking and processing equipment, improve milk handling environment. It must be ensured that the cows are always in good health condition.

References

Abdulkadir, M., & Mugadi, A. G. (2012). Bacteriological examination of fura da nono (fermented milk; cereals mix) sold in some selected areas of Birnin Kebbi Metropolis. *ARPN Journal of Science and Technology*, 2, 333-340.

Adesokan, I. A., Odetoynbo, B. B., & Olubamiwa, A. O. (2008). Biopreservative activity of lactic acid bacteria on suya produced from poultry meat. *African Journal of biotechnology*, 7(20).

AJ, B. (1990). Microbiology of raw milk. *Dairy microbiology*, 1, 163-208.

Bhaskar, S. V. (2017). Foodborne diseases—disease burden. In *Food safety in the 21st century* (pp. 1-10). Academic Press.

Bramley, A. J. (1982). Sources of *Streptococcus uberis* in the dairy herd: I. Isolation from bovine faces and from straw bedding of cattle. *Journal of Dairy Research*, 49(3), 369-373.

Bramley, A. J., McKinnon, C. H., Staker, R. T., & Simpkin, D. L. (1984). The effect of udder infection on the bacterial flora of the bulk milk of ten dairy herds. *Journal of Applied Bacteriology*, 57(2), 317-323.

Brunelle, B. W., Bearson, B. L., & Bearson, S. M. (2015). Chloramphenicol and tetracycline decrease motility and increase invasion and attachment gene expression in specific isolates of

|
multidrug-resistant *Salmonella enterica* serovar Typhimurium. *Frontiers in Microbiology*, 5, 801.

Butler, M. T., Wang, Q., & Harshey, R. M. (2010). Cell density and mobility protect swarming bacteria against antibiotics. *Proceedings of the National Academy of Sciences*, 107(8), 3776-3781.

Dewi, G. (2018). *Investigating the Potential of Lemongrass Essential Oil Against Multidrug-Resistant Salmonella Heidelberg in Broiler Chickens* (Doctoral dissertation, University of Minnesota).

Ensminger ME., (1994). “*Foods and nutrition encyclopedia*” 2.1: 2415.

Falegan, C. R., & Akere, G. A. (2014). Isolation of salmonella spp in ‘wara’(local cheese) from three different locations in ado-ekiti, ekiti state, Nigeria. *The experiment*, 23(4), 1628-1634.

Fenlon, D. R., Logue, D. N., Gunn, J., & Wilson, J. (1995). A study of mastitis bacteria and herdmanagement practices to identify their relationship to high somatic cell counts in bulk tank milk. *British Veterinary Journal*, 151(1), 17-25.

Gajdács, M., & Albericio, F. (2019). Antibiotic resistance: from the bench to patients. *Antibiotics*, 8(3), 129.

Galton, D. M., Petersson, L. G., Merrill, W. G., Bandler, D. K., & Shuster, D. E. (1984). Effects of premilking udder preparation on bacterial population, sediment, and iodine residue in milk. *Journal of dairy science*, 67(11), 2580-2589.

Gebreyes, W. A., Thakur, S., Davies, P. R., Funk, J. A., & Altier, C. (2004). Trends in antimicrobial resistance, phage types and integrons among *Salmonella* serotypes from pigs, 1997–2000. *Journal of Antimicrobial Chemotherapy*, 53(6), 997-1003.

|

Gehring, G. (1980). Multiplication of bacteria during farm storage, in factors influencing the bacteriological quality of raw milk. *IDF Bulletin*, 120.

Gonzalez, R. N., Jasper, D. E., Bushnell, R. B., & Farver, T. B. (1986). Relationship between mastitis pathogen numbers in bulk tank milk and bovine udder infections in California dairy herds. *Journal of the American Veterinary Medical Association*, 189(4), 442-445.

Harshey, R. M. (2003). Bacterial motility on a surface: many ways to a common goal. *Annual Reviews in Microbiology*, 57(1), 249-273.

Helms, M., Vastrup, P., Gerner-Smidt, P., & Mølbak, K. (2002). Excess mortality associated with antimicrobial drug-resistant Salmonella Typhimurium. *Emerging infectious diseases*, 8(5), 490.

Hogan, J. S., Smith, K. L., Hoblet, K. H., Todhunter, D. A., Schoenberger, P. S., Hueston, W. D., ... & Conrad, H. R. (1989). Bacterial counts in bedding materials used on nine commercial dairies. *Journal of dairy science*, 72(1), 250-258.

Jackson, H., & Clegg, L. F. L. (1966). The microflora of raw bulk tank milk. *Canadian Journal of Microbiology*, 12(3), 429-432.

Jeffrey, D.C. and J. Wilson. (1987). *Effect of mastitis-related bacteria on the total bacteria counts of bulk milk supplies*. J. Soc. Dairy Technol. 40(2):23.

Kearns, D. B. (2010). A field guide to bacterial swarming motility. *Nature Reviews Microbiology*, 8(9), 634-644.

Khatun, M. M. (2020). Isolation, Identification and Antibiotic Sensitivity Test of Bacteria in Raw Cow Milk Obtained from Different Sources. *EC Microbiology*, 16, 32-41.

Kurweil, R., & Busse, M. (1973). Total count and microflora of freshly drawn milk. *Milchwissenschaft*, 28, 427.

Lippolis, J. D., Brunelle, B. W., Reinhardt, T. A., Sacco, R. E., Thacker, T. C., Looft, T. P., & Casey, T. A. (2016). Differential gene expression of three mastitis-causing *Escherichia coli* strains grown under planktonic, swimming, and swarming culture conditions. *Msystems*, 1(4), 10-1128.

Mackenzie, E. (1973). Thermotolerant and psychrotrophic organisms on poorly cleansed milking plants and farm bulk milk tanks. *Journal of Applied Microbiology*, 36(3), 457-463.

McKinnon, C. H., Rowlands, G. J., & Bramley, A. J. (1990). The effect of udder preparation before milking and contamination from the milking plant on bacterial numbers in bulk milk of eight dairy herds. *Journal of Dairy Research*, 57(3), 307-318.

Miriam Jamila, Md Zulfekar Ali, Shaharin Sultan, Md Ariful Islam and Mst Minara Khatun. (2020). "Isolation, Identification and Antibiotic Sensitivity Test of Bacteria in Raw Cow Milk Obtained from Different Sources". *EC Microbiology* 16.4 32-41.

Mubarack, H. M., Doss, A., Dhanabalan, R., & Balachander, S. (2010). Microbial quality of raw milk samples collected from different villages of Coimbatore District, Tamilnadu, South India. *Indian Journal of Science and Technology*, 3(1), 61-63.

Oliver, S. P., Jayarao, B. M., & Almeida, R. A. (2005). Foodborne pathogens in milk and the dairy farm environment: food safety and public health implications. *Foodborne Pathogens & Disease*, 2(2), 115-129.

Olson, J.C. Jr., and G. Mocquat. (1980). *Milk and Milk Products*. p. 470. In *Microbial Ecology of Foods*. Vol. II.

|

Palmer, J. (1981). Contamination of milk from the milking environment. *Kieler Milchwirtschaftliche Forschungsberichte*, 33(4), 307-316.

Pankey, J. W. (1989). Premilking udder hygiene. *Journal of dairy science*, 72(5), 1308-1312.

Partridge, J. D., & Harshey, R. M. (2013). Swarming: flexible roaming plans. *Journal of bacteriology*, 195(5), 909-918.

Rehman, M. U., Rashid, M., Sheikh, J. A., & Bhat, M. A. (2014). Molecular epidemiology and antibiotic resistance pattern of enteropathogenic Escherichia coli isolated from bovines and their handlers in Jammu, India. *Journal of Advanced Veterinary and Animal Research*, 1(4), 177-181.

Richardson, G. H. (1985). Standard methods for the examination of dairy products. (*No Title*).

Smith, K., Peter, K., Daniela, H., & Melchior, S. (2007). Food borne pathogenic microorganisms and natural toxins. *Food drug Administration center food safety, Applied Nutrition*, 10, 119-150.

Srinu, B., Kumar, A. V., Kumar, M. S., Narayana, B. V. L., & Rao, T. M. (2012). Assessment of microbiological quality and associated health risks of raw milk sold in and around Hyderabad city. *International Journal of Pharma and Bio Sciences*, 3(4), 609-614.

Thomas, S. B. (1974). THE MICROFLORA OF BULK COLLECTED MILK. I.

Thomas, S. B., Druce, R. G., & King, K. P. (1966). The microflora of poorly cleansed farm dairy equipment. *Journal of Applied Bacteriology*, 29(2), 409-422.

Tolle, A. (1980). The Microflora of the Udder, P4 In Factors Influencing The Bacteriological Quality of Raw Milk. *International Dairy Federation Bulletin*.

Virpari, P. K., Nayak, J. B., Brahmhatt, M. N., & Thaker, H. C. (2013). Study on isolation, molecular detection of virulence gene and antibiotic sensitivity pattern of Escherichia coli isolated from milk and milk products. *Veterinary World*, 6(8).

Zehner, M. M., Farnsworth, R. J., Appleman, R. D., Larntz, K., & Springer, J. A. (1986). Growth of environmental mastitis pathogens in various bedding materials. *Journal of dairy science*, 69(7), 1932-1941.

REVIEWER COMMENTS

1. Not mentioned reference numbers in the manuscript.
2. Mention full form of CDC, CLSI
3. Not mentioned specific type randomization technique

4. Table 6: Demonstrated the microscopic and biochemical characteristics of five bacterial species associated with the raw cow milk- NOT MADE
5. Not mentioned Figure / Graph number in the manuscript but Graph is prepared and displayed
6. Not mentioned about statistical analysis, test and P value etc
7. Not written in Vancouver style **References**

Formatted: Font: Not Bold, Font color: Custom Color(RGB(34,34,34)), English (United Kingdom), Pattern: Clear (White)

Formatted: Indent: Left: 0.3", Tab stops: 0.3", Left