

Effects of Water Deficit Stress at the Tillering Stage of Direct Seeded Rice (Oryza sativa L.)

Abstract

The ten rice ~~varities~~ varieties viz. NAUR-1, Purna, GNR-6, NVSR-2227, NVSR-3093, NVSR-2285, NVSR-2278, NVSR-2852, NVSR-2601, and NVSR-2927 were grown in *kharif* season for observing their efficiency in terms of yield under a deficit of water at tillering stage. The effects of ~~short term~~ short-term water stress on various morpho-physiological and biochemical traits viz. leaf area index, number of panicles, chlorophyll content, number of stomata, relative water content, abscisic acid, proline, glycine betaine, ascorbic acid, and harvest index content were studied and they were observed higher in Purna as compared to other nine ~~varities~~ varieties. However, Plant height and Ionic leakage were recorded lower in Purna as compared to other ~~varities~~ varieties. The abscisic acid as well as proline and glycine betaine content recorded maximum in Purna which might help in keeping hydration in ~~cell~~ cells during water deficit stress. Thus, ~~Short term~~ with short-term water deficit stress at tillering stage, the variety Purna would be better in terms of breeding for tolerant variety as well as yield among ten rice varieties.

Keywords: Water, Stress, Rice, Tillers, ~~Varities~~ Varieties

Introduction

~~The e~~ Cereals have a major role in feeding ~~the~~ billions of people and 50% to 80% of daily calorie intake is provided by rice, especially in Asian countries. Among cereals, rice consumes almost 80% of the total irrigation freshwater resources, and its roots ~~being~~ are submerged during the vegetative and reproductive ~~stages~~ stages. The aerenchymatous tissue is a specialized structure in rice ~~which~~ that makes its growth and development during the submerged condition (Henry et al., 2016). ~~The direct seeded~~ Direct-seeded rice generally ~~grow~~ grows on upland/low water regime ~~face~~ faces water deficit stress, ~~and~~ changes its structure and ~~develops~~ develop deep and thick roots ~~results~~ resulting in the improvement ~~of~~ of hydraulic properties of roots thereby ~~extract~~ extracting water from deeper soil profiles (K. Ji et al., 2012). These ~~develop~~ developed anatomy under drought stress support higher yield in upland rice than lowland rice. Many developmental processes of ~~plant~~ plants such as the number of tillers, panicle initiation, and anthesis are adversely affected by drought stress (R. Serraj et al., 2011). Drought severity ~~activate~~ activates the production of Reactive Oxygen Species (ROS) mainly in chloroplasts ~~and~~ and mitochondria which reduce the

membrane stability index and promote leakage of ions due to membrane disruption (Pandey and Shukla, 2015). The ROS viz. Singlet oxygen, superoxide anion radicals, hydroxyl radicals, and hydrogen peroxide (H₂O₂) are produced and initiate indirect disruption of metabolism (Kant, 2020). The defense mechanism for ROS scavenging is developed in ~~plant-plants for to~~ detoxify and suppress the production of ROS by synthesis of different enzymatic and non-enzymatic antioxidants. The ~~drought-tolerant~~ drought-tolerant genotype of rice dramatically increases antioxidant enzymes viz. superoxide dismutase, peroxidase, and catalase which suppress the production of ROS in the cell (Mishra and Panda, 2027; Kant, 2023). The adverse effects of ROS reduced by the synthesis of compatible solutes/osmolytes such as proline, glycine betaine, sugars (trehalose), and polyols (sorbitol, mannitol, myo-inositol, and pinitol) which exalted hydration in the cell. These osmolytes retain water ~~molecul~~ molecules in their structure and they become shield shields of the osmoprotectants in ~~plant-plants~~ during severe water/salinity stress (Anjum, S. A. et al., 2017).

Proline is one of the best ~~osmolyte~~ osmolytes which support the vitality of the plant during water deficit. It is a neutral imino acid at pH-7, ~~having has~~ rigid structure, and is highly soluble in water. It stabilizes protein structures and inhibits protein unfolding during water stress (Szabados, L. & Savoure A. 2010). The enzymes of the metabolic pathway are alleviated by proline particularly ribulose 1, 5 bisphosphate carboxylase that ~~initiate~~ initiates the Calvin cycle (Kaur and Asthir, 2015). The oxidative stress is the main factor for membrane disruption and ionic leakage, and this occurs when the synthesis of reactive oxygen species (ROS) exceeds their scavenging in the cell. Proline is considered ~~as~~ an effective hydroxyl radical scavenger and promoter of enzymatic antioxidants and its biosynthesis in chloroplasts reached ~~upto~~ up to 80% during stress as ~~compare~~ compared to 5% in normal which ~~facilitate~~ facilitates the electron flow between photosynthetic excitation centers and maintains a low NADPH:NADP ratio in cell. Thus, it protects the plant from photoinhibition and damage to the photosynthetic apparatus because an extra electron in the cell is the root cause of ROS production (Bandurska et al. 2017). Proline accumulation increases cellular osmolarity ~~that~~ drives the influx of water or reduces its efflux resulting in an increase ~~of in~~ pressure potential for cell expansion (Bhaskara et al., 2015). Another biomolecule ~~which that protect~~ protects plant-plants during water stress is glycine betaine which acts as a compatible solute and improves the tissue water status. It is synthesized from choline and glycine when a plant is exposed to drought and salinity. It improves

Ca²⁺-ATPase and Hill reaction activities in the thylakoid membrane thereby promote promoting photosynthesis and translocation of sucrose. It is a quaternary amine with zwitterionic nature and can protect biological membranes from ROS as well as improving improve antioxidants-antioxidant defense system under stress (Nawaz and Wang, 2020). The biosynthetic genes of proline and glycine betaine glycine betaine have been widely used in transgenic production to improve abiotic stress tolerance in plants (Schafleitner et al. 2007, Swamy and Kumar, 2013). Therefore, the present study has been focused on creating short term short-term water deficit stress at the tillering stage of rice for the determination of better yield among different varities varieties.

Materials and Methods

The experiment was carried out at the greenhouse located at N. M. College of Agriculture, Navsari Agricultural University, Navsari, Gujarat, India. The Navsari Agricultural University, Navsari campus is geographically located at 20° 57' N latitude and 72° 54' E longitude at an altitude of 16 m above the mean sea level. The average temperature and relative humidity recorded 31±1⁰C and 80%, respectively. Theseeds of ten rice varities varieties (NAUR-1, Purna, GNR-6, NVSR-2227, NVSR-3093, NVSR-2285, NVSR-2278, NVSR-2852, NVSR-2601, NVSR-2927) subspecies *Indica* groupsown in the pot filled with soil and FYM in the ratio of 2:1 in each pot after soaking in water for 12 hours and treatment with carbendazim 50% WP by immersing for 10 minutes. Both mixtures of soil and FYM were weighed and it was 7.5 Kg per pot. Fertilizer application to the test crop was done adopting the recommended fertilizer dose (RDF) for the Navsari region *i.e.*, N - P₂O₅ - K₂O (100-30-00 kg ha⁻¹). The water deficit stress created at the tillering stage having has the a field capacity of 60%. The morphophysiological and biochemical traits were studied after the creation of stress. Plants height was measured in centimetre centimetres from ground level to the tallest leaf of the plant at different stages of crop growth and the average mean of height was recorded. The leaf area index (LAI) was calculated by dividing the leaf area per plant by the land area occupied by the plant. The number of panicles per plant was counted and the average was worked out. Stomatal Frequency was measured with a microscope and expressed as the number of stomata per micrometer square. Chlorophyll content was measured with a SPAD meter. Ionic leakage was measured following the method of R. Sivakumar *etal.* (2015). Weighed two sets of 200 mg of leaf samples and put them into two test tubes containing 10 ml double distilled water. Kept one test tube at 40°C in a water bath for 30

minutes. Thereafter ~~measured~~ the electrical conductivity (EC) of water containing the sample using a Conductivity Bridge and noted. The other test tube was Incubated at 100°C in the boiling water bath for 15 minutes and measured the EC (C2). Relative Water Content (RWC) was calculated following the method described by R. K. Sairam (1994). ~~Third~~ The third leaf (physiologically functional) from the top is selected for RWC estimation. Floated the leaf discs in water for four hours to attain full turgid, and after ~~taken~~ taking out the leaf discs from the water, and ~~wipe~~ wiped out the water droplets sticking on the leaf surface by using filter paper. Immediately recorded the turgid weight (Tw) after an hour of floating. Transferred the leaf discs to a butter paper cover and then ~~keeps~~ kept the cover in a hot air oven at 80°C for 48 hours. Recorded the dry weight (Dw) after removing from the Oven. $RWC = \frac{Fw - Dw}{Tw - Dw} \times 100$. Harvest index measured in percentage. Harvest index (%) = Economic yield/Biological yield $\times 100$. (Biological yield = Grain yield + Straw yield).

The proline content in the leaves was estimated following the method described by Shivkumar et al. (2015). The 0.1 g of rice leaves ~~was~~ were ground with 5 ml of 3% sulfosalicylic acid, and the mixture was then filtered. To 2 ml of the filtered mixture in a test tube, 2 ml of ~~acid-~~ ninhydrin and 2 ml of glacial acetic acid were added. The mixture was mixed with a Vortex mixer and boiled at 100°C for 1 h. The mixture was then frozen in ice and combined with 4 ml of toluene, mixed, and then left to stand for 5-10 min. ~~Absorbance~~ The absorbance of the reddish pink up- per phase was recorded at 520 nm against a toluene blank.

The glycine betaine (GB) content was determined following the method of Bates et al. (1973) with minor modifications. Finely ground dry material (0.5 g) was mixed with 20 ml of deionized water and maintained with mechanical agitation for 48 h at 25°C; subsequently, the mixture was centrifuged at 10,000 rpm for 10 min using a refrigerated centrifuge. The supernatant was collected and stored in the freezer until the analysis. Samples were thawed and diluted (1:1 v/v) with 2N of sulphuric acid (H₂SO₄). An aliquot of 0.5 ml was collected and incubated for 1 h in ice; after which, 0.2 mL of cold potassium iodide-iodine (KI-I₂) was added and mixed gently. The samples were stored at 0 to 4°C for 16 h; after which, they were centrifuged at 10,000 rpm for 15 min at 0°C. The supernatant was discarded while ensuring not to shake it; during this time, the samples were maintained in ice to allow separation of the periodide complexes from the acid medium. The periodide crystals that formed were dissolved in 9 ml of 1,2-dichloroethane, and the mixture was left to sit for 2 h; after which, the absorbance was read at 365 nm. For the standard

GB curve, a stock comprising 1 mg/ml diluted in 1 N H₂SO₄ was prepared; concentrations of 50-200 mg/ml were used.

Abscisic acid (ABA) content is determined by following the method described by Plessis et al. (2011). 100 mg of the leaves ~~were grinded~~ ground and mixed with 2 ml of extraction solvent (acetone, water, acetic acid, 80/19/1, v/v/v). The supernatant was recovered by centrifugation and the pellet was rinsed with 1 ml of extraction solvent. The extraction solvent was evaporated and the residue was resuspended in 0.5 ml of HPLC solvent (acetonitrile, water, acetic acid, 50/50/0.05, v/v/v) and ABA was quantified.

The Ascorbic acid content in the leaves was estimated following the method described by Shivkumar et al. (2015).

Statistical analysis

The data obtained were converted to the mean value for each parameter with three repetitions under a complete randomized design with factorial concept and statistically analyzed as described by Pans and Sukahtme (1978) factorial concept of the experiment.

Results and Discussion

~~Drought~~ Drought stress at the typical stage of rice led to a reduction in maturity period and yield. Cultivation of ~~direct seeded~~ direct-seeded rice under rainfed ~~condition~~ conditions usually faces a water deficit during its growth stage. Growing of rice genotypes having the tolerance capacity of drought would be the beneficial approach for getting comparatively higher ~~yield~~ yields under ~~short-term~~ short-term water ~~limitation~~ limitations at a specific stage. Hence, 10 ~~varities~~ varieties have been grown and their performance under water deficit stress at the tillering stage was analysed.




Fig. 1. (A) Plant height as affected by water deficit stress at the tillering stage and control. ± Bars indicate SD (n = 10) (B) Leaf area index as affected by water deficit stress at the tillering stage and control. ± Bars indicate SD (n = 10) (C) Number of stomata per micrometer of leaf area as affected by water deficit stress at tillering stage and control. ± Bars indicate SD (n = 10) (D) Number of panicles per plant as affected by water deficit stress at the tillering stage and control. ± Bars indicate SD (n = 10)

The results of 10 varieties of their four morphological traits showed that the variety Purna was lower plant stature (Fig. 1A), higher leaf area index (Fig. 1B), higher number of stomata per micrometer (Fig. 1C), and higher number of panicles per plant (Fig. 1D). These are the traits which support the plants their performance under stage specific stress. Fleury et al. (2010) reported that genotypic variance in wheat increase tolerance capacity of drought. It has been indicated that breeding strategy with genotypic selection might increase the tolerance against drought (T. R. Sinclair, 2011). Morphological traits might have the greater influence on drought tolerance and phenotyping of crops in the genomic era provided higher impact on yield and ecosystem (R. Tuberosa, 2012).




Fig. 2. (A) Ionic leakage as affected by water deficit stress at the tillering stage and control. \pm Bars indicate SD (n = 10) (B)Relative water content as affected by water deficit stress at the tillering stage and control. \pm Bars indicate SD (n = 10) (C) Chlorophyll content as affected by water deficit stress at the tillering stage and control. \pm Bars indicate SD (n = 10) (D) harvest index as affected by water deficit stress at the tillering stage and control. \pm Bars indicate SD (n = 10)

The ionic leakage was observed lower in Purna among 10 treated ~~varities-varieties~~ under water deficit ~~condition-conditions~~ in comparison to control (Fig. 2 A) and relative water content (Fig. 2 B) Chlorophyll content (Fig. 2 C) ~~were-was~~ recorded (69.66%) and (31 SPAD value) respectively, in Purna and they were higher among 10 treated ~~varities-varieties~~ while in control relative water content (RWC) maintained at >80% and chlorophyll content was higher in all 10 varieties compared to treated plants. The percentage harvest index showed that Purna performed better under ~~stage-specific~~ stage-specific water stress among 10 ~~varities-varieties~~ while in control harvest index ~~were-was~~ higher in all cases compared to treated plants (Fig. 2 D). The water deficit stress delayed panicle initiation and increased ionic leakage led to membrane damage, decreased

respiration as well as ATP production in rice (Kadam et al., 2017). Low water potential under drought is ~~the~~ a common ~~phenomena~~ phenomenon and is genotypically regulated as per ~~the~~ tolerance of the plant. It affects the anatomy of leaves and their ultrastructure ~~is~~ highly influenced under low relative water content ~~led~~ leading to a lower number of stomata per micrometer of leaf (Upadhyaya and Panda, 2019). Reduced leaf area and chlorophyll content, leaf rolling, stomatal closure, and cutinized layer on the leaf surface induce early flowering and a decline harvest index (Singh et al., 2012). It has been reported that a short period of drought in rice reduced panicle length and grain filling duration ~~resulted~~ resulting in a loss in grain yield as well as harvest index. ~~Reduced~~ A reduced number of stomata and higher ionic leakage followed by lower stomatal conductance hinders the exchange of gases, resulting in poor seed setting and reduced grain size (Wei et al., 2017). Water deficit ~~at~~ in the short term negatively affected the RWC which created an osmotic imbalance, lipid peroxidation, ~~and~~ membrane damage, and in severe cases, necrosis ~~spot spots~~ has been also reported in rice (Zhu et al., 2020; Panda et al., 2021).

Osmotic imbalance under low RWC is genotypically regulated in ~~plant~~ plants. Some plants have their physiological and biochemical mechanism to synthesize osmolytes and protect from water stress. The simultaneous processes of photosynthesis and transpiration continued with lower stomatal resistance and higher osmoprotectant under ~~water limited~~ water-limited condition ~~conditions~~ are ~~one~~ some of the most important ~~trait~~ traits to select ~~genotype~~ genotypes for different environmental ~~condition~~ conditions (Kamoshita et al., 2008). The acclimatization of genotypes in response to the changing environments is the Genotype-environment interactions ($G \times E$). These two sources of variation might be considered for varietal evaluation for breeding and cultivation across environments (Yan et al., 2003). In ~~the~~ present study, Purna showed better among 10 ~~varities~~ varieties in the synthesis of Proline, Glycine betaine which induced by increased amount of abscisic acid (Fig. 3). The increased amount of ascorbic acid was significant impact on stress mitigation in case of selected 10 ~~varities~~ varieties (Fig. 3D). The content of Proline, Glycine betaine, abscisic acid and ascorbic acid were higher in Purna compared to other nine ~~varities~~ varieties while ascorbic acid was higher in all 10 varieties under control. These two osmolytes-Proline, and Glycine ~~betaine~~ betaine have a greater impact on ~~the~~ hydration capacity of ~~cell~~ cells which ~~led~~ lead to higher ~~yield~~ yields under ~~green house~~ greenhouse condition ~~conditions~~ in comparison to ~~the~~ other nine ~~varities~~ varieties.

Accumulating osmolytes maintain turgidity and ~~exposed-expose~~ the leaves ~~for getting to get~~ photosynthetic active radiation and assimilate translocation. Proline ~~is a well-known well-known~~ vital osmoprotectant increased in rice under water deficit conditions and it helps the plant to continue stomatal conductance and maintain leaf turgidity (Kumar et al., 2016).




Fig. 3. (A) Proline content as affected by water deficit stress at the tillering stage and control. \pm Bars indicate SD (n = 10) (B) Glycine betaine content as affected by water deficit stress at the tillering stage and control. \pm Bars indicate SD (n = 10) (C) Abscisic acid content as affected by water deficit stress at the tillering stage and control. \pm Bars indicate SD (n = 10) (D) Ascorbic acid content as affected by water deficit stress at the tillering stage and control. \pm Bars indicate SD (n = 10)

Proline is one of the important imino ~~acid-acids which that acts~~ actin cellular processes in two ways - a component of protein and an osmoprotectant under drought. Hence, the content of proline would be the-an important tool for screening of plants under drought and might ~~taken be taken~~ be taken as a biochemical marker for the analysis of rice genotypes (Mishra et al., 2019). The increased proline content in Purna resulted in hydration in cell cells, thus ~~increased-increasing~~ increased relative water content

which continued the metabolism under ~~short-term~~ short-term water deficit stress (Fig 3A). The ABA and osmolytes viz. proline, glycine betaine, etc. reduced the oxidative damage and ionic leakage under drought stress in plants (Song, J.-X. et al., 2017). Water deficit stress increased ABA in all 10 ~~varities~~ varieties but it was higher in Purna which might influence the stomata closure and ~~activates-activated~~ the defense mechanisms and synthesis of osmolytes in plant ~~cell~~ cells. It has been reported that ~~an~~ increased concentration of ABA ~~induce-induces~~ the synthesis of ~~of~~ primary and secondary metabolites and ~~enhance-enhances~~ the biomass production (Fahad, S. et al., 2017). The Glycine betaine acts as ~~a~~ compatible solute and it has ~~a~~ major role in improving the water stress tolerance in plants. It has been well documented that Glycine betaine (GB) improved drought tolerance in various crops viz. wheat (Raza et al., 2014) sunflower (Hussain et al., 2008), etc., and increased yield compared to other ~~GB deficient~~ GB-deficient plants. The increased GB content remarkably enhances the osmotic balance, enzymes, and genes associated with drought resistance (Hussain et al., 2013). Hence, the maintained hydration and osmoticum in ~~the~~ cell led to ~~the~~ survival and yield of the plant under ~~short-term~~ short-term water deficit stress.

Conclusion

The abiotic stress particularly water deficit in ~~plant-plants~~ plants under rainfed ~~condition-conditions~~ is the major challenge for marginal ~~farmer-farmers~~ farmers for production and yield of crops. ~~Selection-The selection~~ of ~~drought-tolerant~~ drought-tolerant varieties might reduce severe loss caused by water stress. The variety Purna has been found better under ~~short-term~~ short-term water deficit among 10 ~~varities-varieties~~ varieties of rice. Hence, ~~drought-tolerant~~ drought-tolerant variety-varieties can be produced through ~~a~~ breeding program with the help of Purna.

References:

1. Anjum, S. A. et al. (2017). Drought Induced Changes in Growth, Osmolyte Accumulation and Antioxidant Metabolism of Three Maize Hybrids. Front. Plant Sci.: 08.
2. Bandurska, H., Niedziela, J., Pietrowska-Borek, M., Nuc, K., Chadzinikolau, T., Radzikowska, D. (2017). Regulation of proline biosynthesis and resistance in drought stress in two barley (*Hordeum vulgare* L.) genotypes of different origin. Plant Physiology and Biochemistry, 118, 427-437.

3. Bates, L. S., Waldren, R. P., Teare, I. D. (1973). Rapid determination of free proline for water-stress studies. *Plant and Soil*, 39, 205-207.
4. Bhaskara GB, Nguyen TT, Yang T-H, Verslues PE. (2015). Dynamic proline metabolism and regulation in waterlimited environment. *Frontier in Plant Science*, 6, 384.
5. Fahad, S. et al. (2017). Crop production under drought and heat stress: plant responses and management options. *Front. Plant Sci.* 8, 1147.
6. Fleury, D., Jefferies S., Kuchel, H.P. (2010). Genetic and genomic tools to improve drought tolerance in wheat. *Journal of Experimental Botany*, 61(12), 3211-3222.
7. Henry, A., Wehler, R., Grondin, A., Franke, R., Quintana, M. (2016). Environmental and physiological effects on grouping of drought tolerant and susceptible rice varieties related to rice (*Oryza sativa*) root hydraulics under drought. *Ann. Bot.*, 118 (4): 711-724.
8. Hussain Wani, S., Brajendra Singh, N., Haribhushan, A., Iqbal, Mir, J. (2013). Compatible Solute Engineering in Plants for Abiotic Stress Tolerance - Role of Glycine Betaine. *Curr. Genomics* 14, 157-165.
9. Hussain, M., Malik, M. A., Farooq, M., Ashraf, M. Y., Cheema, M. A. (2008). Improving drought tolerance by exogenous application of glycine betaine and salicylic acid in sunflower. *J. Agron. Crop Sci.* 194, 193-199.
10. K. Ji, Y. Wang and W. Sun et al. (2012). Drought-responsive mechanisms in rice genotypes with contrasting drought tolerance during reproductive stage. *Journal of Plant Physiology*, 169(4): 336-344.
11. Kadam N N, Tamilselvan A, Lawas L M F, Quinones C, Bahuguna R N, Thomson M J, Dingkuhn M, Muthurajan R, Struik P C, Yin X Y, Jagdish S V K. (2017). Genetic control of plasticity in root morphology and anatomy of rice in response to water deficit. *Plant Physiology*, 174(4), 2302-2315.
12. Kamoshita, A., Babu, R. C., Boopathi, N. M. & Fukai, S. (2008). Phenotypic and genotypic analysis of drought-resistance traits for development of rice cultivars adapted to rainfed environments. *Field Crop Res.* 109, 1-23.
13. Kant, K. (2020). Free radicals and their enzymatic scavenger during moisture stress in plant. *International Journal of Science, Environment and Technology*, 9(6), 874-877.
14. Kant, K. (2023). Abscisic acid vis-a-vis water stress in plant. *International Journal of Science, Environment and Technology*, 12(4), 79-82.
15. Kaur, G.; Asthir, B. Molecular responses to drought stress in plants. *Biol. Plant* 2017, 61, 201–209.
16. Kumar A, Basu S, Ramegowda V, Pereira A. (2016). Mechanisms of drought tolerance in rice. In: Sasaki T. *Achieving Sustainable Cultivation of Rice*. Vol. 11. UK: Burleigh Dodds Science Publishing Limited

17. Mishra S S, Behera P. K., Panda D. (2019). Genotypic variability for drought tolerance-related morpho-physiological traits among indigenous rice landraces of Jeypore tract of Odisha, India. *J Crop Imp*, 33, 254-278.
18. Mishra, S. S., and Panda, D. (2017). Leaf traits and antioxidant defense for drought tolerance during early growth stage in some popular traditional rice landraces from koraput, India. *Rice Sci.* 24, 207-217.
19. Nawaz, M., and Wang, Z. (2020). Abscisic acid and glycine betaine mediated tolerance mechanisms under drought stress and recovery in *axonopuscompressus*: a new insight. *Scientific Report*, 10, 6942.
20. Panda, D., Mishra, S. S., and Behera, P. K. (2021). Drought tolerance in rice: focus on recent mechanisms and approaches. *Rice Sci.* 28 (2), 119-132.
21. Pandey, V and Shukla, A. (2015). Acclimation and Tolerance Strategies of Rice under Drought Stress. *Rice Science*, 22(4), 147-161.
22. Plessis, A., Cournol, R., Effroy, D., Silva Perez, V., Botran, L., Kraepiel, Y., Frey A., Sotta, B., Cornic, G., Leung, J., Giraudat, J., Marion-Poll, A., North, H.M. (2011). New ABA-hypersensitive *Arabidopsis* mutants are affected in loci mediating responses to water deficit and *Dickeyadadantii* infection. *Plos One*, 6, e20243.
23. R. Tuberosa (2012). Phenotyping for drought tolerance of crops in the genomics era. *Frontiers in Physiology*, 3.
24. Raza, M. A. S., Saleem, M. F., Shah, G. M., Khan, I. H., Raza, A. (2014). Exogenous application of glycine betaine and potassium for improving water relations and grain yield of wheat under drought. *J. soil Sci. plant Nutr.* 14, 34-364.
25. Sairam R K. (1994). Effect of moisture stress on physiological activities of two contrasting wheat genotypes. *Indian Journal of Experimental Biology*, 32: 594-597
26. Schafleitner, R., Gaudin, A., Rosales, R. O. G., Aliaga, C. A. A., Bonierbale, M. (2007). Proline accumulation and real time PCR expression analysis of genes encoding enzymes of proline metabolism in relation to drought tolerance in Andean potato. *Acta Physiologiae Plantarum*, 29, 19-26.
27. Serraj, R, McNally, K. L. and I. Slamet-Loedin et al. (2011). Drought resistance improvement in rice: An integrated genetic and resource management strategy. *Plant Production Science*, 14(10) 1-14.
28. Shivkumar, R. (2015). *Practical Plant Physiology*. Bio-green books, ISBN 10:9384337242, ISBN 13: 9789384337247.
29. Singh, C. M., Kumar, B., Mehendi, S., and Chandra, K. (2012). Effect of drought stress in rice: a review on morphological and physiological characteristics. *Trends Biosci.* 5, 261-265.

30. Song, J.-X. et al. (2017). Combined foliar application of nutrients and 5-aminolevulinic acid (ALA) improved drought tolerance in *Leymus chinensis* by modulating its morpho-physiological characteristics. *Crop Pasture Sci.* 68, 474-482.
31. Swamy, B. P. M. and Kumar, A. (2013). Genomics-based precision breeding approaches to improve drought tolerance in rice. *Biotechnology Advances*, 31(8), 1308-1318.
32. Szabados, L., Saviouré, A. (2010). Proline: A multifunctional amino acid. *Trends in Plant science*, 15, 89-97.
33. T. R. Sinclair (2011). Challenges in breeding for yield increase for drought. *Trends in Plant Science*, 16(6), 289-293.
34. Upadhyaya, H., and Panda, S. K. (2019). Drought stress responses and its management in rice. *Adv. Rice Res. Abiotic Stress Tolerance*, 177-200.
35. Wei, H., Chen, C., Ma, X., Zhang, Y., Han, J., Mei, H., et al. (2017). Comparative analysis of expression profiles of panicle development among tolerant and sensitive rice in response to drought stress. *Front. Plant Sci.* 8.
36. Yan, W. & Kang, M. S. *GGE Biplot Analysis: A Graphical Tool for Breeders, Geneticists, and Agronomists* (CRC Press, 2003).
37. Zhu, R., Wu, F., Zhou, S., Hu, T., Huang, J., and Gao, Y. (2020). Cumulative effects of drought–flood abrupt alternation on the photosynthetic characteristics of rice. *Environ. Exp. Bot.* 169, 1-14.