

## Effect of Seed Priming Treatments on Germination, Growth and Yield of Ajwain (*Trachyspermum ammi* L.)

### ABSTRACT

**Aim:** To discern the most effective seed priming treatment among several priming treatments, which provide early with maximum germination, prosperous growth and prodigious yield.

**Study Design:** Nine treatments with one control treatment in Randomized Block Design (RBD) with three replications.

**Place and Duration of Study:** Chilli and Vegetable Research Unit, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola (M.S), during the rabi season of 2022.

**Methodology:** The ajwain seeds were primed with treatments comprise of, GA<sub>3</sub> at 100 ppm and 200 ppm, KH<sub>2</sub>PO<sub>4</sub> at 2% and 3%. Moringa leaf extract at 5% and 7.5%, and tulasi leaf extract at 5% and 7.5%, hydropriming (soaking in water), control (without soaking), for 24 hours before sowing. After that, primed seeds shade dried for 4-5 hours. Then, that seeds were sown in the field.

**Results:** The outcomes unveiled fascinating insights. Notably, treating the seeds with 7.5% of moringa leaf extract (T<sub>6</sub>) revealed remarkable results, exhibiting the swiftest germination at 11.67 days, a staggering initial plant stand of 99.38%, and an impressive germination percentage of 93.33%. Meanwhile, the utilization of 200 ppm GA<sub>3</sub> led to maximum plant height of 89.70 cm. Furthermore, the application of 7.5% moringa leaf extract (T<sub>6</sub>) observed the minimum days to 50% flowering at 92.98 days and days to harvesting at 156.17 days, and boasting the maximum primary branches plant<sup>-1</sup> at 11.55. Among the myriad treatments, the application of 7.5% moringa leaf extract (T<sub>6</sub>) emerged as a standout, showcasing unparalleled results with maximum number of umbels and umbellets at 200.72 and 18.03, respectively. A prodigious yield of 10.49g plant<sup>-1</sup>, 2.67 kg plot<sup>-1</sup> and 11.86q ha<sup>-1</sup>.

**Conclusions:** These findings underscore the potency of specific seed priming treatments, particularly, the 7.5% moringa leaf extract (T<sub>6</sub>) and 200 ppm GA<sub>3</sub> (T<sub>2</sub>), in fostering robust ajwain seed germination, growth, accelerated plant development and prodigious yield.

**Keywords:** Ajwain, Seed priming, Moringa leaf extract, Tulasi leaf extract, Germination, Growth and yield.

### 1. INTRODUCTION

Ajwain, scientifically termed as *Trachyspermum ammi* L., stands as an annual herbaceous specimen within the esteemed medicinal cohort of the Apiaceae family. Flourishing predominantly in arid and semi-arid terrains characterized by saline-rich soil compositions, this botanical entity finds its cultivation prevalent across diverse locales such as Iran, Pakistan, Afghanistan, India, and select

regions of Europe. Cultivation methodologies exhibit seasonal nuances, with planting typically undertaken between October-November in India. While, the saline tract area of Vidarbha adjusts its sowing to the latter half of August, culminating in a harvest during December-January with residual moisture and rest of the ajwain growing area concentrating with October- November sowing, with irrigation facility.

The geographical expanse of Ajwain's cultivation in India extends over Gujarat, Rajasthan, Madhya Pradesh, Uttar Pradesh, Maharashtra, Bihar, and West Bengal, encompassing an approximate land area of 37,810 hectares and yielding around 27,920 tons of produce by the year 2020 [1]. This botanical's fruit harbours stimulant, antispasmodic, and carminative attributes, historically revered for its remedial efficacy against an array of ailments encompassing flatulence, dyspepsia, diarrhoea, abdominal pains, piles, bronchial afflictions, lack of appetite, galactagogue, asthma, and amenorrhoea [14].

The optimum developmental milieu for ajwain dictates specific climatic requisites, necessitating temperatures ranging between 15 to 27°C and a relative humidity spanning 60 to 70% during its growth phase. Notably, the phase of seed formation demands a relatively warmer climate. Remarkably, ajwain displays a moderate resistance to drought, enabling its cultivation in both *Kharif* and rainy seasons. Its nutritional matrix showcases pivotal vitamins like riboflavin, thiamine, nicotinic acid, and carotene, coupled with essential minerals encompassing calcium, phosphorus, iron, potassium, and salt [8]. The quintessential essential oil harboured within ajwain, predominantly constituting 35- 60% of its composition, is thymol. This oil not only exudes a redolent fragrance, but also stands as a potent bearer of antiseptic properties.

Seed germination and seedling establishment are critical stages in plant life, and the successful establishment of a plant depends on the rapid and uniform germination of seed under adverse environmental conditions. In ajwain the rate of deterioration of seed quality parameters are rapid during storage, resulting in poor germination, vigour and establishment of crops. Seed priming is an excellent technique, which improves germination and better crop stand. Thus, Seed priming has shown its promise in enabling the seeds to overcome various biotic and abiotic stresses. Therefore, it is a crucial pre-sowing procedure for improving germination and uniform emergence of a crop in normal and less-than-ideal circumstances. By getting the seeds to a point when metabolic processes have already been started and giving them a head start over unprimed seeds, pre-sowing priming is known to boost the performance of the seeds. The primed seeds can continue where they left off in the remaining stages of germination (radical protrusion) after receiving more imbibition. It has been established that, seeds produced from such crop have higher seed quality than the crops which has been raised from non-primed seeds.

## **2.MATERIAL AND METHODS**

The present investigation was carried out during the rabi season of 2022-2023, in the research field of Chilli and Vegetable Research Unit, Dr.Panjabrao Deshmukh Krishi Vidyapeeth,

Akola. The experiment consists of ten treatments and were replicated three times using a Randomized Block Design (RBD).

## 2.1 Treatment details

Treatments includes, T<sub>1</sub> (GA<sub>3</sub> @100 ppm), T<sub>2</sub> (GA<sub>3</sub> @200 ppm), T<sub>3</sub> (KH<sub>2</sub>PO<sub>4</sub> @ 2%), T<sub>4</sub> (KH<sub>2</sub>PO<sub>4</sub> @ 3%), T<sub>5</sub> (Moringa leaf extract @5%), T<sub>6</sub> (Moringa leaf extract @ 7.5%), T<sub>7</sub> (Tulasi leaf extract @ 5%), T<sub>8</sub> (Tulasi leaf extract @ 7.5%), T<sub>9</sub> (Soaking in water / Hydropriming), T<sub>10</sub> (Absolute Control/ Without soaking). The seeds are primed for 24 hours before sowing. Then, the seeds are shade-dried for 4-5 hours, to reduce the excess moisture level. After priming, immediately seeds were sown in the field. In each plot, the seeds are sown in the middle of every ridge with 70 hills and each hill has approximately 25-30 seeds, as a characteristic of the Apiaceae family.

## 2.2 Procedure for preparation of the priming treatments

Initially stock solution of 1000ppm prepared by dissolving 1g GA<sub>3</sub> in 5ml of ethanol, added to 995ml of distilled water. Then, GA<sub>3</sub> solutions of 100ppm and 200ppm were prepared by dissolving 10ml and 20ml of GA<sub>3</sub> in 90ml and 80ml water, respectively. KH<sub>2</sub>PO<sub>4</sub> solutions of 2% and 3% were prepared by dissolving 2g and 3g of KH<sub>2</sub>PO<sub>4</sub> in 100ml of water and the seeds were soaked for 24 hours. Moringa leaf extract and Tulasi leaf extract were prepared according to [9]. Fresh moringa leaves and Tulasi leaves were washed thoroughly in tap water and crushed into fine paste by using a mortar and pestle, each 5g and 7.5g of the sample was ground by a blender with 100ml of water separately. The ground mixture is filtered through a fine cotton cloth, which constitutes 5% and 7.5% of moringa leaf extract and Tulasi leaf extract, respectively. Seeds are soaked for 24 hours before sowing.

## 3. RESULTS AND DISCUSSION

### 3.1 Germination parameters

Table 1 illustrates the impact of various seed priming treatments on key germination parameters, such as initial plant stand (%), days required for germination, and germination percentage (%). The data unequivocally demonstrates that, employing diverse priming methods significantly enhanced seed germination attributes when compared to the control treatment, T<sub>10</sub>. Particularly, seeds primed with moringa leaf extract at 7.5% (T<sub>6</sub>) exhibited the maximum initial plant stand (99.38%), which was at par with KH<sub>2</sub>PO<sub>4</sub> at 3% (T<sub>4</sub>) and GA<sub>3</sub> at 200 ppm (T<sub>2</sub>), which registered 98.56% and 98.08% initial plant stand, respectively. Conversely, ajwain seeds without soaking (absolute control) T<sub>10</sub>, displayed the minimum initial plant stand, significantly only at 80.67%, compared to other priming treatments. This might be due to the fact that, a higher concentration of calcium and other mineral contents in moringa (*Moringa oleifera*) leaves might be responsible for promoting initial plant stand rate. Moreover, a higher zeatin concentration in *Moringa oleifera* leaves (5-200 Hg g of fresh weight) and enhanced mobilizations of metabolites/ inorganic solutes to germinating plumule, which results in enhanced growth. The findings align with previous studies by [3,13,6].

Remarkably, organic priming treatments notably reduced the days required for germination. Seeds primed with 7.5% moringa leaf extract ( $T_6$ ) sprouted in the minimum duration (11.67 days), akin to  $KH_2PO_4$  at 3% ( $T_4$ ) with 12.67 days and moringa leaf extract at 5% ( $T_5$ ) with 13.33 days. In contrast,  $T_{10}$  seeds took 18.33 days for germination, significantly maximum than other treatments. This could be because, calcium in moringa leaf extract promotes quicker germination and functions as an enzyme cofactor during germination. The activity of several phosphatases and kinase enzymes involved in signal translation is modulated by calcium, which also helps in reducing the days to germinate. Similar results were found by [13,6].

Furthermore, seeds primed with 7.5% moringa leaf extract ( $T_6$ ) demonstrated the maximum germination rate at 93.33%, akin to  $KH_2PO_4$  at 3% ( $T_4$ ) with 91.00% and moringa leaf extract at 5% ( $T_5$ ) with 88.33%. Conversely, the control treatment ( $T_{10}$ ) substantially minimum germination percentage to 79.00% compared to other priming methods. This could be due to, the role of phytohormones, amino acids, and mineral elements present in moringa leaf extract, which positively influenced germination parameters. Furthermore, priming treatment promotes the mobilization of seed reserve from endosperm to embryo, resulting in better performance of germination. These outcomes concur with findings from studies conducted by [6].

### 3.2 Growth parameters

Table 1 depicts the influence of various seed priming treatments on growth parameters of ajwain. There were noteworthy differences observed among these treatments. Specifically, seeds treated with  $GA_3$  at 200ppm ( $T_2$ ) resulted in plants with the maximum height at 89.70cm, statistically comparable to treatments involving  $GA_3$  at 100ppm, moringa leaf extract at 7.5% ( $T_6$ ), and  $KH_2PO_4$  at 3% ( $T_4$ ), which produced plant height of 87.17cm, 85.70cm, and 84.83cm, respectively. Contrastingly, the control treatment ( $T_{10}$ ) without soaking exhibited the minimum plant height at 69.27cm. This might be a consequence of  $GA_3$ 's increase in cell division and stem elongation. The phytohormone gibberellin is in charge of controlling plant height. By making the cell wall more flexible,  $GA_3$  treatments are also successful in promoting vegetative development. These findings align with studies by [11,7,5].

Regarding the number of primary branches plant<sup>-1</sup>, statistically significant differences were evident. Seeds treated with 7.5% moringa leaf extract showed the maximum number of primary branches plant<sup>-1</sup> at 11.95, followed by those treated with 3%  $KH_2PO_4$  at 10.70. Conversely, the control treatment ( $T_{10}$ ) exhibited the minimum number of branches plant<sup>-1</sup> at 7.17. This might be because, moringa (*Moringa Oleifera*) leaves have larger concentrations of calcium, zeatin concentration (5-200 Hg g of fresh weight) and other minerals, which might encourage the growth of more branches. Furthermore, this could also be the result of higher or enhanced mobilizations of metabolites/inorganic solutes to germinating plumule, which leads to enhanced growth of primary branches. These results correspond with findings by [12].

The treatment involving moringa leaf extract at 7.5% ( $T_6$ ) remarkably led to the minimum duration of 92.98 days to achieve 50 percent blooming and significantly reduced the harvesting duration to 156.17 days, which was akin to  $KH_2PO_4$  at 3% ( $T_4$ ) which recorded 95.36 days and 158.51 days. And also akin with moringa leaf extract at 5% ( $T_5$ ) required 96.90 days and 160.57 days

for 50 percent blooming and harvesting, respectively. In contrast, the control treatment ( $T_{10}$ ) took the maximum duration at 105.32 days to achieve this stage. This might be due to, moringa leaf extract-primed seeds provided an energizing start, early crop establishment led to early 50% flowering and harvesting. These outcomes coincide with studies by [4,13,6].

### 3.3 Yield parameters

In examining the outcomes depicted in Table 1, notable variations in yield parameters resulting from diverse seed priming treatments compared to the control treatment ( $T_{10}$ ) were evident. Specifically, Treatment ( $T_6$ ) utilizing moringa leaf extract at 7.5% exhibited statistical significance over all other treatments, yielding 200.72 umbels plant<sup>-1</sup>. Following closely was Treatment ( $T_4$ ) involving  $KH_2PO_4$  at 3% with a count of 187.23 umbels plant<sup>-1</sup>. Conversely, the control treatment ( $T_{10}$ ) without soaking displayed a notably minimum of umbels plant<sup>-1</sup> at 162.63. This might be due to, moringa extract is rich in zeatin-like cytokinin that induces cytokinin bio-synthesis improvement and the number of photosynthetic active leaves and number of umbels. These results align with findings by [12].

Furthermore, Treatment ( $T_6$ ) with moringa leaf extract at 7.5% generated 18.03 umbellets umbel<sup>-1</sup>, comparable to  $KH_2PO_4$  at 3% ( $T_4$ ) and  $GA_3$  at 200ppm ( $T_2$ ) that yielded 17.27 and 17.13 umbellets umbel<sup>-1</sup>, respectively. In contrast, the control treatment ( $T_{10}$ ) exhibited the minimum count of 13.36 umbellets umbel<sup>-1</sup>. This might be because, improved vegetative growth which coupled with increased photosynthesis on one hand and greater mobilization of photosynthates towards reproductive sites increased the number of umbellet per umbel on other. These results corroborate with the findings of [12].

Additionally, Treatment ( $T_6$ ) employing moringa leaf extract at 7.5% significantly increased yield plant<sup>-1</sup>, plot<sup>-1</sup>, and hectare<sup>-1</sup>. Seed employed with moringa leaf extract at 7.5% treatment produced the maximum yield plant<sup>-1</sup> at 10.49g, 2.67kg plot<sup>-1</sup>, and 11.86 q ha<sup>-1</sup>, paralleled by Treatment ( $T_4$ ) using  $KH_2PO_4$  at 3%, generating 9.87g plant<sup>-1</sup>, 2.62kg plot<sup>-1</sup>, and 11.61q ha<sup>-1</sup>. Similarly, also with  $GA_3$  at 200ppm ( $T_2$ ) yielded 9.74g plant<sup>-1</sup>, 2.59kg plot<sup>-1</sup>, and 11.47q ha<sup>-1</sup>. In contrast, the control treatment ( $T_{10}$ ) yielded the minimum at 5.30g plant<sup>-1</sup>, 1.74kg plot<sup>-1</sup>, and 7.71q ha<sup>-1</sup>. This might be due to, greater partitioning of photoassimilates to developing grains. Moringa leaf extract's noteworthy benefits might be linked to its impact on growth parameters, which raised photosynthesis and sink capacity by utilizing photoassimilates from leaves, and translocation to produce high-quality and abundant fruit. Furthermore, zeatin-like cytokinin, which increases cytokinin biosynthesis and the number of photosynthetic active leaves, is abundant in moringa extract. It also impacted on the yield. These findings are supported by research conducted by [2,10].

**Table 1. Effect of seed priming treatments on germination, growth and yield parameters of ajwain plant**

Tre.	Treatment details	Initial plant stand (%)	Days to germination	Germination percentage (%)	Plant height (cm)	Number of primary branches	Days to 50% flowering	Days to harvesting	Umbels/plant <sup>1</sup>	Umbellets/umbel <sup>-1</sup>	Yield plant <sup>-1</sup> (g)	Yield Plot <sup>-1</sup> (kg)	Yield ha <sup>-1</sup> (q)
T <sub>1</sub>	GA <sub>3</sub> at 100ppm	96.30	16.33	82.33	87.17	8.07	99.89	165.18	170.87	14.78	6.53	1.88	8.33
T <sub>2</sub>	GA <sub>3</sub> at 200ppm	98.08	14.33	87.67	89.70	10.63	97.90	162.33	182.29	17.13	9.74	2.59	11.47
T <sub>3</sub>	KH <sub>2</sub> PO <sub>4</sub> at 2%	95.42	14.67	84.33	77.01	8.48	99.08	166.10	172.35	14.90	7.23	2.03	9.01
T <sub>4</sub>	KH <sub>2</sub> PO <sub>4</sub> at 3%	98.56	12.67	91.00	84.83	10.70	95.36	158.51	187.23	17.27	9.87	2.62	11.61
T <sub>5</sub>	Moringa leaf extract at 5%	95.08	13.33	88.33	80.60	9.56	96.90	160.57	180.91	15.73	8.30	2.32	10.30
T <sub>6</sub>	Moringa leaf extract at 7.5%	99.38	11.67	93.33	85.70	11.95	92.98	156.17	200.72	18.03	10.49	2.67	11.86
T <sub>7</sub>	Tulasi leaf extract at 5%	92.15	15.67	83.67	73.37	8.15	100.32	166.21	168.67	15.00	6.93	1.92	8.52
T <sub>8</sub>	Tulasi leaf extract at 7.5%	88.94	15.00	86.33	77.10	8.75	98.70	163.91	175.65	15.85	7.43	2.09	9.27
T <sub>9</sub>	Hydropriming (Soaking in water)	85.51	15.67	85.00	76.36	7.99	100.99	167.25	166.68	14.60	6.23	1.86	8.25
T <sub>10</sub>	Absolute control (Withoutsoaking)	80.67	18.33	79.00	69.27	7.17	105.32	171.93	162.63	13.36	5.30	1.74	7.71
	<b>SE (m)±</b>	<b>1.03</b>	<b>0.78</b>	<b>1.60</b>	<b>2.22</b>	<b>0.26</b>	<b>1.33</b>	<b>1.09</b>	<b>1.20</b>	<b>0.31</b>	<b>0.25</b>	<b>0.03</b>	<b>0.15</b>
	<b>CD at 5%</b>	<b>3.08</b>	<b>2.32</b>	<b>4.77</b>	<b>6.62</b>	<b>0.79</b>	<b>3.97</b>	<b>4.45</b>	<b>3.58</b>	<b>0.92</b>	<b>0.76</b>	<b>0.10</b>	<b>0.47</b>

#### 4.CONCLUSION

The investigation results indicates that, all treatments exhibited a broader range of variability in germination, growth and yield parameters compared to the control treatment. Specifically, treating seeds with a 7.5% concentration of moringa leaf extract resulted in significant enhancements in germination metrics such as initial plant stand (%), germination percentage (%), and minimum days required for germination.

Concerning growth factors, priming seeds with GA<sub>3</sub> at 200ppm led to a significant increase in plant height (cm). Additionally, using moringa leaf extract at 7.5% resulted in the maximum number of primary branches plant<sup>-1</sup>, the minimum days to 50 percent blooming, and days to harvesting. Specifically, it also exhibited substantial amplifications in yield parameters such as the number of umbels plant<sup>-1</sup>, umbellets umbel<sup>-1</sup>, yield plant<sup>-1</sup>(g), yield plot<sup>-1</sup>(kg), and yield hectare<sup>-1</sup>(q). Overall, it's evident that employing a 7.5% concentration of moringa leaf extract facilitates early and extensive germination and prodigious yield. While, GA<sub>3</sub> at 200ppm promotes plant height. These results are however, based on one year experimentation. Hence further studies using different priming treatments with different concentration will be helpful for the confirmation of the results.

#### REFERENCES

1. Anonymous. Area and Production of Ajwain, National Research Centre on Seed Spices, Ajmer, Rajasthan, India. 2020. ([www.nrcss.icar.gov.in](http://www.nrcss.icar.gov.in))
2. Azooz MM, Shaddad MA and Abdel-Latef AA. The accumulation and compartment action of proline in relation to salt tolerance of three sorghum cultivars. *Ind J of Plant Physio.* 2004; 5(9): 1-8.
3. BasraSMA, Iftikhar MN and Afzal I. Potential of moringa (*Moringa oleifera*) leaf extract as a priming agent for hybrid maize seeds. *IntJ of Agril and Bio.* 2011;13(6): 1006-1010.
4. Farooq M, Basra SMA, Rehman H and Hussain M. Seed priming with polyamines improve the germination and early seedling growth in fine rice. *J of New Seeds.*2006; 9(2): 145-155.
5. Jamshidian Z and Talat F. Effects of seed priming on morphological and phonological characteristics of the coriander (*Coriandrum sativum* L.). *Adv in Plants &Agril Research.* 2017; 7(6): 411-415.
6. KhanS,Ibrar D, Bashir S, Rashid N, Hasnain Z, Nawaz M, *et al.*, The application of moringa leaf extract as a seed priming agent enhances the growth and physiological attributes of rice seedlings cultivated under a water deficit regime. *Plants.* 2022; 11(3): 261.
7. KumarS, Malik TP, and Tehlan SK. Effect of gibberellic acid on growth and seed yield of coriander (*Coriandrum sativum* L.). *Int J of Curr Microbio AppSci.* 2018; 7(9): 2558-2566.
8. Pruthi JS. *Minor spices and condiments: crop management, and post-harvest technology.* ICAR, New Delhi. 2001. PP. 124-133.
9. Reddy, A. N; A.K. Chaurasia and E. Sutnga, 2020. Assessment of Different Priming Methods on Germination and Quality Parameters of Coriander (*Coriandrum sativum* L.). *Int. J. Curr. Microbiol. App. Sci.* 9(11): 3030-3038.

10. RehmanHU, Basra S, Rady MM, Ghoneim A andWang Q. Moringa leaf extract improves wheat growth and productivity by affecting senescence and source-sink relationship. *Int J of Agriland Bio.* 2017; 19(3): 479–484.
11. Singh P, PandeyAK and Khan AH. Effect of seed priming on growth, physiology, and yield of lentil (*Lens Culinaris* Medie) Cv. Ndl-1. *J of Pharmaco and Phytoche.* 2017; 6(6): 717-719.
12. Soliman YM. Growth and yield of caraway plants (*Carum carvL.*) As influenced by organic fertilizer and moringa leaves extract under newly reclaimed soils. *Int J of Agril Sci.* 2022; 4(4): 187-196.
13. Yasmeen A, Basra SMA, Wahid A, Nouman W and Rehman H. Exploring the potential of Moringa oleifera leaf extract (MLE) as a seed priming agent in improving wheat performance. *Turkish J of Bot.* 2013;37(2): 512-520.
14. Zarshenas MM, Moein M, Samani SM and Petramfar P. An overview on ajwain (*Trachyspermumammi*) pharmacological effects; modern and traditional. *J of Nat Rem,* 2014; 14 (1): 98-105.