

## Original Research Article

### Impact of integrated nutrient management on soil properties under Long-term fertilizer experiment in Vertisol.

#### ABSTRACT

**Aims:** To study the effect of continuous fertilization on soil properties and soil carbon stock, in a long-term experiment under [wheat-soyabean](#) cropping system in Vertisol.

**Study design:** [Experiment](#) was laid in a Randomized block design and consisted of three replication and nine treatments.

**Place and Duration of Study:** Soil samples were collected during April 2022 following the harvest of Wheat from the on-going long-term fertilizer experiment at Jawahar Nehru Krishi Viswa Vidyalaya, Jabalpur, Madhya Pradesh, India

**Methodology:** The soil samples collected from two depths (0-15cm, 15-30cm) and were analyzed for the properties such as pH, electrical conductivity, available N, available P, available K using standard procedures. Further a correlation study was carried out to determine the relationship between all the parameters.

**Results:** The results indicated that out of all the treatments continued application of 100% NPK+ FYM had significant effect (at  $P < 0.05$ ) on all the parameters. This treatment had the highest value for in both 0-15cm and 15-30 cm: SOC% (0.79%, 0.73%), available N ( $319.3 \text{ kg ha}^{-1}$ ,  $275.00$ ), available P ( $40.11 \text{ kg ha}^{-1}$ ,  $37.85 \text{ kg ha}^{-1}$ ), available K ( $326.33 \text{ kg ha}^{-1}$ ,  $302.9 \text{ kg ha}^{-1}$ ).

**Conclusion:** Continued application of both 100% NPK along with FYM has a significant positive impact on build up of nutrients in the soil. This indicates that integrated application of nutrition assures productivity soil in a long.

*Keywords: Long-term fertilizer experiment, Vertisol, integrated nutrient management, chemical fertilizers.*

#### 1. INTRODUCTION

The future of agriculture is tainted with scarcity of land and shortage of water, while being burdened with the responsibility of feeding the expanding population in upcoming years. Under such [scenario](#) use of chemical fertilizers and [high yielding](#) varieties serve as [rescue](#) by increasing the productivity of crops from limited land resources. However, the continued and non-judicious use of chemical fertilizers over [the](#) years may result in serious environmental issues. Relentless use of chemical fertilizers has been reported to damage the physiochemical properties of soil by raising soil acidity, which accelerates the degradation of soil health, productivity, stability, and sustainability [1]. These inorganic fertilizers are also water soluble, which allows them to percolate into the subsurface layers and change the soil's inherent characteristics [2]. Excess application or unbalanced application of nitrogen may result to eutrophication, contamination of ground water with nitrates and emission of greenhouse gases such as nitrous oxides [3]. Application of phosphate

fertilizer over long ~~period-of-time~~time in cultivated land, can be a major source of potentially toxic trace metals like lead, cadmium, and arsenic. These trace elements may build up in soils and move up the food chain [4]. Phosphatic fertilizers like nitrogenous fertilizers when applied to soil over a long ~~period-of-time~~period also contribute to eutrophication significantly [5]. Sole application of chemical fertilizers over ~~the~~ years have also been reported to enhance the deficiency of soil organic carbon as well as secondary and micronutrients which have led to gradual reduction in productivity over years, even after application of fertilizers [6]. Long-term experiments have proved that intensive cropping systems are exhibiting signs of "fatigue," as seen by yields that are stagnant or dropping. One explanation proposed for this yield stagnation is the reduction in the quantity and quality of soil organic matter (SOM)[7]. Thus, an alternative source of nutrient management is necessitated which is efficient enough to fulfill the growing food demand at an affordable cost while maintaining the sustainability of soil by improving the soil organic carbon content.

Organic manures are known to nourish soil health by providing essential nutrients, enriching the soil organic carbon, improving physical properties and boosting microbial activity since ancient times. Since organic manures are locally available and are made of in-farm products they are cost effective and easily available. But neither organic manure nor chemical fertilization alone can sustain ~~the~~ productivity on ~~the~~ long run hence integrated nutrient application turns out to be a promising solution. Numerous studies have been carried out throughout ~~the~~ world to evaluate the impact of integrated nutrient management on soil [8-11]. It has been found that ~~long-term~~long-term application of integrated nutrients not only promotes soil health but also helps in carbon sequestration by enriching ~~the soil~~the soil's organic carbon. Integrated application of nutrients has different ~~impact~~impacts on different cropping systems, soil type, and duration of application. Thus, ~~an~~ assessment of impact of integrated nutrient management on a particular soil order under certain nutrient management ~~system~~systems and cropping system is necessary. Vertisol is one of ~~the~~ most prevalent soil orders in India existing in central India with wheat-soyabean as a common cropping system. ~~Majority~~The majority of farmers in these areas cultivate soybeans without using fertilizer since they are aware of the impact the legume has on the wheat crop that follows. Wheat yields are low in this region due to uneven application of NPK fertilizers and almost non-existent usage of organics, which also degrades soil fertility. Keeping these points under consideration a study was carried out to test the hypothesis that ~~the~~ application of both organic and inorganic fertilizers over ~~the~~ long term has ~~a~~ positive impact on ~~the~~ chemical properties of soil on ~~the~~ long run.

## 2. MATERIAL AND METHODS:

### 2.1 Site description:

The current study was based on a long-term fertilizer experiment under the aegis of ~~the~~ All India Coordinated Research Project (AICRP on LTFE) which was initiated in the year 1972, in a Vertisol at the agricultural research farm of Department of Soil Science and Agricultural Chemistry, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh. The experimental site had ~~a~~ semi-arid and subtropical climate with ~~an~~ average yearly rainfall of 1350 mm and had medium-black soil in the Kheri series of the Typic Haplustert (Vertisol) with fine montmorillonite in ~~the~~ hyperthermic family. The initial soil properties at the initiation of the experiment in the year 1972 ~~is-are~~ given in Table no.1. The experiment consists of 10 treatments laid in randomized block design with four replications. For the present study nine out of ten treatments were selected which are: Control (no fertilization), 50% NPK, 100% NPK, 150%NPK, 100% NPK +Zn, 100% NP, 100% N, 100% NPK+FYM, 100%NPK-S. The recommended dose of fertilizers for soyabean and wheat (20:80:20 and 120:80:40 kg ha<sup>-1</sup> respectively). Under 100% NPK treatment, FYM was applied to the soybean crop at a

rate of 10 mg ha<sup>-1</sup> y<sup>-1</sup> on a dry weight basis. On a dry weight basis, FYM on an average contained 25% C, 0.95% N, 0.55% P, and 0.71% K. Crops were managed as per the standard recommended agronomic practices. Yield data were obtained after threshing of the mature soybean and wheat crops when they had moisture content of 7–10%.

**Table 1: Initial soil characteristics:**

Soil Properties	Value
Soil pH (Soil: water 1:2.5)	7.60
Electrical conductivity (1:2.5, ds m <sup>-1</sup> )	0.18
Organic carbon (g kg <sup>-1</sup> )	5.7
Bulk density (Mg m <sup>-3</sup> )	1.30
Available N (kg ha <sup>-1</sup> )	193.0
Available P (kg ha <sup>-1</sup> ) (Olsen)	7.6
Available K (kg ha <sup>-1</sup> )	370.0

### 2.3 Soil sampling and analysis:

In the current study soil samples were collected from surface (0-15 cm) and sub surface (15-30 cm) from nine different treatments after the harvest of wheat after the 50<sup>th</sup> cropping cycle of wheat-soybean rotation in the year 2022. For analysis of chemical properties of soil ~~the soil~~ the soil samples were collected from four random points within each replication and a composite sample was prepared after mixing them ~~together~~. Further, the sample size of the mixed composite was reduced to 500g by quartering. The soil samples after collection were ~~air-dried~~ air-dried and processed before being subjected to chemical analysis. For determination of bulk density, the soil samples were collected through cylindrical cores. Samples were analyzed for pH (1:2.5 soil: water suspension)[12], electrical conductivity by conductivity meter[12], organic carbon by rapid titration method [13], Available N was estimated by alkaline permanganate method [14] and available P by Olsen's method [15], available potassium by ammonium acetate method [12].

**2.4 Statistical analysis:**The data was analysed using Microsoft Excel 2011 and Analysis of variance (ANOVA) was carried out using SPSS 16.0. Duncan's multiple range test (DMRT) at 5% level of probability was used to test the significance of differences between treatment means [16].

## 3. RESULTS AND DISCUSSION

### 3.1 Soil properties:

#### 3.1.1 Soil pH and electrical conductivity:

Long-term application of both integrated nutrition as well as chemical fertilizers alone significantly affected the soil pH across both the depths. The pH of soil under different nutrient management practices ranged from 7.31 to 7.61 in surface soil while it ranged from 7.35 to 7.66 in sub-surface soil (Table 2). The range of pH found in this study were similar to that reported by Suman et al. (2017)[17]. The stability of the soil reaction may have been caused by the high buffering capacity of the clayey soil and the presence of weak salts, such as carbonates or bicarbonates, which upon dissolution

release free cations. As a result, the pH of surface and subsurface soils was not significantly affected by the long-term application of integrated nutrients over the years [18]. The lowest pH was observed in case of 100% N in both the layers due continuous application of urea alone over the years which results in acidification of soil due to formation of nitrates. The acidification caused due to the sole application of nitrogen has reduced the initial soil pH to maximum extent as compared other treatments. The findings also showed that the pH of the soil was slightly lowered at both soil depths upon the addition of FYM and inorganic nutrients. Likewise, Chouvan et al. (2015)[19] found that the stabilizing effects of FYM may be the reason for stability of soil pH when balanced fertilizers were applied in addition to FYM. The electrical conductivity of soil ranged from 0.15 to 0.18 dSm<sup>-1</sup> in surface soil and 0.14 to 0.18 dSm<sup>-1</sup> in subsurface layer (Table 2). These findings are in congruence with that of Khandagle et al. [20]. It was observed that the EC values did not exhibit any significant variation which can be attributed to the strong buffering capacity of the soil and the low residual effect of applied inputs [21]. Among the treatments electrical conductivity was highest in case of 100% NPK + FYM may be due to the rise in base saturation of soil due to optimum rate of nutrients applied as compared to control [18].

### 3.1.2. Soil organic carbon:

The soil organic carbon under various nutrient management ranged from 0.43% in control to 0.79% in 100% NPK+FYM treatment in surface soil. In subsurface soil it ranged from 0.40% to 0.73% in control and 100% NPK+FYM respectively (Table 2). Similar range of organic carbon was reported by [22]. Increase in soil organic carbon content in 100% FYM +NPK also reported by Santhy et al., (2001) [23] and Singh et al., (2007) [8]. It was found that the organic carbon content of surface soil reduced after 50 years of continuous cultivation in control where as its value had considerably increased the most in case of 100% NPK+FYM followed by 150% NPK and 100% NPK as compared to the initial value. The organic carbon content increased with increase in fertilizer dose following the sequence 50% NPK < 100% NPK < 150% NPK. Soil organic carbon content declined with depth, since continuous cultivation enhanced and boosted the decomposition of plant organic residues at surface level. The highest value of soil organic carbon was observed when organic and inorganic fertilizers were added in combination because it directly added more organic carbon through FYM as compared to inorganic fertilizer alone. Intensive farming with continuous application of fertilizer and FYM treatments results in increased root growth of crops that results in greater residues which may be the cause of this higher SOC content in soil. Consequently, the application of FYM significantly influenced the SOC content of soil [24].

**Table 2: pH, electrical conductivity and soil organic carbon under soybean-wheat cropping system in surface (0-15 cm) and subsurface (15-30 cm) under continuous fertilization for 50 years**

Treatments	pH		Electrical conductivity(dSm <sup>-1</sup> )		Soil organic carbon (%)	
	0-15 cm	15-30cm	0-15 cm	15-30cm	0-15 cm	15-30cm
50% NPK	7.42 <sup>b</sup> ±0.03 <sup>#</sup>	7.53 <sup>b</sup> ±0.02 <sup>#</sup>	0.16 <sup>c</sup> ±0.00	0.09 <sup>a</sup> ±0.01	0.55 <sup>b</sup> ±0.00	0.53 <sup>c</sup> ±0.03
100% NPK	7.48 <sup>c</sup> ±0.04	7.58 <sup>cd</sup> ±0.02	0.16 <sup>c</sup> ±0.00	0.12 <sup>b</sup> ±0.00	0.62 <sup>ab</sup> ±0.00	0.58 <sup>cde</sup> ±0.04
150% NPK	7.59 <sup>ef</sup> ±0.03	7.63 <sup>e</sup> ±0.02	0.16 <sup>bc</sup> ±0.00	0.08 <sup>a</sup> ±0.00	0.70 <sup>b</sup> ±0.10	0.65 <sup>e</sup> ±0.03
100% NPK+ Zn	7.61 <sup>f</sup> ±0.03	7.66 <sup>e</sup> ±0.03	0.18 <sup>e</sup> ±0.00	0.09 <sup>a</sup> ±0.00	0.60 <sup>ab</sup> ±0.02	0.58 <sup>cde</sup> ±0.08
100% NP	7.38 <sup>b</sup> ±0.04	7.42 <sup>b</sup> ±0.03	0.16 <sup>c</sup> ±0.00	0.08 <sup>a</sup> ±0.01	0.57 <sup>a</sup> ±0.08	0.54 <sup>ab</sup> ±0.04
100% N	7.31 <sup>a</sup> ±0.03	7.35 <sup>a</sup> ±0.03	0.16 <sup>a</sup> ±0.00	0.08 <sup>a</sup> ±0.01	0.56 <sup>a</sup> ±0.07	0.51 <sup>cd</sup> ±0.03
100% NPK+FYM	7.53 <sup>cde</sup> ±0.03	7.59 <sup>d</sup> ±0.03	0.18 <sup>f</sup> ±0.00	0.09 <sup>a</sup> ±0.02	0.79 <sup>b</sup> ±0.02	0.73 <sup>f</sup> ±0.03
100% NPK- S	7.55 <sup>de</sup> ±0.02	7.57 <sup>cd</sup> ±0.03	0.17 <sup>d</sup> ±0.00	0.09 <sup>a</sup> ±0.01	0.58 <sup>a</sup> ±0.02	0.55 <sup>de</sup> ±0.04
Control	7.51 <sup>cd</sup> ±0.02	7.55 <sup>cd</sup> ±0.02	0.15 <sup>b</sup> ±0.00	0.09 <sup>a</sup> ±0.01	0.43 <sup>a</sup> ±0.08	0.37 <sup>a</sup> ±0.03
<b>#Mean ± standard deviation</b>						
<b>*Means in a column followed by the same letter do not differ significantly (P&lt; 0.05) by Duncan's multiple range test</b>						

### 3.1.3. Available Nitrogen:

The available N in soil under various nutrient management practices ranged from 191.7 kg ha<sup>-1</sup> to 319.3 kg ha<sup>-1</sup> in surface soil and from 154 kg ha<sup>-1</sup> to 275 kg ha<sup>-1</sup> sub-surface layer (Table 3). The highest amount of available nitrogen was seen in case of application of 100% NPK+FYM while lowest was in case of control. Continuous application of integrated nutrients for 50 years resulted in remarkable increase in the available N content of soil as compared to the initial value (193.0 kg ha<sup>-1</sup>) while reduction of available N was seen in case of control. Application of inorganic fertilizers also led to accumulation of available N to soil as compare to the initial value. The available nitrogen content increased significantly with rise in fertilizer dose following the sequence 50% NPK < 100% N < 100% NPK < 150% NPK. While the rise in available N content with increase in fertilizer level is due to the obvious reasons, the direct addition of organic matter through FYM may have contributed to the increase in available N under NPK+FYM, facilitating in the growth of soil microorganisms and eventually improving the conversion of organically-bound N to mineral form [25]. Low level of available nitrogen in control or 50% NPK can be attributed to the low input of organic residues as well as fertilizers. Available N content in the subsurface soil layer (15-30 cm) had comparatively lesser value than the surface layer which is attributed to lesser organic matter content and lower microbial actions as compared to surface soil layer.

### 3.1. 4. Available Phosphorous:

Long term application of various nutrient management practices had significant impact on Available P in soil and it varied from 7.45 kg ha<sup>-1</sup> to 40.11 kg ha<sup>-1</sup> in surface soil and 6.49 kg ha<sup>-1</sup> to 37.85 kg ha<sup>-1</sup> in subsurface layer (Table 3). The lowest value was seen in case of control with no fertilization and highest in case of NPK+FYM. As compared to the initial value of available P in the soil ( 7.6 kg ha<sup>-1</sup>) there was significant increase in its amount due to application of fertilizers and manures over 50 years, with the maximum increase being witnessed in integrated nutrition. Available P was fairly low in treatment where there was application of only 100% N in both the layers and had value at par to control. The significant

variation in P content seen across different fertility treatments receiving suboptimal, optimum, and super-optimal nutritional dosages, respectively, indicates increased P build up. Among various treatments the highest soil P build-up was reported under NPK+ FYM, which was significantly more than 100% NPK. The annual addition of P and the solubilization of native P through increased release of organic acids in FYM treated plots are responsible for this build-up of P in the soil[26]. The reason behind the increased P accumulation in surface soil as compared to subsurface soil is attributed to the ability of soil to fix applied P and subsequently limit its mobility[27].

### 3.1.5 Available Potassium:

The ammonium acetate extracted K under different nutrient management practices after a period of 50 years ranged from 210 kg ha<sup>-1</sup> to 326 kg ha<sup>-1</sup> in surface soil under 100% N and 100% NPK + FYM respectively (Table 3). Similarly in the sub surface layers it varied from 206 kg ha<sup>-1</sup> to 302 kg ha<sup>-1</sup> in 100% N and 100% NPK+FYM respectively. It was lowest for treatments where K was not applied for years like 100% N followed by 100% NP because of mining of the inherent K present in soil [28]. Similar observation where the K content was in similar range and followed same pattern was reported by Khandagle et al. [20]. The K content declined with increase in depth of soil. The highest K found when FYM was added along with mineral fertilizer because FYM directly adds potassium in soil and also aids in release of K due to interaction of organic matter with clays [29].

**Table 3: Available nutrients (kg ha<sup>-1</sup>) in surface (0-15 cm) and subsurface (15-30 cm) after 50 years of continuous soybean-wheat cropping system**

Treatments	Available N(kg ha <sup>-1</sup> )		Available P(kg ha <sup>-1</sup> )		Available K(kg ha <sup>-1</sup> )	
	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
<b>50% NPK</b>	221.3 <sup>c</sup> ±1.52	193 <sup>bc</sup> ±4.72	21.78 <sup>b</sup> ±1.49	20.24 <sup>b</sup> ±1.55	256.00 <sup>c</sup> ±11.00	243.33 <sup>c</sup> ±5.50
<b>100% NPK</b>	282.0 <sup>f</sup> ±2.00	208 <sup>e</sup> ±2.00	33.60 <sup>d</sup> ±1.45	30.95 <sup>de</sup> ±1.51	286.70 <sup>d</sup> ±6.15	265.33 <sup>e</sup> ±5.13
<b>150% NPK</b>	309.3 <sup>g</sup> ±2.08	251 <sup>f</sup> ±3.51	36.98 <sup>e</sup> ±2.27	33.55 <sup>e</sup> ±1.15	321.00 <sup>e</sup> ±9.64	293.00 <sup>f</sup> ±6.00
<b>100%NPK+</b>	264.7 <sup>e</sup> ±4.66	205 <sup>e</sup> ±4.04	32.40 <sup>cd</sup> ±2.10	30.33 <sup>d</sup> ±1.93	287.33 <sup>d</sup> ±7.76	263.67 <sup>de</sup> ±7.09
<b>Zn</b>						
<b>100% NP</b>	251.7 <sup>d</sup> ±2.08	201 <sup>de</sup> ±4.72	29.07 <sup>c</sup> ±1.39	25.27 <sup>c</sup> ±2.23	218.33 <sup>ab</sup> ±6.80	210.56 <sup>a</sup> ±4.23
<b>100% N</b>	212.3 <sup>b</sup> ±2.08	189 <sup>b</sup> ±5.50	9.12 <sup>a</sup> ±1.60	7.85 <sup>a</sup> ±0.46	210.67 <sup>a</sup> ±4.16	206.00 <sup>a</sup> ±5.56
<b>100%</b>	319.3 <sup>h</sup> ±4.04	275 <sup>g</sup> ±5.68	40.11 <sup>e</sup> ±1.98	37.85 <sup>f</sup> ±3.10	326.33 <sup>e</sup> ±5.50	302.96 <sup>g</sup> ±2.75
<b>NPK+FYM</b>						
<b>100%NPK- S</b>	264.3 <sup>e</sup> ±6.02	198 <sup>cd</sup> ±2.51	31.15 <sup>cd</sup> ±2.61	29.07 <sup>d</sup> ±1.40	285.33 <sup>d</sup> ±3.21	255.67 <sup>d</sup> ±3.05
<b>Control</b>	191.7 <sup>a</sup> ±5.03	154 <sup>a</sup> ±3.05	7.45 <sup>a</sup> ±1.27	6.49 <sup>a</sup> ±0.46	225.67 <sup>b</sup> ±5.03	221.00 <sup>b</sup> ±5.00
<b>#Mean ± standard deviation</b>						
<b>*Means in a column followed by the same letter do not differ significantly (P&lt; 0.05) by Duncan's multiple range test</b>						

The candidate manuscript does not have a robust scientific discussion. I suggest the authors incorporate the suggested paragraphs, in this way it would improve the scientific quality of the manuscript.

The presented study explores the impact of integrated nutrient management on soil properties, specifically focusing on the long-term fertilizer experiment conducted in Vertisol under a wheat-soybean cropping system. The primary objective is to examine the influence of continuous fertilization on soil properties and soil carbon stock. The study is conducted at Jawahar Nehru Krishi Viswa Vidyalaya in Jabalpur, Madhya Pradesh, India, with soil samples collected in April 2022 following the wheat harvest.

Formatted: Font: Not Bold

Formatted: Justified

The results highlight that continuous application of 100% NPK (Nitrogen, Phosphorus, Potassium) along with farmyard manure (FYM) has a significant effect on all measured parameters. This treatment demonstrates the highest values for soil organic carbon (SOC), available nitrogen, phosphorus, and potassium at both soil depths (0-15 cm and 15-30 cm).

The conclusion drawn from the study emphasizes the positive impact of integrated application of nutrients, specifically 100% NPK along with FYM, on the accumulation of essential nutrients in the soil [30, 31, 32]. The findings suggest that this integrated nutrient management approach contributes significantly to the buildup of nutrients in the soil over the long term [33, 34]. The study implies that such integrated strategies are crucial for ensuring sustained soil productivity, providing insights into agricultural practices that can enhance crop production in the studied region [36, 37, 38].

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Comparatively, when considering studies on soil quality and the influence of agro-environmental factors on crop production in tropical territories of Latin America, one should note that regional variations in soil types [39, 40], climate [41, 42], and cropping systems may lead to different outcomes [43, 44, 45]. The scientific relevance of this study lies in its contribution to the understanding of nutrient management practices in Vertisol, which can inform agricultural strategies for improving soil fertility and crop yield in similar agroecological contexts [46, 47]. It adds valuable knowledge to the broader scientific discourse on sustainable agriculture and soil management in tropical regions [48, 49].

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

#### 4. CONCLUSION

It can be concluded from the current study that application of integrated nutrients for 50 years of continuous cultivation of wheat-soyabean in Vertisol had a significant and positive effect on soil carbon stocks as well as the available nutrients. While repeated application of sole chemical fertilizers also had similar impact on soil, their impact was far less as compared to the integrated nutrient application. Further imbalance application of chemical fertilizers had resulted in unwanted impacts on soil pH, mining of K, lower organic carbon etc. Thus application of integrated nutrient management is a safe and productive management system in the long run as compared to application of chemical fertilizer alone

#### REFERENCES

1. Salim, N., & Raza, A. (2020). Nutrient use efficiency (NUE) for sustainable wheat production: a review. *J. Plant Nutr.*, 43(2), 297-315.
2. Baweja, P., Kumar, S., & Kumar, G. (2020). Fertilizers and pesticides: Their impact on soil health and environment. *Soil health*, 265-285.
3. Bashir, M. T., Ali, S. A. L. M. I. A. T. O. N., Ghauri, M., Adris, A., & Harun, R. A. Z. I. F. (2013). Impact of excessive nitrogen fertilizers on the environment and associated mitigation strategies. *Asian J. Microbiol. Biotechnol. Environ. Sci*, 15(2), 213-221.
4. Jiao, W., Chen, W., Chang, A. C., & Page, A. L. (2012). Environmental risks of trace elements associated with long-term phosphate fertilizers applications: a review. *Environ. Pollut.*, 168, 44-53.
5. Huang, J., Xu, C. C., Ridoutt, B. G., Wang, X. C., & Ren, P. A. (2017). Nitrogen and phosphorus losses and eutrophication potential associated with fertilizer application to cropland in China. *J. Clean. Prod.*, 159, 171-179.

Formatted: English (United States)

Formatted: English (United States)

6. Saha, R., Mjshra, V. K., Majumdar, B., Laxminarayana, K., & Ghosh, P. K. (2010). Effect of integrated nutrient management on soil physical properties and crop productivity under a maize (*Zea mays*)–mustard (*Brassica campestris*) cropping sequence in acidic soils of Northeast India. *Commun Soil Sci Plant Anal.* 41(18), 2187-2200.
7. Nayak, A. K., Gangwar, B., Shukla, A. K., Mazumdar, S. P., Kumar, A., Raja, R., ... & Mohan, U. (2012). Long-term effect of different integrated nutrient management on soil organic carbon and its fractions and sustainability of rice–wheat system in Indo Gangetic Plains of India. *Field Crops Res.*, 127, 129-139.
8. Singh, A. K. (2007). Evaluation of soil quality under integrated nutrient management. *Indian Soc. Soil Sci.* 55(1), 58-61.
9. Gogoi, B. (2011). Soil properties and nutrients availability as affected by integrated nutrient management after rainfed cropping sequence. *Indian J. Agric. Res.* 45(4), 346-349.
10. Sharma, G. D., Thakur, R., Chouhan, N., & Keram, K. S. (2015). Effect of integrated nutrient management on yield, nutrient uptake, protein content, soil fertility and economic performance of rice (*Oryza sativa* L.) in a Vertisol. *Indian Soc. Soil Sci.* 63(3), 320-326.
11. Padbhushan, R., Sharma, S., Kumar, U., Rana, D. S., Kohli, A., Kaviraj, M., ... & Gupta, V. V. (2021). Meta-analysis approach to measure the effect of integrated nutrient management on crop performance, microbial activity, and carbon stocks in Indian soils. *Front. Environ. Sci.* 396.
12. Jackson, M. L. (1973). *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd., New Delhi.
13. Walkley, A. and Black, I. A. (1934). An examination of the segtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Sci.* 34: 29-38.
14. Subbiah, B. and Asija, G. L. (1956). Alkaline permanganate method of available nitrogen determination. *Current Sci.* 25: 259
15. Olsen, S. R., Cole, C. V., Watanable, F. S. and Dean, L. A. (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *USDA Circular* 939.
16. Gomez, K. A. and Gomez, A. A. (1984). *Statistical Procedures for Agricultural Research*. John Wiley & Sons, New Delhi
17. Suman, J., Dwivedi, B. S., Dwivedi, A. K., & Pande, S. K. (2018). effect of long term application of nitrogen and sulphur nutrition on nitrogen assimilation in Soybean. *J. Pharmacogn. Phytochem.*, 7(1), 2000-2004.
18. Hati, K. M., Swarup, A., Dwivedi, A. K., Misra, A. K., & Bandyopadhyay, K. K. (2007). Changes in soil physical properties and organic carbon status at the topsoil horizon of a vertisol of central india after 28 years of continuous cropping, fertilization and manuring. *Agric Ecosyst Environ*, 119(1-2), 127-134.
19. Chouhan, B., Rai, H. K., & Suryawanshi, A. (2017). Assessment of chemical properties of a Vertisol under long-term fertilizer experiment in soybean-wheat cropping system. *Int. J. Chem. Stud.*, 5(6), 2107-2111.
20. Khandagle, A., Dwivedi, B.S., Dwivedi, A.K., Panwar, S. and Thakur, R.K. (2020) Nitrogen fractions under long-term fertilizer and manure applications in soybean–wheat rotation in a Vertisol. *Indian Soc. Soil Sci.* 68, 186-193
21. Bahadur, L., Tiwari, D. D., Mjshra, J., & Gupta, B. R. (2012). Effect of integrated nutrient management on yield, microbial population and changes in soil properties under rice-wheat cropping system in sodic soil. *Indian Soc. Soil Sci.* 60(4), 326-329.
22. Bairwa, J., Dwivedi, B. S., Rawat, A., Thakur, R. K., & Mahawar, N. (2021). Long-term effect of nutrient management on soil microbial properties and nitrogen fixation in a vertisol under soybean-wheat cropping sequence. *Indian Soc. Soil Sci.* 69(2), 171-178.
23. Santhy P, Muthuvel P, Selvi D (2001). Status and impact of organic matter fractions on yield, uptake and available nutrients in a Long-Term Fertilizer Experiment. *Indian Soc. Soil Sci.* 49:281-28

Formatted: English (United States), All caps

Formatted: English (United States)

Formatted: English (United States), All caps

Formatted: English (United States)

Formatted: English (United States), All caps

Formatted: English (United States)

Formatted: English (United States), All caps

Formatted: English (United States)

Formatted: English (United States), All caps

Formatted: English (United States)

Formatted: English (United States), All caps

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

24. Meshram, M. K., Dwivedi, B. S., Naik, K. R., Thakur, R., & Keram, K. S. YIELD, NUTRIENT UPTAKE, SOIL FERTILITY AND ECONOMIC.
25. Singh, Muneshwar, Wanjari, R.H., Dwivedi, Anil and Dalal, Ram (2012) Yield response to applied nutrients and estimates of N<sub>2</sub> fixation in 33 year old soybean–wheat experiment on a Vertisol. *Experimental Agriculture* 48, 311-325.
26. Thakur, Risikesh, Sawarkar, S.D., Vaishya, U.K. and Singh, Muneshwar (2011) Impact of continuous use of inorganic fertilizers and organic manure on soil properties and productivity under soybean-wheat intensive cropping of a Vertisol. *Indian Soc. Soil Sci*, 59, 74-8
27. Dubey, L., Dwivedi, A. K., & Dubey, M. (2015). Long term application of fertilizer and manures on physico-chemical properties and N, P and K uptake in soybean-wheat cropping system. *Plant Archives*, 15(1), 143-147.
28. Sawarkar, S.D., Khamparia, N.K., Thakur, Risikesh, Dewda, M.S. and Singh, Muneshwar (2013) Effect of longterm application of inorganic fertilizers and organic manure on yield, potassium uptake and profile distribution of potassium fractions in Vertisol under soybean-wheat cropping system *Indian Soc. Soil Sci*, 61, 9
29. Urkurkar, J.S., Tewari, S., Bajpai, R.K. (2010). Influence of long term use of organic and inorganic manures on soil fertility and sustainable productivity of rice (*Oryza sativa*) and wheat (*Triticum aestivum*) in Inceptisols. *Indian J. Agric. Sci.* 80 (3), 208-212.
30. Araya-Alman, M., Olivares, B., Acevedo-Opazo, C. et al. (2020). [Relationship Between Soil Properties and Banana Productivity in the Two Main Cultivation Areas in Venezuela. \*J Soil Sci Plant Nutr.\*: 20 \(3\): 2512-2524. <https://doi.org/10.1007/s42729-020-00317-8>](https://doi.org/10.1007/s42729-020-00317-8)
31. Lobo, D; Olivares, B; Rey, J.C; Vega, A; Rueda-Calderón, A. [Relationships between the Visual Evaluation of Soil Structure \(VESS\) and soil properties in agriculture: A meta-analysis. \*Scientia agropecuaria\*, 2023; 14 - 1, 67 - 78. <https://doi.org/10.17268/sci.agropecu.2023.007>](https://doi.org/10.17268/sci.agropecu.2023.007)
32. Olivares, B.O., Calero, J., Rey, J.C., Lobo, D., Landa, B.B., Gómez, J. A. (2022a). Correlation of banana productivity levels and soil morphological properties using regularized optimal scaling regression. *Catena*, 208: 105718. <https://doi.org/10.1016/j.catena.2021.105718>
33. Olivares, B. (2016). Description of soil management in agricultural production systems in the Hamaca de Anzoátegui sector, Venezuela. *La Granja: Revista de Ciencias de la Vida*. 23(1): 14–24. <https://n9.cl/yep08>
34. Olivares, B., Verbist, K., Lobo, D., Vargas, R. y Silva, O. (2011). Evaluation of the USLE model to estimate water erosion in an Alfisol. *Journal of Soil Science and Plant Nutrition of Chile*. 11 (2):71-84. <http://dx.doi.org/10.4067/S0718-95162011000200007>
35. Olivares, B.O.; Rey, J.C.; Perichi, G.; Lobo, D. (2022). Relationship of Microbial Activity with Soil Properties in Banana Plantations in Venezuela. *Sustainability* 14, 13531. <https://doi.org/10.3390/su142013531>
36. Campos, B. O. (2023). *Banana Production in Venezuela: Novel Solutions to Productivity and Plant Health*. Springer Nature. <https://doi.org/10.1007/978-3-031-34475-6>
37. Olivares B, Rey JC, Lobo D, Navas-Cortés JA, Gómez JA, Landa BB. (2022). Machine Learning and the New Sustainable Agriculture: Applications in Banana Production Systems of Venezuela. *Agricultural Research Updates*. 42, 133 - 157. Nova Science Publishers, Inc

Formatted: English (United States)

38. Rodríguez, M.F., Olivares, B., Cortez, A., Rey, J.C. and Lobo, D. (2015). Natural physical characterization of the indigenous community of Kashaama for the purposes of sustainable land management. *Acta Nova*. 7 (2):143-164. <https://n9.cl/hakdx>
39. Olivares, B. O., & Franco, E. (2015). Diagnóstico agrosocial de la comunidad indígena de Kashaama: Un estudio empírico en el estado de Anzoátegui, Venezuela. *Revista Guillermo de Ockham*, 13(1), 87-95.
40. Hernández, R., Olivares, B. (2020). Application of multivariate techniques in the agricultural land's aptitude in Carabobo, Venezuela. *Tropical and Subtropical Agroecosystems*, 23(2):1-12. <https://n9.cl/zeedh>
41. Hernández, R.; Olivares, B. Arias, A; Molina, J.C., Pereira, Y. (2018). Agroclimatic zoning of corn crop for sustainable agricultural production in Carabobo, Venezuela. *Revista Universitaria de Geografía*.. 27 (2): 139-159. <https://n9.cl/l2m83>
42. Hernandez, R., Olivares, B., Arias, A, Molina, J.C., Pereira, Y. (2020). Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. *Idesia*. 38(2):95-102. <http://dx.doi.org/10.4067/S071834292020000200095>
43. Hernández, R; Olivares, B., Arias, A; Molina, J.C., Pereira, Y. (2018). Identification of potential agroclimatic zones for the production of onion (*Allium cepa* L.) in Carabobo, Venezuela. *Journal of the Selva Andina Biosphere*., 6 (2): 70-82. [http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2\\_a03.pdf](http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2_a03.pdf)
44. Hernández, R. Olivares, B. (2019). Ecoterritorial sectorization for the sustainable agricultural production of potato (*Solanum tuberosum* L.) in Carabobo, Venezuela. *Agricultural Science and Technology*. 20(2): 339-354. [https://doi.org/10.21930/rcta.vol20\\_num2\\_art:1462](https://doi.org/10.21930/rcta.vol20_num2_art:1462)
45. Olivares, B., Hernández, R; Coelho, R., Molina, J.C., Pereira, Y. (2018). Spatial analysis of the water index: advances in sustainable decision-making in Carabobo agricultural territories, Venezuela *Revista Geográfica de América Central*. 60 (1): 277-299. DOI: <https://doi.org/10.15359/rgac.60-1.10>
46. Rodríguez, M.F., Cortez, A., Olivares, B., Rey, J.C, Parra, R., Lobo, D (2013). Spatio-temporal analysis of precipitation in the state of Anzoátegui and its surroundings. *Agronomía Tropical* 63 (1-2): 57-65. <https://n9.cl/14iow>
47. Olivares, B., Parra, R., Cortez, A. y Rodríguez, M.F. (2012). Patterns of rainfall homogeneity in climatic stations of the Anzoátegui state, Venezuela. *Revista Multiciencias*. 12 (Extraordinario): 11-17. <https://n9.cl/xbslq>
48. Montenegro, E., Pitti-Rodríguez, J, Olivares-Campos, B. 2021. Identification of the main subsistence crops of Teribe: a case study based on multivariate techniques. *Idesia (Arica)*.; 39(3), 83-94. <https://dx.doi.org/10.4067/S0718-34292021000300083>

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

49. Vilorio, J; Olivares, B. et al. 2023. Mapping Projected Variations of Temperature and Precipitation Due to Climate Change in Venezuela. *Hydrology*, 10, 96. <https://doi.org/10.3390/hydrology10040096>

29.

Formatted: Normal, Indent: Left: 0.25", No bullets or numbering

UNDER PEER REVIEW