

Effect of chitosan on increasing beneficial soil microflora and disease suppression in turmeric plants in humid tropics of Kerala, India

ABSTRACT

The present study assessed the population of nitrogen fixers and phosphate solubilizers in the rhizosphere, and suppression of leaf blotch disease caused by *Taphrinamaculans* (Order: Taphrinales, Family: Taphrinaceae) in turmeric plants in response to chitosan application. Leaf blotch disease caused by *Taphrinamaculans* has recorded up to 37.60 % to 52.90 % yield loss in turmeric. Hence, control measures have to be adopted. Chitosan is a biopolymer derived from chitin, which could evoke defense responses in plants. The treatments for this study included, foliar spray of chitosan 2g L^{-1} at monthly intervals (F_1), foliar spray of chitosan 4g L^{-1} at bimonthly intervals (F_2), soil drenching of chitosan 2g L^{-1} at monthly interval (S_1) and soil drenching of chitosan 4g L^{-1} at bimonthly interval (S_2). The untreated plants served as the control. The population of beneficial soil microbes were assessed and soil application of chitosan was observed to have better effect over foliar application with respect to the population of nitrogen fixing bacteria. More population of nitrogen fixing bacteria was recorded in the soil where chitosan 4g L^{-1} was applied as soil drenching at bimonthly interval (S_2). The population of phosphate solubilizers were significantly higher over the control in all the chitosan treatments irrespective of the mode of application. However, the higher population of phosphate solubilizers were observed, when chitosan 2g L^{-1} was given as foliar spray at monthly intervals (F_1). The chitosan treated plants were also observed to have significantly lower incidence of leaf blotch disease compared to control plants. The use of chitosan as a plant biostimulant protects the plant from soil borne pathogens and increases the population of beneficial soil microflora, which have a role in improving the nutrient uptake by plants thereby enhancing the plant growth and yield.

Keywords: Turmeric, Chitosan, Nitrogen fixing bacteria, Phosphate solubilizers, Leaf blotch

1. INTRODUCTION

Soil amendments, either organic or inorganic when added to the soil improve its physiochemical and biological properties. It provides a favourable environment for plant growth by improving the soil structure, soil microflora, water holding capacity and availability of nutrients. Organic amendments are a rich source of organic matter and nutrients. It facilitates the growth and activity of soil microorganisms. Increased activity of soil microbes promotes efficient nutrient cycling and suppress soil-borne pathogens. Soil microbes decompose organic matter, during which they produce enzymes and metabolites that are capable of breaking down salts and other harmful compounds, thus improving soil health [1].

According to Deng et al. [2], organic amendments could increase rhizosphere microbial diversity by providing additional nutrient sources. Organic amendments could facilitate beneficial symbiotic relationship between plants and soil microflora, such as mycorrhizal fungi and nitrogen-fixing bacteria. Alori et al. [3] observed that organic amendments like compost, poultry manure and green manure from *Tithonia diversifolia* improved number and diversity of fungi and bacteria. The plant growth seems to be the outcome of better soil physical properties viz., pore size distribution, particle size, porosity, water-holding capacity and enhanced populations of soil microbes (fungi, bacteria, actinomycetes).

Chitosan is a plant biostimulant that evokes plant growth promotion, soil conditioning, metabolite elicitation and defense responses in plants. Chitosan, extracted from exoskeletons of crustaceans, certain insects and fungal cell wall is a natural, safe and cost effective biopolymer obtained from the deacetylation of chitin, a long chain polymer of N-acetyl-glucosamine [4]. Chitosan could be applied to plants as seed priming, soil drench or as foliar spray. Chitosan when applied to soil has shown great effect on plant growth due to its ability to improve soil fertility and enhancement of mineral nutrients absorption by plant [5]. Due to the presence of nitrogen (~8.9%–9.5%) in its amino group, chitosan serves as a source of nitrogen to the plants. Also, when applied to the plant, it increases the mineral nutrients uptake of nitrogen, phosphorous, potassium, calcium and magnesium [6].

Sharp [7] reported that the addition of chitosan alters rhizosphere conditions to shift the microbial balance in favour of beneficial organisms and detrimental to plant pathogens. The primary function of these beneficial bacteria is to decompose organic matter, provide nutrients to crops, improve soil structure etc. Healthy populations of beneficial bacteria can help suppress pathogens and promote plant growth and increase the yield[8]. In a study conducted by Dzung *et al.* [9], it was found that chitosan spray resulted in improved nutrient uptake. The synergetic effects of many factors, such as suppression of plant diseases, insects, and nematodes, increased biomass and activities of beneficial microbes, improved physical structure of soil and nutrient availability and direct plant growth stimulation makes chitosan as a good soil amendment [10].

Chitosan has been reported to have a positive effect on rhizobacterial growth, where it possesses a symbiotic relation with growth promoting rhizobacteria, thus triggered germination rate and improving plant nutrient uptake [11]. Nitrogen fixers and phosphate solubilizers are important groups of plant growth promoting rhizobacteria having unique role in agriculture. *Azotobacterchroococcum* and *Azospirillumbrasile*, are two free-living aerobic nitrogen fixing bacteria that can be found in most soil and have the ability to convert inert N₂ into available forms for plants. *Bacillus megaterium* and *Pseudomonas fluorescens* are notable for the ability to solubilize unavailable phosphates in soil, as well as produce a wide variety of metabolites like auxin[12].

Turmeric is an important spice cum medicinal plant. The major principle component, curcumin is responsible for therapeutic value of the plant. India is the largest producer, consumer and exporter of turmeric in the world. The annual production of turmeric in India is around 11.61 lakh tonnes per annum. India's contribute to more than 75 % share in world production. Turmeric is grown in more than 20 states in the country. The major states producing turmeric in India are Telangana, Maharashtra, Karnataka, Tamil Nadu, Andhra Pradesh and Madhya Pradesh. India has more than 62 % share of world trade in turmeric [13]. The turmeric is affected by major foliar disease leaf blotch caused by the pathogen *Taphrinamaculans*. Yield loss upto 37.60 % to 52.90 % has been reported in turmeric due to this disease [14]. Hence control measures, specifically biological measures need to be adopted to manage this disease. In this context, the present study was undertaken to assess the population of beneficial soil microflora (nitrogen fixing and phosphate solubilising bacteria) in the rhizosphere of turmeric plants in response to chitosan, when applied both as soil amendment and foliar spray.

2. MATERIALS AND METHODS

Turmeric plants were grown in farmer's field in Trivandrum district, Kerala during June 2021 to January 2022. The varieties, Sobha and Sona released from Kerala Agricultural University were used for the study. The 45 day old protray plantlets were transplanted to the main field at a spacing of 25cm x 25cm in plots of size 3m x 1m and the field was laid out in Randomized block design with four replicates. Crop was raised organically as per Package of Practices recommendations (Organic) [15]. Chitosan (from prawn shell waste) was procured from Matsyafed Chitin and Chitosan Plant, Neendakara, Kollam, Kerala. The chitosan solutions was prepared by dissolving it in aqueous solution of 1 per cent acetic acid. The treatments included, foliar spray of chitosan 2gL⁻¹ at monthly intervals (F₁), foliar spray of chitosan 4gL⁻¹ at bimonthly intervals (F₂), soil drenching of chitosan 2gL⁻¹ at monthly interval (S₁) and soil drenching of chitosan 4gL⁻¹ at bimonthly interval (S₂). The plants that were not exposed to either mode of application of chitosan served as the control. The chitosan treatments were given till 5 months after transplanting (MAT) at specified intervals. Plants were harvested at 7 months after transplanting. Soil samples were collected from the rhizosphere of turmeric plants in sterile polythene bags at the time of harvest. The population assessment of free living nitrogen fixing microorganisms and phosphate solubilizers of rhizosphere soil, collected from the field was carried out by serial dilution followed by plating [16]. The lab experiment was laid out in Completely Randomized Design (CRD) with three replicates. Nitrogen fixing bacteria from the soil samples were counted after growing in different N-free agar media such as N-free malate bromothymol blue (NFB) and Jensen's media. The isolates of phosphate solubilizing bacteria were counted on Pikovskaya's agar. The population of beneficial soil microflora was expressed in log cfu g⁻¹ soil from three replicates. Crop was monitored throughout the growth period and incidence of leaf blotch disease, and disease intensity was assessed by using 0 to 6 disease rating scale, where 0 = Disease free, 1 = 0.10-10.00%, 2 = 10.10-25.00%, 3 = 25.10-50.00%, 4 = 50.10-75.00%, 5 = >75% and 6 = 100% [17]. The data were subjected to analysis of variance using package KAU GRAPES [18].

3. RESULT AND DISCUSSION

3.1 Effect of chitosan on beneficial soil microflora

The nitrogen fixing bacteria could be seen as sugar like crystals on the media. Significant difference was noticed in the population of nitrogen fixing bacteria among the treatments (Table 1). The growth of nitrogen fixing bacteria showed similar trend in both the media. In the variety Sobha, nitrogen fixing bacteria was observed to be the highest in S₂ (soil drenching with chitosan 4g L⁻¹ at bimonthly interval) in both NFB (5.02 log cfu g⁻¹) and Jensen's media (4.89 log cfu g⁻¹). In the variety Sona also same treatment recorded the highest number of colony forming units in NFB (5.12 log cfu g⁻¹) and Jensen's media (5.12 log cfu g⁻¹). The lowest population of nitrogen fixing bacteria was observed in control treatment in both the varieties.

Table 1. Population of nitrogen fixing bacteria in response to chitosan application

Treatments	Population of nitrogen fixing bacteria (log cfu g ⁻¹)			
	NFB medium		Jensen's medium	
	Sobha	Sona	Sobha	Sona
F ₁ : 2g L ⁻¹ monthly foliar application	4.24 ± 0.06 ^c	4.32 ± 0.13 ^c	4.41 ± 0.19 ^b	4.45 ± 0.17 ^b
F ₂ : 4g L ⁻¹ bimonthly foliar application	4.07 ± 0.11 ^d	4.21 ± 0.24 ^c	4.36 ± 0.16 ^b	4.40 ± 0.13 ^b
S ₁ : 2g L ⁻¹ monthly soil application	4.63 ± 0.06 ^b	4.67 ± 0.10 ^b	4.19 ± 0.17 ^b	4.56 ± 0.28 ^b
S ₂ : 4g L ⁻¹ bimonthly soil application	5.02 ± 0.05 ^a	5.12 ± 0.07 ^a	4.89 ± 0.06 ^a	5.12 ± 0.10 ^a
Control (with no chitosan treatment)	4.00 ± 0.07 ^d	4.19 ± 0.10 ^c	4.14 ± 0.16 ^b	4.28 ± 0.16 ^b
SEm (±)	0.043	0.081	0.09	0.103
CD (0.05)	0.135	0.254	0.285	0.323
CV	1.69	3.10	3.564	3.891

The values are mean of 3replications. Values that are followed by the same letter in a column do not differ significantly between them according to LSD

According to Abd-El-Kareem *et al.* [19], soil application of chitosan (0, 2, 4 and 6 g per kg of soil) significantly improved the total count of chitinolytic bacteria at all concentrations, and the *Bacillus* spp., one of the most common nitrogen fixing bacteria was the most noticed genus among the isolated bacteria.

Soil application of chitosan was observed to have better effect over foliar application with respect to the population of nitrogen fixing bacteria. When the same concentration of chitosan was applied as foliar spray and soil drenching, the latter was found to be more effective in improving the microbial population. The multiplication and activities of soil bacterial population would have more when chitosan was directly applied to the soil rather than when applied on the leaves. Chitosan when applied to soil could alter the soil properties in favour of microbial growth and multiplication, in consensus with the findings of Xu and Mou [10], who reported that chitosan could provide a carbon source for microbes in the soil.

Nitrogen is the most essential and restrictive nutrient for plant growth. The unavailable atmospheric nitrogen converts to plant available form as ammonia through nitrogen fixation by a group of nitrogen fixing bacteria in the soil. Nitrogen fixing is essential for plant nutrition by increasing N uptake by the plants and playing a significant role as plant growth promotion [20]. The improvement in the count of nitrogen fixing bacteria by chitosan application in our study indirectly would indicate its effect on plant growth promotion.

The population of phosphate solubilizers also showed significant difference among the treatments tried (Table 2). In Sobha, the highest number of colony forming units of phosphate solubilizing bacteria (3.32 log cfu g⁻¹) were recorded in F₁ and this was on par with all other treatments

except control. In Sona, the highest value ($3.42 \log \text{cfu g}^{-1}$) was recorded in F_1 and on par with F_2 and S_2 . The control treatment recorded a significantly lower value for phosphate solubilizing bacteria in both the varieties.

Table 2. Population of phosphate solubilizing bacteria in response to chitosan application

Treatments	Population of phosphate solubilizing bacteria ($\log \text{cfu g}^{-1}$)	
	Sobha	Sona
F_1 : 2 g L^{-1} monthly foliar application	3.32 ± 0.28^a	3.42 ± 0.10^a
F_2 : 4 g L^{-1} bimonthly foliar application	3.25 ± 0.24^a	3.36 ± 0.10^a
S_1 : 2 g L^{-1} monthly soil application	3.00 ± 0.00^a	3.10 ± 0.17^b
S_2 : 4 g L^{-1} bimonthly soil application	3.10 ± 0.17^a	3.30 ± 0.00^{ab}
Control (with no chitosan treatment)	2.00 ± 0.00^b	2.10 ± 0.17^c
SEm (\pm)	0.105	0.074
CD (0.05)	0.33	0.232
CV	6.175	4.167

The values are mean of 3 replications that are followed by the same letter do not differ significantly between them according to LSD.

Burkholderia gladioli, a phosphate-solubilizing bacteria considerably improved its phosphate-solubilizing capacity when cultured in fermented chitosan medium, which further increased the amount of available phosphorus when amended in the soil [21]. Amerany, et al [22] in his study chitosan 1 mg per plant was applied to the soil in the transplant cavity after three weeks of transplanting of arbuscular mycorrhizal fungi (AMF) inoculated tomato (*Solanum lycopersicum*) seedlings and AMF colonization frequency and intensity reached to higher values after 12 weeks of growth.

Phosphorus is the second most essential nutrient element for plants growth. Phosphorus plays an important role in photosynthesis, development of good root system, respiration, energy storage and transfer etc. The improved count of phosphate solubilizers by chitosan application helps in enhanced phosphorus uptake by plants and thus leads to better plant development. In the present study the population of phosphate solubilizers were found to be significantly higher over the control in all the chitosan treatments irrespective of foliar and soil application in both the varieties.

Kerala Agriculture University recommends the application of *Azospirillum* and phosphate solubilizing bacteria ($20 \text{ g/ bed of } 3\text{m}^2$) as nutrient supplement in turmeric along with chemical fertilizers [23]. From the findings of our study it can be said that chitosan can be used as an alternative to meet this recommendation. In our experiment we found that chitosan application enhanced the count of nitrogen fixing and phosphate solubilizing bacteria in the root zone. These two beneficial soil microbes are actively involved in facilitating plant's nutrient uptake. Thus the use of chitosan as plant biostimulant can supply the nutrients to the plants as well as improve the population of beneficial microflora in soil.

3.2 Effect of chitosan on disease incidence

The incidence of leaf blotch disease caused by *Taphrinamaculans* (Order: Taphrinales, Family: Taphrinaceae) was noticed in the field at 6 MAT and significant variation was observed with respect to per cent disease index among the treatments in both the varieties (Table 3). Significantly lower disease index was recorded in F_1 (12.78 %) in Sobha and F_2 (14.44 %) in Sona and found to be on par with all other chitosan treatments. The highest per cent disease index was noticed in control plot in both the varieties. Leaf blotch disease caused by *Taphrinamaculans* is a serious disease of the

turmeric leaves causing a significant decrease in yield due to loss of photosynthetic properties. The disease is soil and seed borne and the pathogen survive in soil on infected plant debris.

Table 3. Per cent incidence of leaf blotch disease in response to chitosan application

Treatments	Per cent disease index	
	Sobha	Sona
F ₁ : 2 g L ⁻¹ monthly foliar application	12.78 ± 2.84 ^b	15.00 ± 1.36 ^b
F ₂ : 4 g L ⁻¹ bimonthly foliar application	15.56 ± 3.42 ^b	14.44 ± 1.57 ^b
S ₁ : 2 g L ⁻¹ monthly soil application	16.11 ± 2.08 ^b	15.55 ± 3.14 ^b
S ₂ : 4 g L ⁻¹ bimonthly soil application	13.33 ± 1.36 ^b	17.78 ± 2.08 ^b
Control (with no chitosan treatment)	46.11 ± 2.08 ^a	48.34 ± 1.36 ^a
SEm (±)	1.091	1.125
CD (0.05)	3.361	3.468
CV	10.5	10.129

The values are mean of 4 replications that are followed by the same letter do not differ significantly between them according to Least Significance Difference (LSD).

Anusuya and Sathiyabama [24] reported that turmeric plant treated with chitosan (0.1% w/v spraying on leaves) showed increased resistance towards rhizome rot disease caused by *Pythium aphanidermatum*. In a study conducted to determine the effects of chitosan on *Colletotrichum* anthracnose on papaya fruit, more than 60% control of anthracnose was reported when chitosan is applied before pathogen inoculation [25]. Liu et al. [26] demonstrated that chitosan protects rice seedling from sheath blight caused by *Rhizoctonia solani* and causes a 66-91% inhibition of lesion length.

Chitosan is a plant protectant that helps to reduce disease incidence and severity in crops by inhibiting microbial growth and decreasing microbial membrane integrity [10]. Chitosan was also found to have effect on disease suppression by the way of enhancing systemic resistance of the plant. The mechanism of action of chitosan is through direct toxicity or chelation of nutrients and minerals from the pathogen. The biopolymer properties of this compound also helps to form physical barriers around the penetration sites of pathogens thus, preventing them from spreading to healthy tissues. Chitosan induces local and systemic reactions that involve in signalling cascades and activation of defense related antimicrobial compounds and proteins [27].

In the present study, the application of chitosan was found to be effective in reducing the incidence of leaf blotch disease. The foliar spray and soil application of chitosan may help to reduce the population of harmful pathogens in the soil and thus protects the plant from disease incidence. The results of our study is also in consonance with the findings of Sharp [7], who reported that the addition of chitosan alters rhizosphere conditions to shift the microbial balance in favour of beneficial organisms and detrimental to plant pathogens.

4. CONCLUSION

Chitosan, is a natural, safe and cost effective biopolymer extracted from exoskeletons of crustaceans, certain insects and fungal cell wall. The addition of chitosan alters rhizosphere conditions to shift the microbial balance in favour of beneficial organisms and detrimental to plant pathogens. In the present study, the use of chitosan as foliar spray or soil drenching, increased the population of beneficial soil microflora viz. nitrogen fixing bacteria and phosphate solubilizers. These microbes could play a major role in plant's nutrient uptake, which could reflect on plant growth and yield. Chitosan application was also found to be effective in controlling the leaf blotch disease caused by *Taphrinamaculans*. Thus, the use of chitosan is an organic practice of enhancing soil nutrient status

and biological disease management, which is an alternative to chemical use and hence, safe to environment and human health. Chitosan is, thus a safe alternative to agrochemicals for plant disease management and improvement of soil health.

REFERENCE

1. Urra J, Alkorta I, Garbisu C. Potential benefits and risks for soil health derived from the use of organic amendments in agriculture. *Agronomy*. 2019;9(9): 542.
2. Deng S, Wipf HML, Pierroz G, Raab TK, Khanna R, Coleman-Derr D. A plant growth-promoting microbial soil amendment dynamically alters the strawberry root bacterial microbiome. *Sci. Rep.* 2019; 9: 17677.
3. Alori ET, Olaniyan FT, Adekiya AO, Ayorinde BB, Daramola FY, Joseph A, et al. Response of soil microbial community (bacteria and fungi) to organic and inorganic amendments using tomato as a test crop. *Air Soil Water Res.* 2023; 16: 1–13.
4. Domard A, Domard M. Chitosan: structure - properties relationship and biomedical applications. *Polym. Biomater.* 2002. 10: 187–212.
5. Berger L.R., Stamford N.P., Santos, C.E.R.S., Freitas, A.D.S., Franco, L.O. and Stamford, T.C.M. 2013. Plant and soil characteristics affected by biofertilizers from rocks and organic matter inoculated with diazotrophic bacteria and fungi that produce chitosan. *J. Soil Sci. Plant Nutr.* 2013, 13(3), 592-603.
6. Sharma G, Prajapati D, Devi KA, Pal A, Saharan V. Chitosan nanomaterials for delivery of micronutrients in plants. *Role of Chitosan and Chitosan-Based Nanomaterials in Plant Sciences*. Ed: Kumar S, Madhally SV. Academic Press. 2022. pp. 239-253.
7. Sharp RG. A review of the applications of chitin and its derivatives in agriculture to modify plant-microbial interactions and improve crop yields. *Agronomy*. 2013; 3: 757–793.
8. Mishra J, Prakash J, Arora NK. Role of beneficial soil microbes in sustainable agriculture and environmental management. *Clim. chang. environ. sustain.* 2016;4(2): 137-149.
9. Dzung NA, Khanh VTP, Dzung TT. Research on impact of chitosan oligomers on biophysical characteristics, growth, development and drought resistance of coffee. *Carbohydr. Polym.* 2011; 84: 751-755.
10. Xu C, Mou B. Chitosan as soil amendment affects lettuce growth, photochemical efficiency, and gas exchange. *HortTechnology*, 2018 28(4): 476-480.
11. Sharif R, Mujtaba M, Rahman MU, Shalmani A, Ahmad H, Anwar T et al. The multifunctional role of chitosan in horticultural crops; a review. *Molecules*. 2018.23: 872.
12. Wang Z, Chen Z, Xu Z, Fu X. Effects of Phosphate-Solubilizing Bacteria and N₂-fixing Bacteria on Nutrient Uptake, Plant Growth, and Bioactive Compound Accumulation in *Cyclocaryapaliurus* (Batal.) Ijinskaja. *Forests*. 2019. 10: 772
13. PIB [Press Information Bureau] Government of India notifies establishment of National Turmeric Board [online]. 2023. Available: <https://pib.gov.in/PressReleasePage.aspx?PRID=1964083> [04 October 2023].
14. Kumar M, Mishra AK, Kumar P, Kumar B, Maurya S. Capability of leaf blotch sickness of turmeric (*Taphrinamaculans*) at the phenology of crop and its yield attributing characters. *Int. J. Curr. Microbiol. Appl. Sci.* 2020. 10:424-429.
15. KAU [Kerala Agricultural University] Package of Practices Recommendations (Organic): Crops 2017 (2nd Ed.). Kerala Agricultural University, Thrissur. 2017. 328p.
16. Reynolds J. Serial dilution protocols. *American Society for Microbiology*: Washington, DC, USA. 2005 Sep 30:1-7.
17. Nambiar KKN, Sarama YR, Brahma RN. Field reaction of turmeric types (*Curcuma longa*, *Curcuma aromatica*) to leaf blotch disease (caused by *Taphrinamaculans*). *J. Plant. Crops*. 1977; 5: 124.
18. Gopinath PP, Parsad R, Joseph B, Adarsh VS. GRAPES: General Rshiny Based Analysis Platform Empowered by Statistics. 2020. <https://www.kaugrapes.com/home>. version 1.0.0. DOI: 10.5281/zenodo.4923220
19. Abd-El-Kareem F, El-Mougy NS, El-Gamal NG, Fotouh YO. Use of chitin and chitosan against tomato root rot disease under greenhouse conditions. *Res. J. Agric. Biol. Sci.* 2006.2(4): 147-152.
20. Abo-Koura HA, Bishara MM, Saad MM. Isolation and Identification of N - Fixing, Phosphate and 2 Potassium Solubilizing Rhizobacteria and Their Effect on Root Colonization of Wheat Plant. *Int. J. Microbiol. Res.* 2019. 10 (2): 62-76.
21. Sun, X., Shao, C., Chen, L., Jin, X. and Ni, H. 2021. Plant growth-promoting effect of the chitosanolytic phosphate-solubilizing bacterium *Burkholderia gladioli* MEL01 after

- fermentation with chitosan and fertilization with rock phosphate. *J. Plant Growth Regul.* 40: 1674-1686.
22. Ameranya F, Rhazia M, Wahbic S, Taourirteb, M, Meddichc A. The effect of chitosan, arbuscular mycorrhizal fungi, and compost applied individually or in combination on growth, nutrient uptake, and stem anatomy of tomato. *Scientia Horticulturae.* 2019. <https://doi.org/10.1016/j.scienta.2019.109015>.
 23. KAU [Kerala Agriculture University]. Package of Practices Recommendations: Crops 2016 (15th Ed.). Kerala Agricultural University, Thrissur.2016. 392p.
 24. Anusuya, S. and Sathiyabama, M. 2016. Effect of chitosan on growth, yield and curcumin content in turmeric under field condition. *Biocatalysis Agric. Biotechnol.* 6: 102-106.
 25. Bautista-Banos, S., M. Hernandez-Lopez, E. Bosquez-Molina and C.L. Wilson, 2003. Effects of chitosan and plant extracts on growth of *Colletotrichum gloeosporioides*, anthracnose levels and quality of papaya fruit. *Crop Protect.* 22: 1087-1092.
 26. Liu, H., W. Tian, B. Li, G. Wu and M. Ibrahim et al., 2012. Antifungal effect and mechanism of chitosan against the rice sheath blight pathogen, *Rhizoctonia solani*. *Biotechnol. Lett.*, 34: 2291-2298.
 27. Hassan O, Chang T. Chitosan for eco-friendly control of plant disease. *Asian J. Plant Pathol.* 2017;11: 53-70.

UNDER PEER REVIEW