

## **Role of foliar application of Micronutrients on growth and yield of Pulses - A review**

### **Abstract**

Although pulses respond strongly to micronutrients, particularly during the reproductive stages, their application is still uncommon. The results of the current study also suggest that foliar Zn, Fe or Mo application had a favourable impact on growth indicators and grain yield, protein content, total and active nodule counts, and yield components plant<sup>-1</sup>. Molybdenum is essential for the development of nodule tissue and an increase in N<sub>2</sub> fixation, both of which are necessary for Rhizobium to increase the nitrogen fixation process.

**Key words:** *Iron, Molybdenum, Pulses, Yield, Zinc*

### **Introduction**

Pulses are the main source of protein particularly for vegetarians and contribute about 14 per cent of the total protein of an Indian average diet (Meena *et al.*, 2013). The production and consumption of pulses, sometimes referred to as food legumes, is second only to cereals in India. The word pulse is derived from the Latin word Puls means pottage i.e., seed boiled to make porridge or thick soup. Pulse is a rich source of dietary protein (25%), energy, minerals and vitamins for the vegetarian population. As a vital source of both human and animal food and animal feed, it is also crucial for maintaining soil fertility (Kavya *et al.*, 2021).

In addition to crop productivity, micronutrients also improve the amount of crude protein, amino acids, total lipids, energy value, and other nutrients in chickpea, soybean, black gramme, etc. Foliar fertilisation is the most efficient way to raise output, and research has shown that it can increase yield by 12 to 25% when compared to conventional fertilisation (Deshmukh *et al.*, 2017).

Iron is an essential key enzymes Co-factor which performs different function in breakdown of hormone and also take part in many chemical reactions takes place in plants (Kerkeb and Connolly, 2006). Iron is required for the formation of chlorophyll even though it is not a component of it. As a result, a lack of chlorophyll causes iron deficiency and chlorotic plants (Katyal and Randhawa, 1983).

Zinc has been the micronutrient that crops, especially pulses, have required in sufficient amounts (Yashona *et al.*, 2018). Zinc in plants required for biosynthesis of hormone

(Kudi *et al.*, 2018). Zinc is essential for metabolism and participates in the production of nodules, which fix nitrogen (Patel *et al.*, 2011). Zinc is a crucial component of RNA polymers and gives ribosomes their structural integrity (Yashona *et al.*, 2018). Since Zn is required for the synthesis of tryptophan (Brown *et al.*, 1993; Alloway, 2004), Zn is a precursor of IAA, a metal that is also involved in the synthesis of the vital growth hormone Auxin (Brennan, 2005). Zinc is required for pollen function and fertilization (Ali *et al.*, 2017).

The availability of Molybdenum in soil has been shown to enhance the production of leguminous crops (Chandra *et al.*, 2020). Molybdenum, which is essential for several metabolic activities and regulates ascorbate-glutathione in plants. Plants deficient in molybdenum have lower quantities of chlorophyll and abnormal changes to their chloroplasts (Nasar *et al.*, 2017). The synthesis of the nitrate reductase enzyme and the symbiotic nitrogen fixation in legumes depend on the crucial micronutrient molybdenum (Williams and Fraustoda Silva, 2002; Roy *et al.*, 2006). Molybdenum increases plant height, number of branches and pods plant<sup>-1</sup>, number of seeds plant<sup>-1</sup> and nodule production in legume crops. It also enhances yield quality (Togayet *et al.*, 2008).

## **RESULT AND DISCUSSION**

### **Effect on plant height:**

Stoyanova and Doncheva (2002) reported that The end result was the longest plant ever because of the buildup of zinc fertiliser in plant tissues, which changes important developmental processes like photosynthesis and chlorophyll biosynthesis. Alam *et al.* (2010) reported In an earlier trial, the longest plant was produced with a 2.5 kg Zn ha<sup>-1</sup> application. The height of the plants was strongly impacted by the application of 10 kg Zn ha<sup>-1</sup> (47.76 cm and 47.88 cm) in summer mungbean. (Ram and Kattiyar, 2013). El Sayed Hameda *et al.*, (2012) reported that the By applying a combination of microelements to the leaves of pea plants, the yield components indicated by plant height were significantly increased. Zinc in auxin mixture (Basavarajeswari *et al.*, 2018). The plant height of the pigeonpea crop was increased by applying 5 mg kg<sup>-1</sup> Zn added to the soil (Dube *et al.*, 2001). Some enzymes with significant roles in cell division and lengthening were activated by zinc, and as a result, the stem height increased (Seifi Nadergoliet *et al.*, 2011). Habbasha *et al.* (2013) observed 0.2% ZnSO<sub>4</sub> application to chickpea at seed filling stage result in significantly higher plant height (81.43 cm). Zn and Fe may have positive benefits when given topically because of their importance in respiration, photosynthesis, and other biochemical and physiological processes

as well as their function in increasing yields (Rietra *et al.*, 2017; Khan *et al.*, 2022). The increment in plant height of cowpea with zinc spray ( $1 \text{ g L}^{-1}$ ) has been documented by (Salehin and Rahman 2012). The application of molybdenum to the leaves has a considerable impact on plant parameters, including plant height and biological performance. The result was in conformity with those of Manjili *et al.* (2014). Hazra and Tripathi (1998) observed that molybdenum application at the rate of  $1.5 \text{ kg/ha}$  increased forage in calcareous soil. Sahu *et al.* (2008) reported that the application of  $\text{FeSO}_4$  at  $2 \text{ kg ha}^{-1}$  significantly increased the growth characters over control in chickpea. Thapa *et al.* (2003) concluded that the application of micronutrients like Fe (as ferrous sulphate at 0.4%), Mn, Cu and Zn significantly increased the growth attributes in pea.

### **Effect on number of nodules plant<sup>-1</sup>**

Increased infection and rhizobial colonisation in the rhizosphere may be the source of the increased nodule count due to the increased availability of micronutrients (Meena *et al.*, 2012). Chauhan *et al.* (2013) reported 91% higher nodulation in soybean under the application of Zn @  $5 \text{ kg ha}^{-1}$  in consecutive years of study. Ram and Kattiyar (2013) found that the application of Zn @  $10 \text{ kg ha}^{-1}$  to mungbean significantly increase the nodulation. Badar *et al.*, (2015) “Molybdenum is essential for the development of nodule tissue and an increase in  $\text{N}_2$  fixation, both of which are necessary for Rhizobium to increase the nitrogen fixation process. Every bacterium that fixes nitrogen requires molybdenum since it is an essential component of the nitrogenase enzyme reported by Hirpara *et al.* (2019). Iron and molybdenum significantly enhanced nodules number in legumes crops (Brkics *et al.*, 2004). An agreement with Tantawy *et al.* (2013) reported that Fe and Mo significantly increased number of active nodules in broad bean.

### **Effect on dry matter accumulation:**

The production of straw was enhanced by applying multiple foliar sprays of various fertiliser mixtures at distinct growth stages. As previously documented, foliar treatment of macro and micronutrients boosted biological yield by Piri Issa *et al.*, (2012). The improvement in nodulation might have resulted in a higher amount of nitrogen fixation and there by better vegetative growth and dry matter production” Balla *et al.* (2020).

### **Effect on protein content:**

Zinc treatment to the leaves resulted in higher protein content as a result of improved physiological properties and protease enzyme activity Nandan *et al.* (2018), Usman *et al.* (2015) and Lokhande *et al.* (2018). Singh and Singh (2012) also reported 20.2% higher protein content in chickpea grain under the application of 10 kg Zn ha<sup>-1</sup>. Khrogamy and Farnia (2009) on zinc application in chickpea revealed the fact that, the application of ZnSO<sub>4</sub> @ 20 kg ha<sup>-1</sup> improved the protein content (24.02%) in chickpea. Imsande (1998) stated that iron had an important role in the synthesis of chlorophyll and also help in the absorption of other nutrients. Oguz (2004) that increasing Mo application up to 6 g kg<sup>-1</sup> considerably affected the protein content in chickpea. Roy *et al.* (2014) reported that the combined application of 5.5 kg Zn ha<sup>-1</sup> + 0.1% Zn as foliar spray increased the seed crude protein by 26.9% over control. Sharma *et al.* (2010) reported that the treatment of chelated Fe (1 or 2 kg ha<sup>-1</sup>) considerably enhanced the protein content in seed as well as all yield-contributing attributes in the pigeon pea crop.

#### **Effect on yield attributes:**

The beneficial effects of foliar nutrition of micro nutrients on mung bean were also reported by several researchers Kumawat *et al.* (2005). The micronutrients might have enhanced role in seed setting that resulted in improvement in no. of seeds pod<sup>-1</sup>.

#### **Effect on number of pods plant<sup>-1</sup>:**

Kumar *et al.* (2014) reported that seed inoculation with zinc @ 5 kg ha<sup>-1</sup> gave the significantly higher growth and yield attributes. Togay *et al.* (2008) reported that molybdenum increased pods plant<sup>-1</sup>. Valenciano *et al.* (2010) showed pod number plant<sup>-1</sup> were greater with Zn application. Ram and Katiyar (2013) found that the application of 10 kg Zn ha<sup>-1</sup> to mungbean resulted in significant improvement in the pod and seed number. In keeping with our findings, it was reported that micronutrient foliar spraying increased the number of pods plant<sup>-1</sup> in lentil (Zeidan *et al.*, 2006). Shah *et al.* (2016) concluded that the number of seeds pod<sup>-1</sup> and number of pods plant<sup>-1</sup> in pigeon pea significantly increase under the application of Zn @ 20 kg ha<sup>-1</sup>. Increase of pods plant<sup>-1</sup> using Zn spraying reported by Teixeira *et al.* (2004) in common bean. A decrease in flower shedding, a rise in pod formation, and the improvement of photosynthetic material distribution to these organs could all boost the pod plant<sup>-1</sup> (Mady, 2009).

#### **Effect on number of seeds pod<sup>-1</sup>**

Rizk and Abdo (2001) reported that yield components showed a highly significant increase with the application of Zn (0.2 or 0.4 g/l) compared to the control. The reports of other researchers that indicated the foliar application of Zn increased seeds pod<sup>-1</sup> in common bean (Teixeira *et al.*, 2004; Seifi Nadergholiet *al.*, 2011; Nasri *et al.*, 2011) and Faba bean (Bozorgi *et al.*, 2011). Sarkar and Banik (1991) observed the molybdenum application significantly increased seeds pod<sup>-1</sup>.

### Seed yield

Zn application at the rate of 0.3% in combination with Fe increased seed yield and lead to the highest amounts of seed yield. Khrogamy and Farnia (2009) It was noted that seed output and total biomass in chickpea cultivars were significantly affected by the treatment of ZnSO<sub>4</sub> @ 20 kg ha<sup>-1</sup>. Ramakrishna *et al.* (2005) studied the effects of improved production technologies involving application of 10 kg Zn ha<sup>-1</sup> and found increased pigeonpea yield. Patil *et al.* (2006) The application of ZnSO<sub>4</sub> @ 10 kg ha<sup>-1</sup> yielded considerably greater grain (29.55 q ha<sup>-1</sup>), stalk (95.81 t ha<sup>-1</sup>), and bhusa yields (20.58 q ha<sup>-1</sup>) when four different amounts of zinc (0, 10 and 20 kg ha<sup>-1</sup>) were investigated to see how they affected pigeonpea growth and yield. Ibrahim and Ramadan (2015) indicated Zn spraying increased seed yield and growth of dry bean. Malik *et al.* (2015) When the impact of zinc on mungbean growth and yield was examined, it was discovered that the treatment of 20 ppm zinc resulted in the highest seed yield plant<sup>-1</sup> (78.20 g), compared to the control. Shah *et al.* (2016) demonstrated that the treatment with Zn @ 20 kg ha<sup>-1</sup> had the highest pigeonpea yield. We discovered that the increase in bean seed yield brought on by Fe and Zn spraying was comparable to the others (Valenciano *et al.*, 2010; Naseri Rad *et al.*, 2014; Nasri *et al.*, 2011) soybean (Heidarian *et al.*, 2011) and Faba bean (Bozorgi *et al.*, 2011). The grain yield of chickpea was significantly increased with the application of iron and molybdenum (Khan *et al.*, 2014). Zinc fertilization had positive effect on the yield of pulses. Application of Zn @ 5 kg ha<sup>-1</sup> along with S @ 60 kg ha<sup>-1</sup> recorded 67% higher seed yield over control in mung bean (Mali *et al.*, 2003.) Application of FeSO<sub>4</sub> @ 2 kg ha<sup>-1</sup> along with bio-fertilizer inoculation in chickpea led to higher grain and straw yields over control (Sahu *et al.* 2008). Thapua *et al.* (2003) observed that the application of micronutrients like Fe (as FeSO<sub>4</sub> at 0.4%), Mn, Cu and Zn significantly increased the grain yield in pea. Salam *et al.* (2004) conducted a field experiment at Raipur, Chhatisgarh and concluded that the seed yield of Urd bean under application of FeSO<sub>4</sub> at 2-20 kg Fe ha<sup>-1</sup> was maximum over control. Kumawat *et al.* (2006) observed that the soil application of Fe at 25 kg Fe S

O<sub>4</sub>ha<sup>-1</sup> recorded significantly higher seed and straw yield of summer mungbean. Kumaretal. (2009) conducted an experiment at Kanpur and results revealed that the application of 10 kg Fe ha<sup>-1</sup> increase the grain yield of chickpea by 17.3% over control.

## References

- Alam, M.K., Uddin, M.M., Ahmed, M., Latif, M.A. and Rahman, M.M. 2010. Growth and green pod yield of garden pea varieties under different nutrient levels. *Journal of Agroforestry and Environment* **4**(1); 105–107.
- Ali, A., Ahmad, B., Hussain, I., Ali, A. and Shah, F.A. 2017. Effect of phosphorus and zinc on yield of lentil. *Pure and Applied Biology* **6**(4), 1397–1402.
- Alloway, B.J. 2004. Zinc in Soils and Crop Nutrition. Publ. of International Zinc Association. [http://www.iza.com/Documents/Communications/Publications/ALLOWAY\\_PRINT.pdf](http://www.iza.com/Documents/Communications/Publications/ALLOWAY_PRINT.pdf)
- Badar, R., Nisa, Z. and Ibrahim, S. 2015. Supplementation of P with rhizobial inoculants to improve growth of Peanut plants. *International Journal of Applied Research* **1**(4):19-23.
- Balla, H., Singh, V., Tiwari, D. and Shaik, M. A. 2020. Effect of boron and molybdenum on growth rate and yield of groundnut (*Arachis hypogea* L.). *J Pharmacogn Phytochem.* **9**(6):1416-1419.
- Basavarajeswari, 2018. Expression profiling of zinc transporter genes in tomato grown under different concentrations of zinc. *International Journal of Microbiology Research.* **10**(6): 1252-1255.
- Bozorgi, H. R., Faraji, A., Danesh, R. K., Keshavarz, A., Azarpour, E. and Tarighi, F. 2011. Effect of plant density on yield and yield components of rice. *World Applied Sciences Journal*, **12**(11), 2053- 2057. Retrieved from <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=4db23100f9c40e0bf212d5ca1afc0453157aaee2>
- Brennan, R.F. 2005. Zinc Application and Its Availability to Plants. Ph.D. dissertation, School of Environmental Science, Division of Science and Engineering, Murdoch University.
- Brikics, S., Milakovic, Z., Kristek, Z. and Antunovic, M. 2004. Pea yield and its quality depending on inoculation, nitrogen and molybdenum fertilization. *Plant, Soil and Environ.* **50**(1): 39-45.
- Brown, P.H., Cakmak, I. and Zhang, Q. 1993. Form and function of zinc in plants. Chap. 7, In A.D. Robson (Ed). pp 90-106. Zinc in Soils and Plants, Kluwer Academic Publishers, Dordrecht.

- Chandra, G., Gambhir, L. and Upadhyay, R. 2020. Effects of Biofertilizer with and without Molybdenum on Growth and Seed Yield of Chickpea under Doon Valley of Uttarakhand. *Current Journal of Applied Science and Technology*, **39**(15): 133–139.
- Chauhan, S., Titov, A. and Tomar, D. S. 2013. Effect of potassium, sulphur and zinc on growth, yield and oil content in soybean (*Glycine max* L.) in vertisols of central India. *Indian J. Appl. Res.* **3**(6): 489-491.
- Deshmukh, M. S., Zade, S. P., Gourkhede, P. H. and Inamke, P. S. 2017. Effect of High-P-Chelated Micronutrient Foliar Spray on Yield and Quality of Chickpea. *International Journal of Tropical Agriculture*, **35**(3): 707-712.
- Dube, B.K., Sharma, C. P. and Chatterjee, C. 2001. Response of Pigeonpea to Applied Zinc in Ustifluent Soils of Western Uttar Pradesh. *J. Indian Soc. Soil Sci.* **49**(3): 471-475.
- El Sayed Hameda, E.A., Amen El-Sh. A., EI-Morsy, A.H. and Tolba, M.H. 2012. Effects of foliar spraying with microelements and different fertilizer sources on quality and yield of *Pisum sativum* L. plant. *International Research Journal of Agricultural Science and Soil Science* **21**(1): 17–24.
- Habbasha, S. F., M. H. Mohamed, E. M. Abd El-Lateef, B. B. Mekki and M. E. Ibrahim, 2013. Effect of combined zinc and nitrogen on yield, chemical constituents and nitrogen use efficiency of some chickpea cultivars under sandy soil conditions. *World J. Agric. Sci.* **9**(4): 354-360.
- Hazra, C.R., Tripathi, S.B. 1998. Effect of secondary micronutrient on yield and quality of forages. *Fertilizer News* **43**(12): 77–82.
- Heidarian, A. R., Kord, H., Mostafavi, K., Parviz Lak, A. and Amini Mashhadi, F. 2011. Investigating Fe and Zn foliar application on yield and its components of soybean (*Glycine max* (L.) Merr.) at different growth stages. *Journal of Agricultural Biotechnology and Sustainable Development*, **3**(9):189-197. Retrieved from [https://academicjournals.org/article/article1379436818\\_Heidarian%20et%20al.pdf](https://academicjournals.org/article/article1379436818_Heidarian%20et%20al.pdf)
- Hirpara, D. V., Sakarvadia, H. L., Jadeja, A. S., Vekaria, L. C. and Ponkia, H. P. 2019. Response of boron and molybdenum on groundnut (*Arachis hypogea* L.) under medium black calcareous soil. *J PharmacognPhytochem.* **8**(5):671-677.
- Ibrahim, E. A. and Ramadanb, W. A. 2015. Cross effect of zinc foliar spray alone and combined with humic acid and chitosan on growth, nutrient elements content and yield of dry bean

(*Phaseolus vulgaris* L.) plants sown at different dates. *Scientia Horticulturae*, 184, 101–105.  
<https://doi.org/10.1016/j.scienta.2014.11.010>

Imsande, J. 1998. Iron, sulphur and chlorophyll deficiencies: A need for an integrative approach in plant physiology. *Plant Physiology*.**103**:139-144.

Katyal, J. C and Randhawa, N. S. 1983. Micronutrient. F.A.O. Fertilizer and plant nutrition. Bulletin No. 7. United Nation.

Kavya, P., Singh, S., Hinduja, N., Tiwari, D. and Sruthi, S. 2023. Effect of Foliar Application of Micronutrients on Growth and Yield of Green gram (*Vigna radiata* L.) *Legume Research - An International Journal* **44**(12): 1460-1464.

Kerkeb, L. and Connolly, E. 2006. Iron transport and metabolism in plants. *J. Gent. Eng.* **27**: 119-140.

Khan, M. M. H., Ahmed, N., Naqvi, S. A. H., Ahmad, B., Dawar, K., Rahie, A. A. and Danish, S. 2022. Synchronization of zinc and boron application methods and rates for improving the quality and yield attributes of *Mangifera indica* L. on sustainable basis. *Journal of King Saud University - Science*, **34**(8), 102280. <https://doi.org/10.1016/j.jksus.2022.102280>

Khan, N., Tariq M., Ullah, K., Muhammad, D., Khan I., Rahatullah, K., Ahmed, N. and Ahmed S. 2014. The Effect of Molybdenum and Iron on Nodulation, Nitrogen Fixation and Yield of Chickpea Genotypes (*Cicer arietinum* L.). *Journal of Agriculture and Veterinary Science*. **7**(1): 63-79.

Khorgamy, A. and Farina, A. 2009. Effect of phosphorus and zinc fertilisation on yield and yield components of chick pea cultivars. *African Crop Science Conference Proceedings*. **9**: 205-208.

Kudi, S., Swaroop, N., David, A.A., Thomas, T., Hasan, A. and Rao, S. 2018. Effect of different levels of sulphur and zinc on soil health and yield of Greengram (*Vigna radiata* L.) Var. Patidar-111. *Journal of Pharmacognosy and Phytochemistry***7**(3): 2271–2274.

Kumar, V., Dwivedi, V. N. and Tiwari, D. D. 2009. Effect of phosphorus and iron on yield and mineral nutrition in chickpea. *Ann. Plant Soil Res.* **11**:16-18.

Kumar, R., Deka, B.C., Kumawat, N. and Ngachan, S.V. 2014. Effect of integrated nutrition, biofertilizers and zinc application on production potential and profitability of garden pea (*Pisum sativum* L.) in Eastern Himalaya, India. *Legume Research***37**(6): 614–620.

- Kumawat, R. N., Rathore, P. S., Pareek, N. 2006. Response of mung bean to sulphur and iron nutrition grown on calcareous soil of Western Rajasthan. *Indian J. Pulse Res.* **19**:228-230.
- Kumawat, R.N., Rathore, P.S. and Talwar, H.S. 2005. Effect of sulphur and iron on crop growth attributes in summer green gram. *Indian Journal of Plant Physiology.* **10**(1): 86-89.
- Lokhande, P. B., Indulkar, B. S., Vaidya, P. H., Padghan, A. D., Wagh, C. B., Ingole, A. J., Patil, N. M. and Aundhkar, A. 2018. Effect of phosphorus and zinc on yield and quality of greengram (*Vigna radiata* L.) in *inceptisol*. *International Journal of Engineering Science and Computing.* **8**(7): 18647-18649.
- Mady, M. A. 2009. Effect of foliar application with yeast extract and zinc on fruit setting and yield of Faba bean (*Vicia faba* L.). *Journal of Biological, Chemical and Environmental Science*, **4**(2), 109-127. Retrieved from [https://www.bu.edu.eg/portal/uploads/Agriculture/Botany/1299/publications/Mohamed%20Ahmed%20Mohamed%20Mady\\_5.pdf](https://www.bu.edu.eg/portal/uploads/Agriculture/Botany/1299/publications/Mohamed%20Ahmed%20Mohamed%20Mady_5.pdf)
- Mali, G.C., Sharma, N.N., Acharya, H.K., Gupta, S.K. and Gupta, P.K. 2003. Response of pigeon pea to S and Zn fertilization *Vertisols* in South eastern plain of Rajasthan. Proceedings of National symposium on Arid Legumes for Food Nutrition Security and Promotion Trade, Hisar, India, 15-16 May 2002. Pp. 267-271.
- Malik, K., S. Kumar and K. P. S. Arya, 2015. Effect of zinc, molybdenum and urea on growth and yield of mungbean (*Vigna radiata* L. Wilczek). *Adv. Res. J. of Crop Impro.* **6**(1): 58-62.
- Manjili, M. J., Bidarigh, S. and Amiri, E. 2014. Study the effect of foliar application of Nano chelate molybdenum fertilizer on the yield and yield components of peanut. *Botany.* **5**(1):67-71.
- Meena, K. K., Meena, R. S. and Kumawat, S. M. 2012. Effect of sulphur and iron fertilization on yield attributes, yield and nutrient uptake of mungbean (*Vigna radiata*). *Indian Journal of Agricultural Sciences.* **83**(4): 472-6.
- Meena, K. K., Meena, R. S. and Kumawat, S. M. 2013. Effect of sulphur and iron fertilization on yield attributes, yield and nutrient uptake of mungbean (*Vigna radiata*). *Indian Journal of Agricultural Sciences* **83** (4): 472
- Nandan, B., Sharma, B. C., Chand, G., Bazgalia, K., Kumar, R. and Banotra, M. 2018. Agronomic fortification of Zn and Fe in chickpea an emerging tool for nutritional security - a global perspective. *Acta Scientific Nutritional Health.* **2**(4):12-19.

- Nasar, J. and Shah, Z. 2017. Effect of iron and molybdenum on yield and nodulation of lentil. *ARNP: Journal of Agriculture and Biological Sciences*, **12**(11), M332–339
- Naseri Rad, H., Sayad, V. and Naseri Rad, A. 2014. Effect of *rhizobium* bacteria (*Rhizobiumleguminosarum*) and nano-Iron application on yield and yield components of different pinto beans genotypes. *Agricultural Communication*, **2**(2), 22-27. Retrieved from [https://www.researchgate.net/publication/261885406\\_Effect\\_of\\_Rhizobium\\_Bacteria\\_Rhizobium\\_leguminosarum\\_and\\_Nano-Iron\\_Application\\_on\\_Yield\\_and\\_Yield\\_Components\\_of\\_Different\\_Pinto\\_Beans\\_Genotypes](https://www.researchgate.net/publication/261885406_Effect_of_Rhizobium_Bacteria_Rhizobium_leguminosarum_and_Nano-Iron_Application_on_Yield_and_Yield_Components_of_Different_Pinto_Beans_Genotypes)
- Nasri, M., Khalatbari, M. and Aliabadi Farahani, H. 2011. Zn-foliar application influence on quality and quantity features in Phaseolous Vulgaris under different levels of N and K fertilizers. *Advances in Environmental Biology*, **5**(5), 839-846. Retrieved from <http://www.aensiweb.com/old/aeb/2011/839-846.pdf>
- Oguz, F. 2004. Research on the effects of different levels Molybdenum application on the yield and some yield components in Chickpea varieties on dry and irrigation conditions. M.Sc. Thesis, YuzuncuYil University Van, Turkey (unpublished).
- Patel, M.M., Patel, I.C., Patel, R.I. and Acharya, S. 2011. Effect of Zinc and Iron on yield and yield attributes of rainfed cowpea (*Vigna unguiculata* L. Walp). *Annals of Arid Zone*, **50**(1): 17–19.
- Patil, H. M., Shete, B. T. and Gaikwad, C. B. 2006. Effect of potash and zinc levels on growth and yield of pigeonpea. *Ecology*, **12**(2): 341-344.
- Piri Issa, 2012. Effect of phosphorus and micronutrients foliar application on sorghum yield. *Annals of Bio. Res.* **3**(8): 3998-4001.
- Ram, S. and Katiyar, T. P. S. 2013. Effect of sulphur and zinc on the seed yield and protein content of summer mungbean under arid climate. *Int. J. Sci. Nature*, **4**(3): 563-566.
- Ramakrishna, A., Wani, S. P., Rao, S. C. and Reddy, S. U. 2005. Effect of improved crop production technology on pigeonpea yield in resource poor rainfed areas. *J. SAT Agric. Res.*, **1**(1): 1-2.
- Rietra, R. P., Heinen, M., Dimkpa, C. O. and Bindraban, P. S. (2017). Effects of nutrient antagonism and synergism on yield and fertilizer use efficiency. *Communications in Soil Science and Plant Analysis*, **48**, 1895–1920. <https://doi.org/10.1080/00103624.2017.1407429>.

- Rizk, W.M. and Abdo, F.A. 2001. The response of two mungbean cultivars to zinc, manganese and boron II. Yield and chemical composition of seeds. *Bulletin of Faculty of Agriculture, Cairo University* **52**(3): 467–477.
- Roy, P. D., Narwal, R. P., Malik, R. S., Saha, B. N. and Kumar, S. 2014. Impact of zinc application methods on green gram (*Vigna radiata* L.) productivity and grain zinc fortification. *J. Environ. Biol.* **35**(5): 851-855.
- Roy, R.N., Finck, A., Blair, G.J. and Tandon, H.L.S. 2006. Plant nutrition for food security. A guide for integrated nutrient management. *FAO Fertilizer and Plant Nutrition Bulletin* 16. FAO, Rome, Italy. 368
- Sahu, S., Lidder, R. S. and Singh, P. K. 2008. Effect of micronutrients and biofertilizers on growth, yield and nutrient uptake by chickpea (*Cicer arietinum* L.) in *Vertisols* of Madhya Pradesh. *Adv. Plant Sci.* **21**:501-503.
- Salam, P. K., Rajput, R. S., Mishra, P. K., Anita and Shrivastava, G. K. 2004. Effect of micronutrient fertilization on productivity potential of mungbean. *Ann. Agric. Res. New Series* **25**:329-332.
- Salehin, F. and Rahman, S. 2012. Effect of zinc and nitrogen fertilizer and their application method on yield and yield components of *Phaseolus vulgaris* (L.) *Agric. Sci.* **3**(1): 9-13.
- Sarkar, R.K., Banik, P. 1991. Response of green gram to nitrogen, phosphorus and molybdenum. *Indian Journal of Hill Farming* **5**(1), 75–76.
- Seifi Nadergholi, M., Yarnia, M. and Rahimzade Khoei, F. 2011. Effect of zinc and manganese and their application method on yield and yield components of common bean (*Phaseolus vulgaris* L. CV. Khomein). *Middle-East Journal of Science and Research*, **8**(5): 859-865. Retrieved from [https://www.idosi.org/mejsr/mejsr8\(5\)11/3.pdf](https://www.idosi.org/mejsr/mejsr8(5)11/3.pdf)
- Shah, K. A., Gurjar, R., Parmarand, H. C. and Sonani, V. V. 2016. Effect of sulphur and zinc fertilization on yield and quality of pigeonpea in sandy loam soil. *Green Farming*, **7**(2): 495-497.
- Sharma, A., Nakul, H. T., Jelgeri, B. R. and Surwenshi, A. 2010. Effect of micronutrients on growth, yield and yield components in pigeon pea (*Cajanus cajan* L.). *Res. J. Agric. Sci.* **1**:142-144.

- Singh, G., H. S. Sekhon and Harpreet Kaur, 2012. Effect of farmyard manure, vermicompost and chemical nutrients on growth and yield of chickpea (*Cicer arietinum* L.). *Int. J. Agric. Res.* **7**(2): 93-99.
- Stoyanova, Z., Doncheva, S., 2002. The effect of zinc supply and succinate treatment on plant growth and mineral uptake in pea plant. *Brazilian Journal of Plant Physiology***14**(2):111–116.
- Tantawy, E. L. E. M and Dalia, A. S. N. 2013. Nodulation, growth, photosynthetic pigments and yield of broad bean plant (*Vicia faba* L.) as affected by nitrogen source, *rhizobium* inoculation and iron foliar application. *J. Applied Sciences Research.* **9**(1): 974-987.
- Teixeira, I. R., Boren, A. B. and Araujo, G. A. A. 2004. Manganese and zinc leaf application on common bean grown on a cerrado soil. *Scientia Agricola*, **61**, 77-81. <https://doi.org/10.1590/S0103-90162004000100013>
- Thapu, U., Rai, P., Suresh, C. P. and Pal, P. 2003. Effect of micronutrients on the growth and yield of pea in gangetic alluvial of West Bengal. *Environ. Ecol.* **21**:179-182.
- Togay, Y., Togay, N. and Dogan, Y. 2008. Research on the effect of phosphorus and molybdenum applications on the yield and yield parameters in lentil (*Lens culinaris* Medic.). *African Journal of Biotechnology***7**(9), 1256–1260.
- Togay, Y., Togay, N., Dogan, Y., 2008. Research on the effect of phosphorus and molybdenum applications on the yield and yield parameters in lentil (*Lens culinaris* Medic.). *African Journal of Biotechnology***7**(9): 1256–1260
- Usman, M., Tahir, M. and Majeed, M. A. 2014. Effect of zinc sulphate as soil application and seed treatment on greengram (*Vigna radiata* L.). *Pakistan Journal of Life and Social Sciences.***12**(2):87- 91.
- Valenciano, J. B., Miguélez-Frade, M. M., Marcelo, V. and Reinoso, B. 2010. Response of irrigated common bean (*Phaseolus vulgaris*) yield to foliar zinc application in Spain. *New Zealand Journal of Crop and Horticultural Science*, **35**(3): 325-330. <https://doi.org/10.1080/01140670709510198>
- Williams, R.J.P., Frausto da Silva, J.J.R., 2002. The involvement of molybdenum in life. *Biochemical and Biophysical Research Communications* **292**, 293–299
- Yashona, D.S., Mishra, U.S. and Aher, S.B. 2018. Response of Pulse Crop to Sole and Combined mode of Zinc Application. *Journal of Soils and Crops***28**(2): 249–258

Zeidan, M. S., Hozayn, M. and Abd-El-Salam, M. E. E. 2006. Yield and quality of lentil as affected by micronutrient deficiencies in sandy soils. *Journal of Applied Sciences Research*, **2**(12), 1342- 1345. Retrieved from <http://www.aensiweb.com/old/jasr/jasr/1342-1345.pdf>

UNDER PEER REVIEW