

Original Research Article

Antibiogram of *Pseudomonas aeruginosa* Clinical Isolates Tested for Pan, Extensive and Multidrug Resistances

ABSTRACT

Background: *Pseudomonas aeruginosa* usually cause nosocomial infections with concurrent morbidity and mortality and is generally resistant to many antibiotics.

Aim: This is an interesting study that aimed to determine the proportion of pan-drug-resistant (PDR), extensively drug-resistant (XDR), and multidrug-resistant (MDR) *P. aeruginosa* strains recovered from human samples.

Methodology: The retrospective study was conducted in the University of Nigeria Teaching Hospital Enugu in 2023. Clinical samples obtained from patients between September 2017 and July 2019 were analysed. A total of 100 *Pseudomonas aeruginosa* isolates recovered from 780 clinical samples were used. Standard microbiological techniques were used to identify and categorize the isolates. Antibiotic susceptibility pattern was determined using the Kirby Bauer disc diffusion method by Clinical and Laboratory Standard Institute (CLSI) guidelines.

Results: Isolates recovered from wound was 38%, voided urine, catheter tip urine, ear swab, and high vaginal swab samples recorded 29%, 11%, 10% and 4%, respectively. Ceftazidime recorded the highest level of resistance (70.0%) and the least was Colistin (20%). Resistance patterns showed that 32(32.0%) bacterial strains were MDR, 68(68.0%) were XDR and no PDR was recorded.

Conclusions: For the best selection of empirical therapy, *P. aeruginosa* susceptibility monitoring is essential due to the high prevalence of antibiotic resistance. The resistance pattern raises the possibility of misuse of broad-spectrum antibiotics. Treatment for bacterial infections should be directed by the results of antimicrobial susceptibility tests.

Keywords: *Pseudomonas aeruginosa*, antibiotic, nosocomial, resistance, infections

1. INTRODUCTION

Pseudomonas species belong to the family Pseudomonadaceae, they are aerobic, Gram-negative, rod-shaped, and polar-flagellated organisms. The most widespread species of medically important bacteria is *Pseudomonas aeruginosa*, an organism that is present in a wide variety habitats and it is responsible for nosocomial infections in clinical settings [1, 2]. The organism is an opportunistic organism that is a principal contributor to morbidity and mortality, particularly among people with cystic fibrosis and other immune system disorders [3]. It causes infections in the blood, surgical sites, eye, external ear, urinary tract, respiratory tract, and wounds (particularly in burn victims) [1, 2].

The goal of initial antimicrobial regimen for patients suspected of severe *P. aeruginosa* infections is to decrease mortality. This therapy includes mono therapy and combination therapy [4, 5]. Nevertheless, because *P. aeruginosa* has the ability to resist the majority of the currently available antibiotics, treating *P. aeruginosa* infections has grown to be a significant concern [6]. *Pseudomonas aeruginosa* is one of the bacteria that was classified among 12 bacteria by the World Health Organization lists of antibiotic-resistant "priority pathogens", and was included in the first category of critical Priority 1[7]. The occurrence of multidrug resistance in *P. aeruginosa* is accelerated by the misuse of antibiotics during treatment, rendering the empirical antimicrobial treatment ineffective against these bacteria [8]. *Pseudomonas aeruginosa* shows resistance to a wide range of antimicrobials [9, 3]. There are three main mechanisms *P. aeruginosa* uses to resist antimicrobials including, acquired, intrinsic, and adaptive resistance. Intrinsic resistance involves creation of efflux pumps that remove drugs from the cells, low outer membrane permeability, and the production of enzymes that render antibiotics inactive[10]. Adaptive resistance is concerned with the formation of biofilm [11] which can support the growth of multidrug-tolerant persister cells that survive the antimicrobial therapy [12].

Antimicrobial resistance in *P. aeruginosa* and other organisms has been on the increase globally [13, 14]. Based on the degree of their resistance, the isolates have been labeled as multidrug-resistant (MDR), extremely drug-resistant (XDR), and pan-drug-resistant (PDR). Since there are few effective antibiotic treatments available, infections with these resistant strains may lead to a rise in morbidity and mortality [8, 15].

Reviews on MDR *P. aeruginosa* have shown a wide range of definitions[16, 17]. The absence of definition for MDR in clinical research methods makes it hard to compare data [18]. However, in most of the research, multidrug resistance was defined as resistance to at least three drugs from different antibiotic group categories, including carbapenems, antipseudomonal penicillins, aminoglycosides, cephalosporins, and fluoroquinolones [8].

A team of experts got together to propound standardized international nomenclature for determining acquired resistance profiles in all bacteria that commonly cause nosocomial infections and are at risk for multidrug resistance [18]. For each bacterium, epidemiologically significant antimicrobial categories were developed. The US Food and Drug Administration (FDA), the European Committee on Antimicrobial Susceptibility Testing (EUCAST), and the Clinical Laboratory Standards Institute (CLSI) data and breakpoints were utilized. Acquired resistance to at least one antimicrobial agent across three or more different antimicrobial groups is referred to as MDR, whereas XDR refers to resistance to at least one antimicrobial agent across all categories except two or fewer. Resistance to all antimicrobial agents was defined as PDR. To be certain that these definitions are accurately applied, all or almost all of the antimicrobials mentioned under the antimicrobial categories must be employed. Additionally, results must be reported accurately [18].

In Nigeria, most studies on MDR in *P. aeruginosa* did not include these criteria and there is, therefore, a dearth of information on the prevalence of these phenotypes, hence this study. The purpose of this study, therefore, was to investigate the antimicrobial resistance profile of *Pseudomonas aeruginosa* and to evaluate the prevalence of MDR, XDR, and pan drug-resistance phenotypes in *P. aeruginosa* from human samples.

2. METHODOLOGY

2.1 Study Area

This study was conducted in Enugu Metropolis. It is located in southeast Nigeria with a population of 820,000 based on the last census. There are three tertiary hospitals including the University of Nigeria Teaching Hospital (UNTH) Ituku/Ozalla, National Orthopedic Hospital, and Enugu State University Teaching Hospital.

2.2 Bacteriological Analysis

The isolates were recovered from clinical samples that were submitted to the Microbiology Laboratory of the University of Nigeria Teaching Hospital Ituku-Ozalla. They were sub-cultured from the stock cultures unto nutrient agar (Neogen, LAB008, KR) and then cultured unto Centrimide agar (HiMedia, MH024, IND). *Pseudomonas aeruginosa* were isolated from the various clinical samples using the cetrimide Agar selective medium. The formation of pigment by the production of fluorescein and pyocyanin appearing as greenish colonies was a prominent confirmatory identification characteristic of the colonies after 24 hours incubation at 37°C. The oxidase enzyme test for the presence of cytochrome oxidase enzymes was also carried out. The change of tetra-methyl- p-phenylenediamine dihydrochloride reagent to deep purple colour indicated positive. Other Biochemical confirmatory tests were carried out according to standard methods [19]. A total of 100 non-duplicates *Pseudomonas. aeruginosa* were positive among 780 clinical isolates from samples of urine, wound swab, high vaginal swab, catheter tip, ear swabs, and sputum.

2.3 Antimicrobial Susceptibility Testing

The Kirby Bauer's disk diffusion technique as set out in the recommendations of the Clinical Laboratory Standard Institute (CLSI) was used [20]. The disc diffusion technique involved swabbing a standardized inoculum onto the surface of Muller Hinton agar (Oxoid, CMO337B, UK). The agar surface was inoculated by using a swab dipped in the *Pseudomonas* cells suspension adjusted to the turbidity of a 0.5 McFarland standard and was spread evenly over the surface, avoiding the edges of the plates during the swabbing. There after the antibiotic disks were picked by sterile forceps and dropped on the surface of the agar seeded with the *Pseudomonas* inoculum. The antibiotics were allowed to diffuse into the agar before incubation. The plates were then incubated at 37°C for 18 to 24 hours. After the incubation, the Inhibition Zone Diameter (IZD) was measured and recorded. The zone sizes were read using standardized chart to record the result as sensitive, resistant, or intermediate. The following antibiotics were included; ceftazidime (30µg), Cefepime (30µg), Piperacillin-tazobactam (110 µg) Amikacin (30µg), gentamicin (10µg), levofloxacin (5µg), ciprofloxacin (5µg), imipenem (10 µg), Meropenem (10 µg) Aztreonam (30µg), and colistin (10 µg). *P. aeruginosa* ATCC 27853 was used for the quality control. It was done in parallel with test isolates for each susceptibility test.

2.4 Detection of MDR, XDR, and PDR

MDR, XDR, and PDR in *P. aeruginosa* isolates were defined following a structured global report [17], and by the outcome of antimicrobial susceptibility profile against all antimicrobial categories. Non-susceptibility to at least one agent in ≥3 antimicrobial categories was recorded MDR, non-susceptibility to at least one agent in ≥ 6 antimicrobial categories were reported XDR, and the isolates that showed non-susceptibility to all the antibiotic categories were called PDR. Phosphonic acids (Fosfomycin) were not used because of the absence of susceptibility breakpoints for the drug against *P. aeruginosa*.

2.5 Statistical Analysis

For all statistical calculations, SPSS for Windows version 20 (SPSS, Chicago, IL, USA) was used. Descriptive statistics (frequencies and percentages) were used to describe categorical variables. Pearson's Chi-square test (X^2) was used to test for significant association between variables at a 95% confidence interval. Statistical significance was defined as a P-value of 0.05 or lower.

3. RESULTS

Table 1 shows the distribution of clinical isolates according to source, of the 100 isolates of *Pseudomonas aeruginosa*, the highest number of isolates was recovered from wound 38 (38.0%), followed by voided

urine 29 (29.0%). Others were catheter tips 11 (11.0%), ear swab 10 (10.0%), sputum 8 (8.0%) and the least was from HVS 4 (4.0%).

Table 1: Distribution of Isolates According to the Source

Isolate Source	No/ %
Wound	38 (38.0)
Urine	29 (29.0)
Sputum	8 (8.0)
Catheter tip	11 (11.0)
Ear swab	10 (10.0)
High Vaginal Swab (HVS)	4 (4.0)
Total	100 (100.0)

Table 2 shows the resistance rates of the antimicrobials, Gentamicin, Amikacin, Imipenem, Meropenem, Ceftazidime, Cefepime, Ciprofloxacin, Levofloxacin, Piperacillin-tazobactam, Aztreonam, and Colistin as 40.0%, 44.0%, 68.0%, 62.0%, 70.0%, 55.0%, 50.0%, 51.0%, 55.0%, 58.0%, and 20.0%, respectively.

Table 2: Antibigram of *P. aeruginosa* from different clinical samples

Antimicrobial categories	Antimicrobial agents	Susceptible No/%	Intermediate No/%	Resistant No/%
Aminoglycosides	Gentamicin	50 (50.0)	10 (10.0)	40 (40.0)

	Amikacin	52 (52.0)	4 (4.0)	44 (44.0)
Carbapenems	Imipenem	31 (31.0)	1 (1.0)	68 (68.0)
	Meropenem	34 (29.0)	4 (3.0)	62 (62.0)
Cephalosporin	Ceftazidime	26 (26.0)	4 (4.0)	70 (86.0)
	Cefepime	40 (40.0)	5 (5.0)	55 (55.0)
Fluoroquinolones	Ciprofloxacin	48 (48.0)	2 (2.0)	50 (50.0)
	Levofloxacin	46 (46.0)	3 (3.0)	51 (51.0)
Penicilins/ β -lactamase inhibitors	Piperacillin/tazobactam	40 (60.0)	5 (5.0)	55 (35.0)
Monobactams	Aztreonam	32 (32.0)	10 (10)	58 (58.0)
Polymyxins	Colistin	80 (80.0)	0 (0)	20 (20.0)

Table 3 shows the prevalence of resistant types according to the source of the isolate and the different resistance patterns rates recorded. Out of the 38 isolates from the wound, 20 (62.5%) were MDR, and 18 (26.5%) were XDR. Urine isolates recorded 3 (9.4%) MDR and 26 (38.3%) were XDR. The highest number of isolates with MDR pattern was from wound 20(62.5%), followed by catheter tips 5% 15.6%, and the least was from ear swabs and HVS 2(6.3%) each. The highest number with XDR pattern was from Urine 26 (38.2%), followed by Wound 18(26.5%), and the least was from HVS 2(2.9%). Out of the 100 isolates of *P. aeruginosa*, XDR ranked highest 68(68.0%) and MDR was the lowest 32 (32.0%). There was no PDR isolates.

Table 3: Prevalence of Resistance Types According to the Source of Isolates

Source of Isolate	MDR	XDR	Total
Wound	20 (62.5)	18 (26.5)	38 (38.0)
Urine	3 (9.4)	26 (38.2)	29 (29.0)

Sputum	0 (0.0)	8 (11.8)	8 (8.0)
Catheter tip	5 (15.6)	6 (4.4)	11 (11.0)
Ear swab	2 (6.3)	8 (11.8)	10 (10.0)
High Vaginal Swab (HVS)	2 (6.3)	2 (2.9)	4 (4.0)
Total	32 (32.0)	68 (68.0)	100 (100.0)
	X²= 16.63		

4. DISCUSSION

Pseudomonas aeruginosa continues to be a significant hospital-acquired organism known for its morbidity and mortality, especially in immunocompromised people and susceptible patients in intensive care units [21]. It is possible to isolate the bacterium from any clinical sample. In this work, majority of the isolates were mostly found in wounds and this is in line with what was reported in Ethiopia [1]. This may be as a result of contaminated surgical tools and the environmental proliferation of *P. aeruginosa* in healthcare facilities [1; 22].

Our research revealed that carbapenems, such as imipenem (68%), and meropenem (62.0%), had higher rates of resistance than cephalosporins and fluoroquinolones. The evolvement of carbapenemase-producing bacteria has restricted the use of carbapenems, which were once very effective anti-pseudomonal drugs [23]. Our findings were closely related to those reported in Mexico 70% and 54% [24] and 53% and 63% in India [25]. A lower resistance rate was recorded in Ethiopia 18% and 13% [1]. The higher resistance rate observed in our research may be attributable to prescription practices used in our clinical setting, inappropriate use of broad-spectrum antibiotics, and a special trait of *P. aeruginosa* that makes it susceptible to acquiring resistance such as low cell wall permeability, formation of inducible cephalosporinases and efflux pumps, and a poor affinity to the target sites [26]. Our study also showed higher resistance among imipenem as against meropenem. This is consistent with the study of Addis *et al* and Kateete *et al* [1; 23]. This might be due to differences in the chemical structures of the two drugs. Meropenem is more effective against *P. aeruginosa* because it penetrates the outer membrane porin-D (OprD) more quickly. However, due to its increased risk of membrane selection, imipenem has been less effective [1].

The isolates encountered during this investigation displayed considerable cephalosporin resistance, especially to ceftazidime (70%), and cefepime (55%). This is in consonance with what was reported in other nations. In Uganda, 69% and 55% for ceftazidime and cefepime, 63% and 62% in Egypt, 65 and 55% in Mexico, 66% and 63% in India were reported [22; 23; 24; 26]. However, lower prevalence has been reported in Iran 35% and 38%, and in Ethiopia, 35% and 31% [1; 28]. Over-production of beta-lactamases in particular may be the cause of the increased resistance that has been observed in Nigeria and other nations. Improper prescription of cephalosporins causes the pathogen to undergo a genetic change.

This present study also showed that combination drug penicillin/ beta-lactamase inhibitor, Piperacillin-tazobactam recorded 55% resistance and was better than carbapenems, and cephalosporins.

The fluoroquinolones, ciprofloxacin (50.0%) and levofloxacin (51.0%) were more effective than cephalosporins. The result is consistent with what was reported in Uganda 64% and India 67% for ciprofloxacin [23; 24] but at variance with what was reported in Ethiopia 18% and 24% for ciprofloxacin and levofloxacin respectively. This disparity may be caused by the diverse sample sizes, various study settings, and vast geographical differences [1].

Aminoglycoside was more potent than cephalosporins, fluoroquinolones, and carbapenem. Amikacin was 44.0% and Gentamicin 40.0%. Our findings were comparable to what was reported in Mexico (58.0% and 52%, in Uganda 31% and 69%) but higher than what was reported in Ethiopia [1], which was 2% and 7%. The effectiveness of this drug might be because of lower prescription practice in our setting.

In this study, colistin was the most potent drug for *P. aeruginosa* infection despite having an overall resistance rate of 20%. The nephrotoxicity and neurotoxicity of this drug deterred clinicians from using it in the past. It is a reserved drug used to treat confirmed or suspected infections caused by MDR pathogens [1]. Our colistin-resistance rate of 20% was consistent with the 23% reported in Egypt [27] but higher than the 6% and 9% reported in Ethiopia and Iran respectively [1; 28]. The discrepancy may be caused by methodological differences and/or the presence of colistin-resistant bacteria as a result of improper colistin use in veterinary medicine, where it has been frequently utilized to promote growth in animal husbandry [29]. Since no alternative antibiotics may be utilized, the rise of colistin-resistant *P. aeruginosa* strains is extremely worrisome and poses a severe global problem.

In this investigation, the prevalence of MDR and XDR *Pseudomonas aeruginosa* isolates was 32% and 68%, respectively. There was no PDR in this study. The definition of the acronyms was done according to the accepted global report for *P. aeruginosa* (18) except for Fosfomycin that was not included because of the absence of susceptibility breakpoints for fosfomycin against *P. aeruginosa*. Saderi and Owlia [28] reported 54.5% and 33% for MDR and XDR and there was no PDR recorded among 88 clinical isolates in Iran whereas Addis and his colleagues [1], reported 23%, 9%, and 2% for MDR, XDR, and PDR respectively in Ethiopia. There are few published data on multi-drug resistance using a proper definition of MDR. However, a high and lower prevalence of MDR was reported in many countries [29]. Variations in prevalence have been due to inappropriate use of antimicrobials, and geographical variations

5. CONCLUSION

Multidrug resistance is becoming a severe problem in hospital settings, increasing the incidence of nosocomial infections as its prevalence rate rises and spreads globally. Antibiotic resistance to formerly highly sensitive medications is on the rise. To identify trends in resistance, the sensitivity pattern should be periodically monitored over time. Additionally, a strategic approach to the limited and responsible use of antipseudomonal medicines is urgently needed to combat the morbidity and mortality of these strains. Antimicrobials work differently depending on where they are used, hence it is essential to update local treatment guidelines using this kind of research. Results of tests for antibiotic susceptibility should be used to guide treatment of bacterial infections

REFERENCES

1. Addis T, Araya S, Desta K. Occurrence of Multiple, Extensive and Pan Drug-Resistant *Pseudomonas aeruginosa* and Carbapenemase Production from Presumptive Isolates Stored in a Biobank at Ethiopian Public Health Institute. *Infection and Drug Resistance* 2021;14, 3609-3618.
2. Karami P, Mohajeri P, Mashouf RY. et al Molecular characterization of clinical and environmental *Pseudomonas aeruginosa* isolated in burn center. *Saudi J. Biol Sci.* 2019; 26 (7): 1731-1736. doi: 10.1016/j.sjbs.2018.07.009
3. Pang Z, Raudonis R, Glick BR, Lin T, Cheng Z (2019). Antibiotic resistance in *Pseudomonas aeruginosa*: mechanisms and alternative therapeutic strategies. *Biotechnology Advances* 2019 37(1) 177-192. Doi.org/10.1016/j.biotechadv.2018.11.013
4. El Solh AA, Alhajhusain A. Update on treatment of *Pseudomonas aeruginosa* pneumonia. *J. Antimicrob. Chemther.* 2009 Aug;64(2):229-38 doi:10.1083/jac/dkp201.
5. Park, S.Y., Park, H.J., Moon, S.M., Park, K.H., Chong, Y.P., et al. Impact of adequate empirical combination therapy on mortality from bacteremic *Pseudomonas aeruginosa* pneumonia. *BMC Infect Dis.* 2012;12: 308
6. Lister, P.D., Wolter, D.J., Hanson, N.D. Antibacterial-resistant *Pseudomonas aeruginosa*: clinical impact and complex regulation of chromosomally encoded resistance mechanisms. *Clin Microbiol Rev.* 2009; 22, 582–610

7. World Health Organization. WHO Publishes List of Bacteria for Which New Antibiotics Are Urgently Needed(2017) (accessed 17 Sep 2021) <https://www.who.int/News/item>
8. Hirsch, E.B., Tam, V.H. Impact of multidrug-resistant *Pseudomonas aeruginosa* infection on patient outcomes. *Expert Rev Pharmacoecon Outcomes Res.* 2010;10, 441–451.
9. Hancock, R.E., Speert, D.P. Antibiotic resistance in *Pseudomonas aeruginosa*: mechanisms and impact on treatment. *Drug Resist Update.* 2000; 3, 247–255.
10. Breidenstein, E.B., de la Fuente-Nunez, C., Hancock, R.E. *Pseudomonas aeruginosa*: all roads lead to resistance. *Trends Microbiol.* 2011; 19, 419–426.
11. Drenkard, E., Antimicrobial resistance of *Pseudomonas aeruginosa* biofilms. *Microbes Infect.* 2003; 5, 1213–1219.
12. Mulcahy, L.R., Burns, J.L., Lory, S., Lewis, K. Emergence of *Pseudomonas aeruginosa* strains producing high levels of persister cells in patients with cystic fibrosis. *J Bacteriol.* 2010; 192: 6191–6199.
13. Tenover FC. Mechanisms of antimicrobial resistance in bacteria. *Am J Infect Control* 2006; 34(5):10.
14. Strateva T, Yordanov D. *Pseudomonas aeruginosa* a phenomenon of bacteria resistance. *J Med Microb* 2009;58: 1133-48.
15. Morales E, Cots F, Sala M, Comas M, Belvis F, Riu M, *et al.* Hospital costs of nosocomial multi-drug resistant *Pseudomonas aeruginosa* acquisition. *BMC Health Serv Res* 2012; 12: 122.
16. Falagas ME, Koletsi PK, Bliziotis IA. The diversity of definitions of multidrug-resistant (MDR) and pandrug-resistant (PDR) *Acinetobacter baumannii* and *Pseudomonas aeruginosa*. *J Med Microbiol* 2006; 55: 1619-29.
17. Obritsch MD, Fish DN, MacLaren R, Jung R. Nosocomial infections due to multidrug-resistant *Pseudomonas aeruginosa*: epidemiology and treatment options. *Pharmacotherapy* 2005; 25 (10): 1353-64.
18. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, *et al.* Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012; 18 (3): 268-81
19. Cheesbrough M (ed) (2000) *Pseudomonas* and related organisms; Biochemical test to identify bacteria; Antimicrobial susceptibility testing. In: *District Laboratory Practice in Tropical Countries Part II*. Low price edition 2000. Cambridge University Press, Cambridge, pp. 193–194; 63–70; 132–143.

20. CLSI. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Third Informational Supplement. CLSI document M100–S23. Wayne, PA: Clinical and Laboratory Standards Institute; 2013.
21. Isichei-Ukah OB, Enabulele, OI. Prevalence and Antimicrobial Resistance of *Pseudomonas aeruginosa* recovered From Environmental And Clinical Sources In Benin City. *Ife Journal of Science*; 2018 20(3):1734-1738.
22. Davane M, Suryawanshi N, Pichare A, Nagoba BS. *Pseudomonas aeruginosa* from hospital environment. *J Microbiol Infect Dis*. 2014;4:01. doi:10.5799/ahinjs.02.2014.01.0124
23. Kateete DP, Nakanjako R, Namugenyi J, Erume J, Joloba ML, Najjuka CF. Carbapenem resistant *Pseudomonas aeruginosa* and *Acinetobacter baumannii* at Mulago Hospital in Kampala, Uganda (2007–2009). *Springerplus*. 2016;5(1):1–11. doi:10.1186/s40064-016-2986-7
24. Uc-Cachón AH, Gracida-Osorno C, Luna-Chi IG, Jiménez- Guillermo JG, Molina-Salinas GM. High prevalence of antimicrobial resistance among gram-negative isolated Bacilli in intensive care units at a tertiary-care hospital in Yucatan Mexico. *Medicina*. 2019;55(9):588. doi:10.3390/medicina55090588
25. Kumari M, Khurana S, Bhardwaj N, Malhotra R, Mathur P. Pathogen burden & associated antibiogram of *Pseudomonas* spp. in a tertiary care hospital of India. *Indian J Med Res*. 2019; 149(2):295. doi:10.4103/ijmr.IJMR_14_18
26. Prakash V, Mishra PP, Walia PA, Dhawan S, Kumar A. Increasing incidence of multidrug resistant *Pseudomonas aeruginosa* in inpatients of a tertiary care hospital. *Int J Res Med Sci*. 2014;2(4):1302-1306
27. Abd El-Baky RM, Masoud SM, Mohamed DS, Waly NG, Shafik EA, Mohareb DA et al. Prevalence and some possible mechanisms of colistin resistance among multidrug-resistant and extensively drug-resistant *Pseudomonas aeruginosa*. *Infect Drug Resist*. 2020; 13:323. Doi:10.2147/IDR.S238811
28. Saderi H, Owlia P. Detection of multidrug resistant (MDR) and extremely drug resistant (XDR) *P. aeruginosa* isolated from patients in Tehran, Iran. *Iranian j Pathol*. 2015; 10(4):265.
29. World Health Organization. Global Antimicrobial Resistance Surveillance System (GLASS): the detection and reporting of colistin resistance; 2018. Available from: <https://apps.who.int/iris/handle/10665/277175/> Accessed May 31, 2023.