

## Original Research Article

### **Effects of different vesicular - arbuscular mycorrhizal (VAM) fungi on the seedling growth of *Tecomella undulata***

#### **Abstract**

*Tecomella undulata* (Sm.) Seem. (family Bignoniaceae) is a tree found in desert parts of India, and Arabia. The cultivation of high-quality seedlings is vital for establishing a successful plantation and the contribution of arbuscular mycorrhizal fungi (AMF) in the soil plays a vital role in the nursery phase. A study conducted during the periods of 2021-22 and 2022-23 scrutinised the effect of four distinct species of *Glomus* spp. (*G. mosseae*, *G. intraradices*, *G. fasciculatum* and *Glomus hoi*) on *Tecomella undulata* seedlings in AMF-inoculated soil. The inoculation involved applying of 400-500 sporocarps/kg of soil during sowing and the evaluation encompassed growth and survival parameters. Results revealed that among the 4 *Glomus* species, soil inoculated with *Glomus fasciculatum* at 180 days after sowing, significantly increased the root colonization percentage with (50.69 and 46.71 %) in year (2021-22) and (2022-23) compared to the uninoculated control. Additionally, seedlings exhibited significantly higher no. of spores per 100g of soil, when treated with (*Glomus fasciculatum*) inoculated soil, which was statistically at par with soil inoculated with (*Glomus intraradices*) at (103.67 and 98.65). During both the experimental year, in terms of germination percentage, root length and root: shoot ratio were found non-significantly with (44.45 %), root length at 360 DAS, (26.67 and 25.74) respectively, root shoot ratio at 360 days after sowing. While, plant survival percentage (30.63 and 25.41%) at 90 DAS. Whereas, at 360 DAS, collar diameter (6.14 and 10.07 mm), shoot length (33.36 and 32.19 cm), plant biomass (8.31 and 8.05 g), leaf area (196.29 and 185.46 cm<sup>2</sup>) was observed significantly higher in treatment T3 (*Glomus fasciculatum*) which was statistically at par with treatment T1 (*G. intraradices*). It is concluded from study that *Glomus fasciculatum* were found best for growth and survival of seedlings followed by *Glomus intraradices* as compare to control.

Key words: Seedlings, *Glomus fasciculatum*, Arbuscular mycorrhizal fungi, *G. intraradices*

#### **1. INTRODUCTION**

*Tecomella undulata* (Sm.) Seem., belonging to the Bignoniaceae family, is a tree of significant economic and pharmaceutical importance found in the arid regions. The plant is renowned for its diverse therapeutic effects, including hepatoprotective, antibacterial,

antimicrobial, antifungal and anti-inflammatory activities. Symbiotic microorganisms in the mycorrhiza play a vital role in plant adaptation and survival. Mycorrhizal interactions enhance the overall absorption capacity of plant roots through morphological and physiological modifications. These modifications involve an increase in the absorption surface area, exploration of a larger soil area (as the fungus acts as an extension of the root), prolonged lifespan of absorbing roots, improved utilization of low-availability nutrients, and enhanced retention and storage of soluble nutrients. (Dehne, 1982). It reduces interactions with soil colloids and minimizes leaching losses. (Muchovej, 2002). AM fungi confer a distinct advantage to their host compared to non-host species by offering nutrients and defending against various pathogenic organisms (Kothamasiet *al.*, 2001). Among these, the primary significance lies in nutrient absorption, preventing nutrient leaching, contributing to soil structure, and serving as hyphal highways for the dispersion of bacteria (Marcel *et al.*, 2009). Mycorrhizal fungi play a role in inhibiting the growth of pathogens. However, the establishment of mycorrhizal associations is influenced by various factors such as soil moisture, soil pH, organic carbon content, soil phosphorus, and soil nitrogen. These factors affect both root colonization by mycorrhizal fungi and the density of spores. Additionally, seasonal variations impact mycorrhizal associations, while the use of modern pesticides and foliar fungicides can reduce the incidence of mycorrhizal presence. The natural occurrence of VAM (vesicular-arbuscular mycorrhizal) association in desert plants is crucial in arid zone forestry. These associations play a vital role by offering not only additional mineral support to the plants but also aiding in their survival under challenging climatic conditions. Prasad and Mertia (2005) proposed that the biological activity in the rhizosphere of trees and the presence of VAM fungi associated with tree roots could serve as criteria for selecting agroforestry trees in arid ecosystems. Limited research has been conducted on the application of arbuscular mycorrhizal fungi (AMF) in cultivating seedlings of Rohida. Consequently, the current study was designed with the aim of identifying a suitable AMF that can be artificially introduced into the soil prior to sowing.

## **2. MATERIALS AND METHODS**

The investigation carried out in the nursery of the Forestry Department, CCS Haryana Agricultural University, located in Hisar (latitude 29.10° N, longitude 75.46° E, altitude 215 meters above sea level) in the semi-arid region of North-Western India. The soil employed in the experiment was of sandy loam composition. Hisar, situated in Haryana, experiences a semi-arid climate characterized by intense heat and dry, desiccating winds, often

accompanied by high-velocity dust storms in the summer. Pure cultures of four different AMF species, namely *Glomus intraradices*, *Glomus mosseae*, *Glomus fasciculatum* and *Glomus hoi* with control. The samples of four different AM fungi of *Glomus spp.* were obtained from the Department of Plant Pathology, CCS Haryana Agricultural University, Hisar. These samples were then multiplied on pearl millet (*Pennisetum typhoides*) and wheat (*Triticum aestivum*) under controlled conditions within a screen house. The soil and rootlets from the root horizon of plants inoculated with *G.intraradices*, *G.mosseae*, *G.fasciculatum* and *Glomus hoi* were utilized in various treatments before sowing the seeds, as outlined below. The seeds were planted at a depth of 2-3 cm in sterile sandy soil individually inoculated with *G.intraradices*, *G.mosseae*, *G.fasciculatum* and *Glomus hoi* at a rate of 450-500 sporocarps/kg of soil. This was carried out in polythene bags measuring 22.5 x 12.5 cm in the months of august. A control was also included by sowing seeds in uninoculated soil. The experiment was set up in a completely randomized design, with each treatment being replicated five times. Fifteen seedlings per replication were raised and maintained. Growth parameters of the seedlings, mycorrhizal colonization in roots, and sporocarp numbers in the soil were assessed at 90, 180, 270 and 360 days after sowing (DAS). Mycorrhizal colonization was determined using the method outlined by Phillips and Hayman (1970) and sporocarps were quantified following the procedure of Gerdeman and Nicolson. The study was conducted at the Forestry Nursery of CCS, Haryana Agricultural University, Hisar, during the experimental year, (2021-22 and 2022-23). The treatments for soil inoculation with *Glomus spp.* before sowing and the control (uninoculated) AMF were as follows:

T1: *Glomus intraradices*

T2: *Glomus mosseae*

T3: *Glomus fasciculatum*

T4: *Glomus hoi*

T5: Control (uninoculated)

### **3. RESULTS AND DISCUSSION**

#### **3.1 Effect of different AMF on seed germination and seedling growth parameters**

The information provided in Table 1 to 7 discloses the growth parameters, including plant survival %, collar diameter (mm), root length (cm), shoot length (cm), plant biomass (g), leaf area (cm<sup>2</sup>)/ plant and root : shoot ratio for *Tecomella undulata* seedlings at 90, 180, 270, and 360 days after sowing (DAS). These parameters were observed in soil treated with various *Glomus spp.* during the years 2021-2022 and 2022-2023, respectively. The Plant survival % of *Tecomella undulata* seedlings was significantly higher in all the treatments having soil



T1	<i>Glomus intraradices</i>	28.41	25.14	23.12	22.23	26.45	23.25	21.89	19.82
T2	<i>Glomus mosseae</i>	25.28	22.98	21.78	19.64	23.21	20.20	18.63	16.20
T3	<i>Glomus fasciculatum</i>	30.63	27.09	25.84	23.36	28.55	25.41	24.64	22.68
T4	<i>Glomus hoi</i>	27.57	23.12	22.31	20.24	24.49	22.35	19.21	18.31
T5	Control	24.12	20.85	19.29	17.22	22.56	19.12	17.56	15.68
	C.D at 5 %	2.44	2.15	2.03	1.86	2.24	1.98	1.84	1.67

There is a paucity of studies focusing on the selection of a suitable arbuscular mycorrhizal fungus (AMF) for artificial soil inoculation before sowing, especially for the production of robust seedlings with high vigour to enhance establishment under various biotic and abiotic stresses in field conditions. Khaiperet *et al.*, (2023) examined that among of all the treatments, *Glomus mosseae* was found effective for enhancing plant growth parameters and maximizing the biomass of *Melia azedarach* seedlings. Similar observations were observed by Singh and Chugh [10] reported that enhanced survival percentage, root length, shoot length, total biomass, in trees like *Dalbergia sissoo*, *Eucalyptus tereticornis*, *Azadirachta indica*, and *Ailanthus excelsa* when sown in soils containing mycorrhiza compared to non-inoculated soil. Similarly, studies by Gulab *et al.*, (2022) demonstrated that significantly higher plant growth parameters survival percentage, root length, shoot length and plant biomass in *Melia azedarach* in the presence of *G. mosseae*, *G. fasciculatum* as compared to control. Giri *et al.* [14] reported that to enhancing the survival rate, the inoculation of arbuscular mycorrhizal (AM) fungi resulted in improved seedling growth, including plant height and collar diameter. Among the AM fungi treated, *Glomus macrocarpum* exhibited higher efficacy compared to *G. fasciculatum*, both in the nursery seedlings. Additionally, Kaushik *et al.* revealed that *Acacia nilotica* and *Dalbergia sissoo* seedlings exhibited significantly higher biomass, root and shoot length when inoculated with *G. mosseae* compared to the control. Singh *et al.* investigated that percentage of plant survival was elevated in soil treated to arbuscular mycorrhizal fungi (AMF) treatment, potentially attributed to the safeguarding of seedlings from detrimental pathogens facilitated by *Glomus* spp. Khaiperet *et al.*, (2022) reported that the interaction between arbuscular mycorrhizal fungi (AMF) and the roots of trees, which benefits the plant for better germination and growth. dormancy of seeds or a delay in growth can be identified as a disadvantage in a plantation program. Hamidi *et al.* investigated the impact of endomycorrhizae, specifically *Glomus etunicatum*, on cork oak plant (*Quercus suber*) seedlings. They noted that plants inoculated with mycorrhizae exhibited a stimulative influence on the aerial part, aerial part weight, average root length, average fresh weight, and leaf count compared to the control group. Sumana and Bagyaraj (33) carried out an experiment utilizing eight different VAM (vesicular-arbuscular mycorrhizal) fungi. They

aimed to identify the most suitable symbiont for *Dalbergia latifolia* and found that plants inoculated with these fungi displayed increased plant height, stem girth, dry weight, and phosphorus content in comparison to non-inoculated plants. Among the VAM fungi, *Glomus fasciculatum* were particularly effective in enhancing the growth of *D. latifolia*. The presence of AMF in the soil contributes to enhanced survival rates for rohida. The alteration of the soil environment by AMF enveloping the roots could also be a factor contributing to the improved seedling growth, survival rates, more effective root colonization, and increased multiplication.

**Table 2. Effect of arbuscular mycorrhiza fungi on Collar diameter of Rohida in nursery:**

Treatments		Collar diameter (mm)							
		2021-2022				2022-2023			
		90 DAS	180 DAS	270 DAS	360 DAS	90 DAS	180 DAS	270 DAS	360 DAS
T1	<i>Glomus intraradices</i>	5.96	7.29	9.35	10.09	4.98	6.67	8.70	9.38
T2	<i>Glomus mosseae</i>	4.71	5.98	7.45	7.91	3.67	5.03	7.04	7.46
T3	<i>Glomus fasciculatum</i>	6.14	7.96	9.61	10.44	5.93	7.68	9.02	10.07
T4	<i>Glomus hoi</i>	5.21	6.46	8.33	8.92	4.10	6.06	7.96	8.39
T5	Control un-inoculated)	4.09	4.88	6.76	6.93	3.03	4.34	6.33	6.49
	<b>C.D at 5 %</b>	<b>0.47</b>	<b>0.60</b>	<b>0.75</b>	<b>0.80</b>	<b>0.40</b>	<b>0.55</b>	<b>0.70</b>	<b>0.76</b>

**Table 3. Effect of arbuscular mycorrhiza fungi on Root length of Rohida in nursery**

Treatments		Root length (cm)							
		2021-2022				2022-2023			
		90 DAS	180 DAS	270 DAS	360 DAS	90 DAS	180 DAS	270 DAS	360 DAS
T1	<i>Glomus intraradices</i>	15.67	18.67	22.51	25.84	14.57	17.36	21.45	24.08
T2	<i>Glomus mosseae</i>	14.12	17.59	21.79	24.91	13.11	16.62	19.67	23.11
T3	<i>Glomus fasciculatum</i>	16.50	19.33	23.41	26.67	15.92	18.65	22.59	25.74
T4	<i>Glomus hoi</i>	14.33	18.03	22.02	25.13	13.69	17.22	21.03	23.78
T5	Control(uninoculated)	13.89	16.69	20.15	23.79	12.76	16.03	19.13	22.67
	<b>C.D at 5 %</b>	<b>1.33</b>	<b>1.61</b>	<b>1.96</b>	<b>N.S</b>	<b>1.25</b>	<b>1.53</b>	<b>1.85</b>	<b>N.S</b>

**Table 4. Effect of arbuscular mycorrhiza fungi on Shoot length of Rohida in nursery**

		Shoot length (cm)	
		2021-2022	2022-2023

Treatments		<b>90 DAS</b>	<b>180 DAS</b>	<b>270 DAS</b>	<b>360 DAS</b>	<b>90 DAS</b>	<b>180 DAS</b>	<b>270 DAS</b>	<b>360 DAS</b>
T1	<i>Glomus intraradices</i>	23.10	26.33	28.53	32.67	22.17	24.45	26.53	30.38
T2	<i>Glomus mosseae</i>	19.45	22.95	24.65	30.12	19.21	21.69	23.29	28.46
T3	<i>Glomus fasciculatum</i>	25.00	27.12	29.12	33.36	24.13	26.17	28.10	32.19
T4	<i>Glomus hoi</i>	21.20	25.19	27.39	30.55	20.89	22.33	25.45	29.18
T5	Control	18.89	21.55	23.83	29.55	18.16	20.67	22.16	28.33
	C.D at 5 %	1.93	2.20	2.38	2.77	1.88	2.07	2.25	2.64

**Table 5. Effect of arbuscular mycorrhiza fungi on root : shoot ratio of Rohida in nursery.**

Treatments		<b>Root: shoot ratio</b>							
		2021-2022				2022-2023			
		<b>90 DAS</b>	<b>180 DAS</b>	<b>270 DAS</b>	<b>360 DAS</b>	<b>90 DAS</b>	<b>180 DAS</b>	<b>270 DAS</b>	<b>360 DAS</b>
T1	<i>Glomus intraradices</i>	0.68	0.71	0.79	0.79	0.66	0.71	0.81	0.79
T2	<i>Glomus mosseae</i>	0.73	0.77	0.88	0.83	0.68	0.77	0.84	0.81
T3	<i>Glomus fasciculatum</i>	0.66	0.71	0.80	0.80	0.66	0.71	0.80	0.80
T4	<i>Glomus hoi</i>	0.68	0.72	0.80	0.82	0.66	0.77	0.83	0.81
T5	Control (un-inoculated)	0.74	0.77	0.85	0.81	0.70	0.78	0.86	0.80
	C.D at 5 %	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S

**Table 6. Effect of arbuscular mycorrhiza fungi on Plant Biomass of Rohida in nursery**

Treatments		<b>Plant Biomass (g)</b>							
		2021-2022				2022-2023			
		<b>90 DAS</b>	<b>180 DAS</b>	<b>270 DAS</b>	<b>360 DAS</b>	<b>90 DAS</b>	<b>180 DAS</b>	<b>270 DAS</b>	<b>360 DAS</b>
T1	<i>Glomus intraradices</i>	2.15	4.36	6.12	7.58	2.00	4.20	6.00	7.41
T2	<i>Glomus mosseae</i>	1.53	3.92	4.97	5.08	1.15	3.17	4.46	4.93
T3	<i>Glomus fasciculatum</i>	3.26	5.67	7.65	8.31	3.11	5.34	7.27	8.07
T4	<i>Glomus hoi</i>	1.85	4.00	5.71	6.21	1.32	3.95	5.33	6.11
T5	Control (un-inoculated)	1.25	2.58	3.67	4.97	1.10	2.28	3.53	3.93
	C.D at 5 %	<b>0.39</b>	<b>0.48</b>	<b>0.54</b>	<b>0.59</b>	<b>0.35</b>	<b>0.43</b>	<b>0.46</b>	<b>0.57</b>

**Table7. Effect of arbuscular mycorrhiza fungi on Leaf area (cm<sup>2</sup>) per plant of Rohida in nursery**

	<b>Leaf area (cm<sup>2</sup>) / plant</b>

Treatments		2021-2022				2022-2023			
		90 DAS	180 DAS	270 DAS	360 DAS	90 DAS	180 DAS	270 DAS	360 DAS
T1	<i>Glomus intraradices</i>	112.63	153.13	183.76	193.29	108.53	150.37	178.27	182.32
T2	<i>Glomus mosseae</i>	107.56	150.36	177.43	188.45	100.67	146.63	170.36	178.51
T3	<i>Glomus fasciculatum</i>	115.67	157.49	184.79	196.29	112.15	152.39	181.33	185.46
T4	<i>Glomus hoi</i>	110.15	152.27	180.35	191.37	105.46	149.21	173.29	180.67
T5	Control	103.19	148.75	175.13	186.65	97.31	142.16	168.61	172.79
	C.D at 5 %	2.18	2.47	2.82	3.05	2.03	2.41	2.72	2.96

### 3.2 Impact of Soil Application of Different *Glomus* spp. on root colonization and numbers of sporocarp per 100 g of soil

The information provided in Table 8 and Table 9 illustrates the root colonization percentage and the number of sporocarp per 100 g of soil in *Tecomella undulata* seedlings at 90, 180, 270, 360 days after sowing (DAS) in soil treated with various *Glomus* spp. in the periods of 2021-2022 and 2022-2023, respectively. In both years, when each of the 4 *Glomus* species was individually inoculated into the soil before sowing seedlings *G. fasciculatum* consistently resulted in significantly higher root colonization at 180 DAS, with (50.69% and 46.71%) respectively. were closely followed by *G. intraradices* treated soil exhibited root colonization percentages with (42.71% and 39.66 %) at 180 DAS during 2021-22 and 2022-23, respectively. In contrast, no root colonization was observed in the control (uninoculated) soil. Regarding the number of sporocarps per 100 grams of soil, *G. fasciculatum* treated soil exhibited significantly higher values at 360 DAS (103.67 and 98.65) sporocarps per 100 g of soil during 2021-2022 and 2022-2023, respectively. which was followed by *G. intraradices*, treated soil recorded at (94.33 and 90.45) sporocarps per 100 g of soil during 2021-2022 and 2022-2023, respectively at 180 DAS. Similar observations on increase in sporocarp numbers in soil and root colonization in *Dalbergia sissoo* have also been reported by Kumar, *et al.* []. The AMF root colonization and spore density in soil is an important parameter to determine the status of the association and studies have shown that higher colonization rates are often positively correlated with plant growth and nutrition. Tapwalet *al.* (2015) reported maximum growth and germination in *Shorea robusta* on inoculation with arbuscular mycorrhiza fungi. The root exudates of seedlings may also have influenced the better root colonization, faster multiplication of AMF and hence better growth and survival of seedlings.

**Table 8. Effect of different arbuscular mycorrhiza fungi on root colonization % of Rohida**

Treatments		Root colonization %			
		2021-2022		2022-2023	
		90 DAS	180 DAS	90 DAS	180 DAS
T1	<i>Glomus intraradices</i>	36.27	42.71	32.48	39.66
T2	<i>Glomus mosseae</i>	22.45	26.33	18.61	22.49
T3	<i>Glomus fasciculatum</i>	43.56	50.69	40.37	46.71
T4	<i>Glomus hoi</i>	27.67	35.27	24.53	32.33
T5	Control(uninoculated)	00.00	00.00	00.00	00.00
	<b>C.D at 5 %</b>	<b>2.84</b>	<b>3.38</b>	<b>2.54</b>	<b>3.07</b>

**Table 9. Effect of arbuscular mycorrhiza fungi on number of spores of Rohida in nursery:**

Treatments		No. of spores per 100 gm of soil			
		2021-2022		2022-2023	
		90 DAS	180 DAS	90 DAS	180 DAS
T1	<i>Glomus intraradices</i>	82.47	94.33	78.66	90.57
T2	<i>Glomus mosseae</i>	70.39	81.54	64.51	76.51
T3	<i>Glomus fasciculatum</i>	95.61	103.67	91.37	98.65
T4	<i>Glomus hoi</i>	76.55	87.45	72.33	84.36
T5	Control (un-inoculated)	00.00	00.00	00.00	00.00
	<b>C.D at 5 %</b>	<b>5.63</b>	<b>5.75</b>	<b>4.09</b>	<b>5.38</b>

#### 4. CONCLUSION

The effect of different *Glomus* spp. (*Glomus intraradices*, *Glomus mosseae*, *Glomus fasciculatum* and *Glomus hoi*) on the growth characteristics (plant survival %, shoot length, root length, collar diameter, plant biomass, leaf area) and survival of *Tecomella undulata* seedlings was investigated. The results indicated that *G. fasciculatum* exhibited the highest effectiveness compared to the other species. Additionally, *G. fasciculatum* treatment resulted in the highest root colonization of seedlings followed by *G. intraradices* and *G. hoi* treatments. The multiplication rate of *G. fasciculatum* was also found to be the highest, followed by *G. intraradices* and *Glomus hoi*. In conclusion, applying *G. fasciculatum* to the soil before sowing *Tecomella undulata* seeds proved to be highly effective for seedling establishment in nursery conditions.

## REFERENCES

- Singh MK, Chugh RK. Impact of Arbuscular Mycorrhizal fungi on the growth parameters and nutrient content on different tree species. *Indian Journal of Pure Applied Bioscience*. 2019;7(6):244-248.
- Gerdemann, JW, Nicolso TH. Spores of mycorrhizal Eadogone. species extracted from soil by wet sieving and decanting. *Transaction of British Mycology Society*.1963;46:235-244.
- Giri B, Kapoor R. Mukherji KG. Effect of the arbuscular mycorrhizae *Glomus fasciculatum* and *G. macrocarpum* on the growth and nutrient content of *Cassia siamea* in a semi-arid Indian wasteland soil. *New Forests*. 2005;29:63-73.
- Phillips JM, Hayman DS. Improved procedure for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Transactions of British Mycological Society*. 1970;55:158-161.
- Muchovej, R. N . Importance of Mycorrhiza for Agriculture crops. *Journal of Ecology* .2002;97:1139-1150.
- Marcel, G.A., V. Heijden and T.R. Horton Socialism in soil ? The importance of mycorrhizal fungal networks for facilitation in natural ecosystems. *Journal of Ecology*. 2009; 97:11 39-1 150.
- Prasad R, Mertia RS. Dehydrogenase activity and VAM fungi in tree-rhizosphere of agroforestry systems in Indian arid zone. *Agroforestry systems*. 2005;63:219-23.
- Ahlawat KS, Dhanda SK, Chugh R, Mehta SK, Sirohi C, Dalal V, Poonia P. Impact of Different *Glomus* Species on Growth and Survival of *Acacia nilotica* Seedlings. *International Journal of Plant & Soil Science*. 2022; 18:74-81.
- Khaiper M, Dhanda SK, Ahlawat KS, Poonia PK, Kumar A, Verma P, Chugh R, Jangra M. Unlocking the Growth Potential of *Melia azedarach* Seedlings: The Synergistic Impact of *Glomus mosseae* and Pre-sowing Treatments. *Biological Forum – An International Journal*. 2023; 15(8): 371-377.
- Gulab, Verma R, Ahlawat KS, Chugh R, Sirohi C, Mehta SK, Singh C, Khaiper M. Impact of Different Arbuscular Mycorrhizal Fungi on Nutrient Status and Their Uptake by *Melia azedarach* Seedling. *International Journal of Plant & Soil Science*. 2022; 28:89-96.
- Giri B, Kapoor R. Mukherji KG. Effect of the arbuscular mycorrhizae *Glomus fasciculatum* and *G. macrocarpum* on the growth and nutrient content of *Cassia siamea* in a semi-arid Indian wasteland soil. *New Forests*. 2005;29:63-73.
- Tapwal, A., Kumar, R. and Borah, D. Effect of mycorrhizal inoculations on the growth of *Shorea robusta* seedlings. *Nusantara bioscience*. 2015;7(1):1-5
- Hamidi O, Talbi Z, Chliyah M, Touhami AO, Selmaoui K, Benkirane R, Douira A. Effect of Endomycorrhizal Inoculation on the Young Cork Oak Plants (*Quercus suber*) Growth. *Annual Research & Review in Biology*. 2017;13:1-11.
- Sumana DA, Bagyaraj DJ. Selection of efficient VA-mycorrhizal fungi for inoculating Neem. In *Proceedings of the National conference of Mychorrhiza*. Singh Barkatulla University, Bhopal 1999;10.

Khaiper M, Kumar Poonia P, Kumar Dhanda S, Ahlawat KS, Chugh R, Kumar A, Verma P, Jangra M, Kumar Kagra A, Nanda K. Efficacy of Arbuscular Mycorrhizal Inoculations on Tree Species Germination and Growth. *International Journal of Plant & Soil Science*. 2022; 34:930-936.

Dehne, H.W. Interaction between vesicular-arbuscular mycorrhizal fungi and plant pathogens. *Phytopath*. 1982; 72(8): 115-118.

Kothamasi, D., R.C. Kubad and C.R. Babu. Arbuscular mycorrhizal inplant survival strategies. *Tropical Ecology*. 2001; 42(1):1-13.

UNDER PEER REVIEW

UNDER PEER REVIEW