

POTENTIAL CONTRIBUTION OF EXPOSURE TO INSECTICIDE USED IN NIGERIA TO THE DEVELOPMENT OF DYSLIPIDAEMIA AND HORMONAL DISORDER IN MALE AND FEMALE ALBINO WISTAR RATS

Abstract

Background: The use of household insecticides for the eradication of insects especially mosquitoes in Nigeria is increasing. These insecticides are used without consideration of their adverse effect on human health.

Aim: This study was aimed at accessing the effect of common household insecticides used in Nigeria on male reproductive hormones of Wistar rats.

Methodology: Thirty male Wistar rats were divided into five groups of six each and kept in different rooms. Rats in group 1 were exposed to Rambo, those in group 2 were exposed to Mortein, those in group 3 were exposed to Raid, those in group 4 were exposed to Baygon while those in group 5 were not exposed to any insecticide and served as the control group. The exposure was done twice a day via inhalation route. Throughout the experiment, animals were fed *ad libitum* with standard feed and drinking water. After twenty-one days of exposure, they were sacrificed after an overnight fast under diethyl ether as anesthesia. Blood samples were collected by cardiac puncture. Follicle stimulating hormone (FSH), luteinizing hormone (LH), and testosterone levels were determined using enzyme-linked immunosorbent assay (ELISA).

Results: Results showed that insecticides caused an increase in FSH but decrease in LH and testosterone levels significantly when compared to those in control animals at $P < 0.05$.

Conclusion: The results indicate that prolonged exposure to common household insecticides used

Keywords: Albino wistar rat, reproductive and lipidemic effect, Raid, insecticide

1.0 INTRODUCTION

Excessive use of various chemicals including insecticides has become a public health concern. The Use of insecticides and other organophosphates is one of the major ways through which manufacturing workers and farmers are exposed to toxicants and this has impact on the ecosystem and public health [1]. Insecticides are natural or man-made preparation that is used to kill or otherwise control insects such as mosquitoes, cockroaches, bees, wasp, and bedbug and many others.

They can be grouped into two major classes: systemic insecticides, which are known to have residual or long-term activity and secondly contact insecticides, that has no residual activity. They are in form of sprays, dust, gel, baits, smokers, fumigant and powders [2]. A common active chemical in insecticides are pyrethroids, derivatives of pyrethrins, natural substances obtained from the flowers of pyrethrum species ([2 a and 2b]. The most common active ingredient in insecticide are synergist, carbamate, whose common name is propoxur, pyrethrin (or synthetic pyrethroids), D-trans-allevrin, permethrin, tetramethrin, deltamethrin, cyfluthrin, imiprothrin, chlorpyrifos, Diazinon, Malathion, Silica gel, Boric acid, Arsenicals, paradichlorobenzene, Naphthalene, N,N-diethyl meta-toluamide (3Deet), Dimethylphthalate [2b]. The mechanism of action of insecticide can be important in understanding whether an insecticide will be toxic to unrelated species, such as fish, birds and mammals. In terms of brand, insecticides' may be repellent or non-repellent ([3]. Domestic insecticides are known to be quite different from non-insecticidal repellents, which only keep away insects.

In Nigeria, systemic insecticides are the most common and they include; Rambo manufactured by Gongoin and Co, Mortein manufactured by Reckitt Benckiser, Raid and Baygon both manufactured by S.C. Johnson and Co. The common household Insecticides are distinct from non-insecticidal repellents, which only repel but do not kill [4]. However common household insecticides used in Nigeria are systemic in action and these include Rambo produced by Gongoin and Co, Mortein produced by Reckitt Benckiser, Raid and Baygon both produced by S.C. Johnson Co. [4]

Mosquitoes and man cohabit in most homes in Nigeria and thus it becomes a herculean task to completely eliminate mosquitoes in many homes in our country. The resurgence of mosquitoes few minutes/hours after application of insecticide show that either the insecticide is not effective or it is a sign of resistance of the mosquitoes to the applied insecticide. This

confirms that mosquitoes cannot be completely eradicated in Nigeria. This limitation of the insecticide is not known to many and can lead to abuse or indiscriminate use of insecticide in most homes without consideration of the negative effect on their health.

Lipids are class of compounds that are soluble in organic solvent and nearly insoluble in water. They have also been described as naturally occurring esters of glycerol and fatty acids which are solid at room temperature as opposed to oils which are similar esters but liquid at room temperature [5]. Structurally lipids consist of heterogeneous group of compounds soluble in non-polar solvents such as methanol, ether, chloroform and benzene. Hydrolysis of lipids yields fatty acids or complex alcohols that may combine with the fatty acids to form esters. Some lipids are more complex by the possession of other groups like phosphoric, amine, sugar, or sulphate groups which confers *amphiphatic* character on them. Lipids are ubiquitous in the body tissue, playing important role in virtually all the physiological processes of the body which include, lipids serves as hormone or hormone precursors, aids digestions, facilitates energy storage, provides metabolic fuels, acts as functional and structural components in cell membrane, and also forms insulation to allow nerve conduction or prevent heat loss [6] Simple, Compound and Derived derivatives are the three majors classes of lipid fraction. Each of the fractions has a specific physiological and metabolic role in the body, the alteration of the equilibrium concentration of these lipid fraction may precipitate a pathological disorder in the body.

The reproductive hormones are essential physiological requirements for procreation and they include; Follicle stimulating hormone (FSH). Leuteinizing hormone (LH) Oestrogen and Progesterone, each of them contributing specific role in human in procreation. LH and FSH are gonadotrophins synthesized by the pituitary gland [7] but released by the Gonadotropin releasing hormone (GnRH) a product of the hypothalamus. Testosterone a

reproductive hormone is a product of the extra tubular Leydig cells regulated by LH, a glycoprotein synthesized by the pituitary [7]. FSH synthesized by the pituitary is another gonadotropic hormone that controls sperm production by the regulation of the activities of the both the germinal epithelium and Sertoli cells [8]. Physiological concentration of these hormones is under the control of hypothalamic pituitary gonadal axis [9]. Testosterone improves sperm motility and epididymis function. Oestrogen and Progesterone are female sex hormone which act synergistically to regulate the female accessory sex organs and female secondary sex characteristics, the menstrual cycle, breast, uterine growth and in the maintenance of pregnancy [9].

These hormones are produced by the ovarian follicles and the corpus luteum and during pregnancy by the placenta.

Combined physiological contribution and function of FSH and LH is required for the synthesis of Testosterone in males and Ovarian hormones (oestrogen and progesterone) in female which are essential and inevitable requirements that facilitates fertility [10]

Human Chorionic gonadotropin (hCG) synthesized in the syncytiotrophoblast cells of the placenta is a glycoprotein containing a protein core with branched carbohydrate side chains which usually terminate with sialic acid. hCG is composed of two non-identical, non-covalently bound glycoprotein subunits, the alpha (α) and the beta (β) subunits. [11]. The chorion of the developing placenta begins to secrete hCG shortly after implantation of the fertilized egg and its concentration rises steadily in plasma and urine from the first few days after conception until the tenth or twelfth week of pregnancy [12,13] ([12, 13]. The detection of

hCG in urine or serum is a biochemical confirmation of establishing pregnancy as early as 7 to 10 day post conception [13]. The action of hCG is similar to that of LH because it also stimulates the corpus luteum to produce progesterone. The aim of this study is to evaluate the effect of the common household insecticide used in Nigeria on the Lipids and reproductive hormones in male and female albino wister rat.

MATERIALS AND METHODS

2.1 Collection of Insecticides

Raid (Liquid brand) insecticide was purchased directly from a super market in Ile-Ife. It was preserved at room temperature for use during the experiment.

2.2 Ethical Approval.

Ethical approval was obtained from Animal Care and Use Research Ethics Committee (ACUREC), of Obafemi Awolowo University Ile-Ife. (OAU),

2.3 Animal Selection

Healthy male and female albino rats of the Wistar strain (*Rattus norvegicus*), with an average weight of 110 ± 20 g, obtained from the Animal Breeding House of the Obafemi Awolowo University (OAU), Ile-Ife were acclimatized for one week prior to the commencement of the project. The rats were restricted to a clean quiet, well ventilated and temperature controlled experimental animal cages for 12/12 hours except during the period

of exposure to the insecticide. The experimental work was performed in accordance with the guidance for care and use of laboratory animals. The median lethal dose (LD50; 25mg/kg body weight) for commercial formulations of Raid, (a domestic insecticide) on wister albino rats from previous research was adopted for this study. They were randomly divided into five groups of six rats ,eachand kept in different rooms.

Twenty male and twenty female wistar rats weighing between 109 and 133 kg were used for this study. They were randomly divided into five groups and each group contained eight animals acclimatized for seven (7) days to laboratory conditions before the commencement of the experiment. Prior to and during the experiment, the animals were acclimatized and the male were separated from the females and both were fed *ad libitum* with standard feed and drinking water in the clean cages placed in well-ventilated housing conditions (under humid tropical conditions) throughout the experiment. The animals were given humane care according to the criteria outlined in the 'Guide for the Care and Use of Laboratory Animals' prepared by the National Academy of Science and published by the National Institute of Health.

The rats in group 1 were exposed once to the insecticide and tagged acute group. The rats in group 2 were exposed trice to the insecticide at interval of 48 hours and was tagged sub-chronic group. The third groups of the rats were exposed four times to the insecticide at interval of 48 hours and was labelled the chronic group. The rats in the fourth group were exposed four times to the insecticide at the interval of 48 hours but were left for two weeks undisturbed except for feeding. This group was labelled post exposure group and the animals in the group were sacrificed for sample collection 7 days post exposure. While the rats in group 5 were not exposed at all. (The control group). All animals received humane care in compliance with the guidelines of the Obafemi Awolowo University Ile-Ife Animal Care and Use Research Ethics Committee (ACUREC). The exposure was done twice daily via inhalation route. During the duration of the experiment the albino rats were fed *ad libitum* with standard feed and hygienic water. the albino rats were fed *ad libitum* with standard feed and hygienic water. The rats in each group were sacrificed under diethyl ether as anaesthesia after an overnight fast

Blood was collected from each rat in the group through cardiac puncture with the aid of a needle and syringe. Two milliliters (2ml) of blood was dispensed into K-EDTA anticoagulant bottle for lipid estimation while 3ml was dispensed in to plain bottle for the hormonal assay.

2.5 Assay of Reproductive hormones

The serum progesterone, testosterone beta hCG, FSH, and LH and were measured with the aid of enzyme-linked immunosorbent assay (ELISA) according to the methods described by [14]Hidayat and Patricia, (2021)

Assay of lipid profile.

Plasma cholesterol and triglyceride were assayed using Enzymatic method described by [15]Li-Hua, 2019 and [16] while the plasma HDL, LDL and VLDL- cholesterol were measured by[17,18, 19]

UNDER PEER REVIEW

3.0 RESULTS

Table 1 The distribution of weight change among the Albino rats used for this experimental research before and after exposure to the domestic insecticide is shown in Table 1. Among the exposed group, the sub chronic had the highest weight loss (122%) while the acute exposed group had the lowest weight loss (52.1%) when compared with the unexposed control group. The unexposed control group gained weight significantly when compared to all the exposed groups. The comparative loss of weight by the exposed albino rats might be due to the combined negative effects of the insecticide which might have been precipitated through insecticide inhalation causing irritation. loss of appetite, distortion and probably inactivation of the digestive system.

Table1: Mean Initial and Final Weight of the all the Exposed and Unexposed Albino rats to Insecticide (Groups 1-5)

Variables	Grp 1 (Acute exposure)	Grp2(Sub-chronic exposure)	Grp3 (Chronic Exposure)	Grp4 (Post chronic Exposure)	Grp 5 (Control)	p-value
Initial weight(Kg)	109.53±5.4	135. ±895.2	92.20±6.0	136.75±4.5	133.55±5.2	P<0.05
Final Weight(Kg)	166.60±8.8	300.86±5.8	168.24±7.9	214.01±9.7	525.20±4.20	P<0.05
Change in weight(Kg)	57.07±3.5	164.97±4.7	76.04±4.9	77.26± 4.8	392.05±7.5	P<0.05
% Chang in weight (Kg)	52.10	122.20	82.47	56.50	293.56	P<0.05

Table 2 shows the Mean \pm S.E.M plasma lipid profile in both the exposed and non –exposed Albino rats to the domestic insecticide.

All the lipid profile indices in the exposed group (groups 1 to 4) were not statistically different ($P>0.05$) when compared to the unexposed groups (group5). Comparison of the lipid profile indices among different groups of exposed albino rats (groups 1 to 4) showed no statistically significant difference ($P>0.05$).

Table 2 Lipid profile in the different groups of exposed and non –exposed of albino rats to insecticide

Variables	Grp 1 (Acute exposure)	Grp2(Sub-chronic exposure)	Grp3 (Chronic Exposure)	Grp4 (Post chronic Exposure)	Grp 5 (Control)	p-value
TG (mmol/L)	1.35 \pm 0.13	0.93 \pm 0.10	1.13 \pm 0.10	1.20 \pm 0.22	1.15 \pm 0.13	$P>0.05$
TCHOL (mmol/L)	2.35 \pm 0.16	2.58 \pm 0.17	3.01 \pm 0.13 *	2.8 \pm 0.12	2.65 \pm 0.12	$P>.05$
LDLC (mmol/L)	0.70 \pm 0.08	1.18 \pm 0.35	0.78 \pm 0.33	0.41 \pm 0.17	0.80 \pm 0.23	$P>0.05$
HDLC (mmol/L)	1.20 \pm 0.07	0.83 \pm 0.10*	1.78 \pm 0.25 *	1.9 \pm 0.24	1.33 \pm 0.15	$P>0.05$

The hormonal profiles in the different groups of the exposed and non- exposed female albino rats are shown in Table 4

There was no statistical significant difference ($P>0.05$) in the FSH and progesterone concentrations values between the control and exposed groups. However there was statistically significant difference ($P<0.05$) in LH, Oestrogen and β - hCG values between the control and the exposed groups. Comparison of the reproductive hormones among different groups of exposed female albino rats (1 to 4) showed no statistically significant difference ($P>0.05$) for all the reproductive hormones

Table 3 : Reproductive hormones profile in the different groups of exposed and non –exposed female albino rats

Variables	Grp 1 (Acute exposure) (Mean \pm SD)	Grp2(Sub-chronic exposure) (Mean \pm SD)	Grp3 (Chronic Exposure) (Mean \pm SD)	Grp4 (Post chronic Exposure) (Mean \pm SD)	Grp 5 (Control) (Mean \pm SD)	p-value

LH	1.2±0.10	0.83±0.10	1.56±0.12	1.7±0.82	1.62±0.10	P<0.05
FSH	0.51±0.10	0.52±0.01	0.54±0.01	0.53±0.82	0.51±0.10	p>0.05
PROG	23.3±0.32	9.03±0.17	16.25±0.96	5.00±0.82	4.48±0.10	p>0.05
OEST	40.6±0.36	42.702±0.27	32.50±1.29	47.30±0.90	62.72±0.82	P<0.001
β- HCG	6.5±0.08	1.302±0.19	2.48±0.17	2.75±0.06	4.12±0.10	P<0.001

The mean ± S.E.M. of the reproductive hormones in the exposed male group only were presented in Table 4. Comparison of the reproductive hormones among different groups of exposed male albino rats (groups 1 to 5) showed no statistically significant difference (P>0.05) for all the reproductive hormones except for the LH which was significantly reduced (P>0.05) when compared with the non-exposed control group

Table 4: Reproductive hormone level profile in exposed and non- exposed ,male albino rats to insecticide.

Variables	Grp 1 (Acute exposure) (Mean ±SD)	Grp2(Sub-chronic exposure) (Mean ±SD)	Grp3 (Chronic Exposure) (Mean ±SD)	Grp4 (Post chronic Exposure) (Mean ±SD)	Grp 5 (Control) (Mean ±SD)	p-value
LH	1.71±0.08	1.70±0.08	1.55±0.13	1.15±0.13	1.90±0.08	P<0.05
FSH	0.60±0.07	0.56±0.01	0.48±0.01	0.54±0.02	0.56±0.08	p>0.05
Testosterone	0.30±0.08	2.15±0.13	0.58±0.01	9.68±0.17	8.1±0.08	p>0.05

4.0 DISCUSSION

Exposure to different insecticide has been reported to cause adverse effects on the physiological equilibrium of the biochemical and the biological systems of different tissues [20] Exposure to Raid, a domestic insecticide significantly induced weight loss in the exposed rats relative to the un-exposed group. (Table1). The weight loss by the exposed rats might be due to the combined negative effects of the insecticide which might have been precipitated through insecticide inhalation causing irritation loss of appetite and probably inactivation of the digestive system

The holistic values obtained for the lipid and reproductive hormones suggests that exposure to the insecticide has no significant adverse effect on the hormonal and lipid indices among the exposed albino rats (Table 2&4). Our study showed no statistically significant difference ($P > 0.05$) between the values obtained for the lipid indices among the exposed groups compared to the controls therefore this study shows no correlation between the duration of exposure and the lipid indices. However, this is in contrast with the reports of [21] and [22] who reported significant ($p < 0.05$) increase in the levels of lipids in the exposed albino wistar rats in relation to the duration of exposure to the raid insecticide. We observed significant decrease ($P < 0.05$) in the values of LH, Oestrogen and β -hCG in the the female exposed groups only than the in the unexposed female rats but the values were within the reference range. This result agrees with [23] who reported decreased levels of the reproductive hormones among exposed wistar albino rats.

In contrast, the values obtained for the FSH and progesterone concentrations among the female rats in this study showed no significant difference ($P > 0.05$) between the exposed wistar rats and the control group and they are within the reference values. This result is at variance with the findings of [24] who reported significant concurrent reduction in the gonadotropic hormone, oestradiol, and progesterone levels in insecticide treated rats.

In this study, we also obtained a significantly ($P < 0.05$) reduced LH and a non-significant difference ($p > 0.05$) in the testosterone value among the male albino rats compared to the controls male controls, however the values were within the reference range [25]. This is in contrast with [26] who reported significant decrease in LH, FSH and testosterone levels between the exposed and non-exposed control male albino rats.

Conclusion

We conclude from the findings of this study that exposure to insecticides have no significant effect on the lipids and reproductive hormones concentrations. The results generated from this research further indicate that exposure to the common domestic insecticide used in Nigeria is not associated with dyslipidemia and hormonal disorder. However, cohabitation with the aerosol of the insecticides in a fumigated room should be discouraged.

CONSENT:

This is not applicable.

ETHICAL APPROVAL: Ethical permission was collected and preserved by the author(s).

Limitations: The authors could not assay additional markers due to paucity of fund.

REFERENCES

1. Pathak, V. M., Verma, V. K., Rawat, B. S., Kaur, B., Babu, N., Sharma, A., Dewali, S., Yadav, M., Kumari, R., Singh, S., Mohapatra, A., Pandey, V., Rana, N., & Cunill, J. M. (2022). Current status of pesticide effects on environment, human health and its eco-friendly management as bioremediation: A comprehensive review. *Frontiers in Microbiology*, *13*, 962619. <https://doi.org/10.3389/fmicb.2022.962619>
2. Rasheed S L, E Rasheed A S. (2017) Formulation of Essential Oil Pesticides Technology and their Application. *Agri Res & Tech: Open Access J.* 9(2): 555759. DOI: 10.19080/ARTOAJ.2017.09.555759
2. Sudo, M, Yamanaka, T, Miyai, S. Quantifying pesticide efficacy from multiple field trials. *Population Ecology*. 2019; 61: 450–456. <https://doi.org/10.1002/1438-390X.12019>

3. Hugh Smith, Adam Dale, and Julien Beuzelin (2022) understanding insecticide modes of action and resistance management in florida

horticulture. <https://edis.ifas.ufl.edu> for the currently supported version of this publication. <https://edis.ifas.ufl.edu/publication/IN1379>

4. Augustine I. Airaodion^{1*}, Ada C. Ngwogu², Anthony U. Megwas³,

John A. Ekenjoku⁴ and Kenneth O. Ngwogu (2019) *International Journal of Research and Reports in Gynaecology* 2(1): 1-8, 2019; Article no.IJRRGY.55113

5. Saha, S.K., Pathak, NN. (2021), Lipids In: Fundamentals of Animal Nutrition, springer,

Singapore. https://doi.org/10.1007/978-981-15-9125-9_8

6. Guo R, Chen Y, Borgard H, Jijiwa M, Nasu M, He M, Deng Y. The Function and Mechanism of Lipid Molecules and Their Roles in The Diagnosis and Prognosis of Breast Cancer. *Molecules*. 2020 Oct 21;25(20):4864. doi: 10.3390/molecules25204864. PMID: 33096860; PMCID: PMC7588012.

7. Marsh, C. (Ed.). (2021). Reproductive Hormones. Intech Open. Doi: 10.5772/intechopen.91499.

8. Oduwole, O. O., Peltoketo, H., & Huhtaniemi, I. T. (2018). Role of Follicle-Stimulating Hormone in Spermatogenesis. *Frontiers in Endocrinology*, 9, 427599. <https://doi.org/10.3389/fendo.2018.00763>.

9. Hanlon, C., Takeshima, K., & Bédécarrats, G. Y. (2021). Changes in the Control of the Hypothalamic-Pituitary Gonadal Axis Across Three Differentially Selected Strains of Laying Hens (*Gallus gallus domesticus*). *Frontiers in Physiology*, 12, 651491. <https://doi.org/10.3389/fphys.2021.651491>.

10.Nassar GN, Leslie SW. Physiology, Testosterone. [Updated 2023 Jan 2]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK526128/>

11.Betz D, Fane K. Human Chorionic Gonadotropin. [Updated 2023 Aug 14]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK532950/>.

12.Makrigiannakis, A., Vrekoussis, T., Zoumakis, E., Kalantaridou, S. N., & Jeschke, U. (2017). The Role of HCG in Implantation: A Mini-Review of Molecular and Clinical Evidence. *International Journal of Molecular Sciences*, 18(6). <https://doi.org/10.3390/ijms18061305>

13.Oyatogun, O., Sandhu, M., Barata-Kirby, S., Tuller, E., & Schust, D. J. (2021). A rational diagnostic approach to the “phantom hCG” and other clinical scenarios in which a patient is thought to be pregnant but is not. *Therapeutic Advances in Reproductive Health*. <https://doi.org/10.1177/26334941211016412>

14.Hidayat, R., & Patricia Wulandari. (2021). Enzyme Linked Immunosorbent Assay (ELISA) Technique Guideline. *BioscientiaMedicina : Journal of Biomedicine and Translational Research*, 5(5), 447-453. <https://doi.org/10.32539/bsm.v5i5.228>

15.Li-Hua Li Ewelina P. Dutkiewicz, Ying-Chen Huang, Hsin-Bai Zhou, Cheng-Chih Hsu. (2019).Analytical methods for cholesterol quantification. *Journal of Food and Drug Analysis*, {27},{2},{375-386}. <https://doi.org/10.1016/j.jfda.2018.09.001>

16.Kawano M, Hokazono E, Osawa S,(2019) A novel assay for triglycerides using glycerol dehydrogenase and a water-soluble formazan dye, WST-8. *Annals of Clinical Biochemistry*. 56(4):442-449. doi:10.1177/0004563219830715

17. Friedwald, Levy & Fredrickson equation (1972).
18. Hafiane, A., & Genest, J. (2015). High density lipoproteins: Measurement techniques and potential biomarkers of cardiovascular risk. *BBA Clinical*, 3, 175-188. <https://doi.org/10.1016/j.bbacli.2015.01.005>.
19. Islam SMT, Osa-Andrews B, Jones PM, Muthukumar AR, Hashim I, Cao J. Methods of Low-Density Lipoprotein-Cholesterol Measurement: Analytical and Clinical Applications. *EJIFCC*. 2022 Dec 12;33(4):282-294. PMID: 36605300; PMCID: PMC9768618.
20. Pathak, V. M., Verma, V. K., Rawat, B. S., Kaur, B., Babu, N., Sharma, A., Dewali, S., Yadav, M., Kumari, R., Singh, S., Mohapatra, A., Pandey, V., Rana, N., & Cunill, J. M. (2022). Current status of pesticide effects on environment, human health and its eco-friendly management as bioremediation: A comprehensive review. *Frontiers in Microbiology*, 13, 962619. <https://doi.org/10.3389/fmicb.2022.962619>
21. Yan TL, Zhu XJ, Du HH, Ding XW, Niu DS, Li J. (2022) [Relationship between pesticide exposure and lipid metabolism in population]. *Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi*. 2022 Jan 20;40(1):24-27. Chinese. doi: 10.3760/cma.j.cn121094-20210126-00053.
22. Pothu, U. K., Thammisetty, A. K., & Nelakuditi, L. K. (2019). Evaluation of cholinesterase and lipid profile levels in chronic pesticide exposed persons. *Journal of Family Medicine and Primary Care*, 8(6), 2073-2078. https://doi.org/10.4103/jfmpe.jfmpe_239_19
23. Nguyen HT, Polimati H, Annam SSP, Okello E, Thai QM, et al. (2022) Lobaric acid prevents the adverse effects of tetramethrin on the estrous cycle of female albino Wistar rats. *PLOS ONE* 17(7): e0269983. <https://doi.org/10.1371/journal.pone.0269983>

24. Ghosh R, Banerjee B, Das T, Jana K, Choudhury SM. Antigonadal and endocrine-disrupting activities of lambda cyhalothrin in female rats and its attenuation by taurine. *Toxicology and Industrial Health*. 2018;34(3):146-157. doi:10.1177/0748233717742291
25. Suresh C. Joshi, Bhawna Bansal & Nakuleshwar D. Jasuja (2011) Evaluation of reproductive and developmental toxicity of cypermethrin in male albino rats, *Toxicological & Environmental Chemistry*, 93:3, 593-602, DOI: 10.1080/02772248.2010.537441
26. Airaodion, augustinei. airaodion, ada c. ngwogu2, anthony u. megwas
john a. ekenjoku4 and kenneth o. ngwogu. (2019) effect of common household insecticides used in Nigeria on rat male reproductive hormones. *international journal of research and reports ingynaecology* 2(1): 1-8
27. Mazaheri, F., Aliabad, K. K., Kalantar, S. M., Ziya, N., Khoradmehr, A., & Anvari, M. (2020). Effects of phosalone plant pesticide on sperm parameters and sexual hormone levels in Wistar rats: An experimental study. *International Journal of Reproductive Biomedicine*, 18(9), 785-794. <https://doi.org/10.18502/ijrm.v13i9.7683>