

## Original Research Article

# **A Growing Prosperity of Okra Cultivation (*Abelmoschus esculentus* L.) under Mahaneem (*Ailanthus excelsa*) based Agro-Silviculture System in a Semi-Arid Region of India**

### Abstract

Production of food to feed the rapidly growing population of India is the current challenge. To overcome such global issues of food security and climate change, there is a need to develop a good economically feasible agroforestry system in the tropics. Besides that, Indian farmers also facing the effect of climate change on the monocropping system and lack of optimum land utilization for their economic sustainability. Hence, the economic feasibility of okra cultivation in the agro-silviculture system of mahaneem (*Ailanthus excelsa*) in the arid zone was conducted at the Regional Research Station in Bawal, Haryana, India during two consecutive rainy seasons from July 2020 to October 2020 and July 2021 to October 2021. To create an economically prosperous agro-forestry system, okra varieties Varsha Uphar and Hisar Naveen were planted under various planting densities of Mahaneem in an agri-silvicultural system. Nevertheless, agroforestry is a compelling alternative that encourages social, economic, ecological and agricultural sustainability in response to these challenges. The ratio of net return to cultivation cost was used to compute the benefit-cost ratio for the test of the economic feasibility of the proposed agroforestry system in the semi-arid zone climate of India. The present study revealed that 10 x 10 m planting spacing with the Hisar Naveen okra variety calculated the highest net return (108528 Rs./ha & 123884 Rs./ha in the consecutive years) and benefit-cost ratio (1.90 & 1.86 in the consecutive years) which was highest profitable for farmers to get economic and ecological security. Hence, it is recommended that the Mahaneem-based agro-silviculture system, specifically incorporating the Hisar Naveen variety, proves to be economically more feasible compared to sole cropping which offers practical guidance for farmers and policymakers seeking effective strategies to enhance agricultural productivity amidst evolving environmental conditions.

**Keywords:** Agro-silviculture, Mahaneem, Okra, Agroforestry, Economic feasibility, Arid Zone Agricultural Sustainability.

## **INTRODUCTION**

India is facing the challenge of a population that is rapidly increasing, with a current count of approximately 1.39 billion people and growing at an alarming rate (Conway, 2012). In light

of global issues like food security and climate change, the pursuit of efficient and sustainable farming practices has become crucial. Due to the increased demand for necessities like food, fuel and fodder due to this population expansion, the natural ecological systems of arid and semi-arid regions that include water, land and vegetation are under tremendous strain. Ahmad *et al.* (2019) highlighted that agroforestry is a compelling alternative that encourages social, economic and ecological sustainability in response to these challenges. Growing agricultural crops under tree-based systems has been widely promoted in the tropical regions of the world as a natural resource management strategy that attempts to sustainable goals of agricultural development with the conservation of soils, water, climate and biodiversity (Schroth *et al.* 2004). Nevertheless, agroforestry also provides direct and indirect benefits such as alternative sources of income and employment to the rural poor (Balooni 2003; Puri and Nair 2004; Samra *et al.* 2005; Handa *et al.* 2016; Mehta and Kaushik 2023). Therefore, an exciting opportunity to increase agricultural output is present through the convergence of commercially significant crop cultivation with new agro-silviculture systems in the Arid Zone Region of India. **The climatic conditions in the arid region of India are** critical to growing tree crops and agricultural crops.

Agriculture crop *i.e.* Okra (*Abelmoschus esculentus* L.), cultivation under the shade of Mahaneem (*Ailanthus excelsa* L.) in the semi-arid region of India, is a prime example of one such potential combination in action (Bhuyan *et al.*, 2021). Okra develops in a healthy environment because of the special properties of Mahaneem, which include its ability to replenish the soil and provide a substantial amount of shade. When analysing the complexities of this mutually beneficial relationship, it becomes apparent that the Mahaneem Agro-Silviculture concept improves the resilience of the agro-ecosystem overall in along with making okra production cheaper.

Hence, Mahaneem Agro-Silviculture-based agro-system is important because it can help with the immediate issue of food security. In addition to directly supplying food, Quandt *et al.* (2018) have said that this integrated method shows the ability to improve soil fertility and produce revenue for food purchases. India is confronted with a range of agricultural and environmental challenges, including deforestation, soil erosion, increasing livestock and human population pressure and a declining land-to-man ratio. Fast-growing tree species, especially *Ailanthus excelsa* (Roxb.), popularly called "Ardu" or "Mahaneem," must be incorporated in order to increase the nation's tree cover which is now less than 20% and meet the raw material requirements of wood-based industries (Mehta and Kausik, 2023). Kaushik

*et al.* (2017) highlight *Ailanthus excelsa* as a key element in agroforestry systems and show the sustainability and economic potential of growing trees alongside crops.

The establishment of Mahaneem plants in agroforestry systems that extend across community land, farm boundaries and road avenues is essential to maintaining the stability of ecosystems because it reduces the fluctuations in climate parameters carried on by climate change (Jat et al. 2011). In order to meet the needs of human and livestock populations in a sustainable manner without depleting land or natural resources, *Ailanthus excelsa* integration into farming systems is especially well-suited for the degraded soils of arid and semi-arid agroecosystems (Kaushik *et al.*, 2017).

Nevertheless, *Ailanthus excelsa* L. is acknowledged for its economic significance due to gained popularity in sustainable development, improving the standard of living for resource-poor farmers in semi-arid areas. Mahaneem is a great addition to sustainable practices because of its fast growth, adaptability, and tolerance to diverse pressures. Its wood is used in industries, and its leaves provide valuable feed for tiny ruminants. In order to further support, sustainable agricultural practices in the area, the current study focuses on identifying appropriate combinations of various arable crops under the *Ailanthus excelsa*-based agroforestry system in the semi-arid zones of India. Hence, the present research study provides the economic feasibility of okra under the agro-silviculture system in complex relationships between the semi-arid climate of Haryana to insight into the sustainable practices that propel the increasing wealth of the farmers.

## **MATERIALS AND METHODS**

**Experimental Site:** The present study was conducted at the Regional Research Station of the College of Agriculture, Bawal, Haryana, India which is geographically situated at 28.1°N latitude and 76.5°E longitude. The trial was conducted in two consecutive rainy seasons *i.e.*, from July 2020 to October 2020 and July 2021 to October 2021.

**Experimental Design and Details:** Nine-year-old plantations of *Ailanthus excelsa* served as the tree component of this experiment which planted four distinct spacing configurations (10 × 20 m, 10 × 10 m, 10 × 6.5 m, and 10 × 5 m) with six-month-old *Ailanthus excelsa* seedlings. In order to establish an agri-silvicultural system in agro-forestry, the second component of the system involved intercropping, especially the okra variety *i.e.*, Varsha Uphar and Hisar Naveen, sowed in conjunction with Mahaneem in all the spacings.

**Cost-Benefit Ratio Analysis:** The agroforestry system based on *Ailanthus* underwent a multi-step economic evaluation. The entire cost of cultivation was computed, including labour pay (for clearing land, planting seeds, weeding, watering, harvesting intercrops, and pruning trees), as well as costs for seeds, fertiliser, marketing, and other miscellaneous expenses. The gross return was expressed in rupees per hectare of the intercrop yield at various Mahaneem planting spacings. The cost of cultivation was then reduced from the gross return per hectare to arrive at the net return. The ratio of net return to cultivation cost was then used to compute the benefit-cost ratio.

**Statistical Analysis:** A two-factorial randomised block design with ten treatment combinations was used in the experiment. Standard errors (S.E.m.±) were computed for each situation based on ten treatment combinations that were used. Using the Panse and Sukhatme (1967) method, a crucial difference (CD) at a 5 per cent probability level was calculated to compare treatment means when the treatment effects were considered significant. Data presented in the Bar Diagram in the graphical form were prepared using MS Excel Spreadsheet.

## RESULTS AND DISCUSSION

Evaluating the economic feasibility of cultivation and intercropping is pivotal for the judicious utilization of new agroforestry models. In this study, we conducted a comparative analysis of a Mahaneem (*Ailanthus excelsa* L.) based agroforestry systems with okra intercropping under different planting spacing of mahaneem. The economic information of okra-based agroforestry systems showed in Table 1 that the maximum cost of cultivation & gross return was found in 10 x 10 m planting of mahaneem agro-silviculture system whereas, the least cost of cultivation and gross return was calculated in control (without tree). In the case of higher net return and B: C ratio, 10 x 10 m planting spacing while lease in control (without tree) in the years 2020-21 and 2021-22 (Table 1). When comparing okra varieties, maximum gross return, net return and benefit-cost ratio were observed okra variety Hisar Naveen as compared to the okra variety Varsha Uphar in both years (Table 1).

The highest profitability in terms of gross return (250232 & 294111 Rs./ha) was observed when the Hisar Naveen variety of okra was sown with densely planted Mahaneem (10 × 5 m spacing) which was depicted in Figure 2. The highest net return (108528 & 123884 Rs/ha) was obtained under 10 × 10 m spacing with the variety Hisar Naveen. In contrast, sole planted Varsha Uphar was the least profitable in terms of gross return (124273 & 130588 Rs/ha) and net returns (25207 & 19776 Rs/ha). Despite variations in net and gross returns, the

Benefit-Cost ratio was maximum at  $10 \times 10$  m with Hisar Naveen (1.89 & 1.86), while the lowest ratio was recorded in sole Varsha Uphar cropping (1.25 & 1.18) which was shown in figure 1. Notably, net returns and the Benefit-Cost ratio decreased in the second year of the experiment compared to the first, whereas the cost of cultivation and gross returns increased. When compared to agroforestry systems, sole okra growing showed the lowest cultivation costs (99066 & 110812 Rs/ha) in 2020–21 and 2021–22 (figure 1). Our results show that when compared to solo cropping, the agroforestry system based on *Ailanthus excelsa* produced greater net returns. In various planting spacings, a rise in the number of trees per hectare has resulted in an increase in the cost of cultivation, gross return, and net return, with or without trees. In okra variety Hisar Naveen, the maximum benefit-to-cost ratio was noted with  $10 \times 10$  m plant spacing in all systems. A higher number of trees per hectare, which would result in more wood and fodder production, could be the cause of the improved net returns (123884 Rs) for okra var. Hisar Naveen with  $10 \times 10$  m plant geometry. In agreement with Banerjee *et al.* (2010), pruning and lopping of *A. excelsa* also contributed to greater wood yield. Pratap and Pant (2020) revealed that the net returns (Rs. 177672 ha<sup>-1</sup>) were higher under wider spacing ( $8 \times 5$  m) of *Melia composita*-based agroforestry system when okra intercropped as compared to the sole cropping system.

The increased crop productivity and accompanied rise in market prices are responsible for the improved returns under agroforestry systems. Kaushik *et al.* (2014) observed that horticultural systems with optimum returns are gained for field and vegetable crops. Furthermore, research conducted in North-West India's dry region showed that agri-silvi-horti systems produced higher net returns than solo cropping (Kaushik *et al.*, 2017). These results have been supported by Ahlawat *et al.* (2021), who conclude that under an agroforestry system based on *Ailanthus excelsa*, the highest net returns were achieved in the wheat crop, followed by the onion crop, during the Rabi season of 2014–15.

## CONCLUSION

The findings reveal that Mahaneem planting at a spacing of  $10 \times 10$  m, particularly when coupled with the Hisar Naveen variety of Okra, yields the highest net realization and benefit-cost ratio. This outcome suggests that the Mahaneem-based agro-silviculture system, specifically incorporating the Hisar Naveen variety, proves to be economically more feasible compared to sole cropping. Consequently, this study offers practical guidance for farmers and policymakers seeking effective strategies to enhance agricultural productivity amidst evolving environmental conditions.

## REFERENCES


- Ahlawat, K.S., Johar V., Kumar, V. and Kumar, S. (2021). Performance of agricultural crops under *Ailanthus excelsa* based agroforestry system in semi-arid region of Haryana. *The Pharma Innovation Journal*, **10**(6): 660-662.
- Ahmad, F., Goparaju, L., and Qayum, A. (2019). FAO guidelines and geospatial application for agroforestry suitability mapping: case study of Ranchi, Jharkhand state of India. *Agroforestry Systems*, **93**(2): 531-544.
- Balooni K (2003) Economics of wasteland afforestation in India, a review. *New Forest* 26:101–136.
- Conway, G., 2012. *One billion hungry: can we feed the world?* Cornell University Press.
- Bhuyan Bandan Kumar, C.L. Thakur, Harish Sharma and Dhirender Kumar (2021). Integrated nutrient and performance of okra (*Abelmoschus esculentus* (L.) Moench) under *Morus* based agroforestry system in mid-hills of Himachal Pradesh. *Indian Journal of Agroforestry* 23(2), 25-29 (2021)
- Handa AK, Toky OP, Dhyani SK, Chavan SB (2016) Innovative agroforestry for livelihood security in India. *World Agric* 7:7–16
- Jat, H.S., Singh, R.K. and Mann, J.S. (2011). Ardu (*Ailanthus sp.*) in arid ecosystem: A compatible species for combating with drought and securing livelihood security of resource poor people. *Indian Journal of Traditional Knowledge*, **10**(1): 102-113.
- Kaushik, N., Kumari, S., Singh, S. and Kaushik, J.C. (2014). Productivity and economics of different agri-silvi-horti systems under drip irrigation. *Indian Journal of Agricultural Sciences*, **84**(10): 1166-1171.
- Kaushik, N., Tikoo, A., Yadav, P.K., Deswal, R.P.S. and Singh, S. (2017). Agri–silvi–horti systems for semiarid regions of north-west india. *Agriculture Research*, **6**(2): 150–158.
- Mehta Kajal and N. Kaushik (2023). Impact of planting geometry on *Ailanthus excelsa* L. based silvoarable systems for food and biomass production. *Agroforest Systems*, 97:739–749
- Pratap, U.B. and Pant, K.S. (2020). Performance of okra as influenced by tree spacing and organic manures under *Melia composita* based Agrisilviculture systems. *The Pharma Innovation Journal*, 9(10): 277-281.
- Panse, V. G. and Sukhatme, P.V. (1967). Statistical methods for agricultural workers. *ICAR Publications*, New Delhi, pp. 359.
- Puri S, Nair PKR (2004) Agroforestry research for development in India: 25 years of experiences of a national program. *Agrofor Syst* 61:437–452.
- Quandt, A., Neufeldt, H. and McCabe, J.T. (2018). Building livelihood resilience: what role does agroforestry play. *Climate and Development*, pp: 1-16.
- Samra JS, Kareemulla K, Marwaha PS, Gena HC (2005) Agroforestry and livelihood promotion by cooperatives. National Research Centre for Agroforestry, India, p 104
- Schroth G, Fonseca GAB, Harvey CA, Gascon C, Vasconcelos HL, Izac A-MN (2004) Agroforestry and biodiversity conservation in tropical landscapes. Island Press, Washington, p 523

UNDER PEER REVIEW


**Table 1: Economics of Planting Spacing of Mahaneemunder Agri-Silviculture System (2020-21 & 2021-2022)**

Tree Spacing (m) / Okra varieties	Cost of cultivation (Rs./ha)			Gross return (Rs./ha)			Net return (Rs./ha)			B:C Ratio		
	Varsha Uphar	Hisar Naveen	Mean	Varsha Uphar	Hisar Naveen	Mean	Varsha Uphar	Hisar Naveen	Mean	Varsha Uphar	Hisar Naveen	Mean
<b>2020-21</b>												
<i>10 × 20</i>	113022	113022	113022	161233	167901	164567	48211	54879	51545	1.43	1.49	1.46
<i>10 × 10</i>	121922	121922	121922	224227	230450	227339	102305	108528	105417	1.84	1.90	1.87
<i>10 × 6.5</i>	132574	132574	132574	225931	238224	232078	93357	105650	99504	1.71	1.80	1.75
<i>10 × 5</i>	142917	142917	142917	236084	250232	243158	93167	107315	100241	1.65	1.75	1.70
<i>Control</i>	99066	99066	99066	124273	128382	126328	25207	29316	27262	1.26	1.30	1.28
<i>Mean</i>	121900	121900		194350	203038		72449	81138		1.58	1.65	
<b>C.D. (5%)</b>												
<i>Tree Spacing</i>	2258			4340			5301			0.053		
<i>Varieties</i>	NS			2745			3353			0.033		
<i>Tree Spacing × Varieties</i>	NS			NS			NS			NS		
<b>2021-22</b>												
<i>10 × 20</i>	129518	129518	129518	182000	189044	185522	52482	59526	56004	1.41	1.46	1.43
<i>10 × 10</i>	144081	144081	144081	262013	267965	264989	117932	123884	120908	1.82	1.86	1.84
<i>10 × 6.5</i>	159966	159966	159966	268211	281217	274714	108245	121251	114748	1.68	1.76	1.72
<i>10 × 5</i>	175541	175541	175541	279289	294111	286700	103748	118570	111159	1.59	1.68	1.63
<i>Control</i>	110812	110812	110812	130588	135044	132816	19776	24232	22004	1.18	1.22	1.20
<i>Mean</i>	143984	143984		224420	233476		80437	89493		1.53	1.59	
<b>C.D. (5%)</b>												
<i>Tree Spacing</i>	2354			7252			8049			0.06		
<i>Varieties</i>	NS			4587			5091			0.038		
<i>Tree Spacing × Varieties</i>	NS			NS			NS			NS		

Note: NS= non-significant



**Fig. 1:** Gross Return and Net Return of Interaction of Planting Spacing of Mahaneem and Intercrop Okra Varieties under Agri-Silviculture System and Okra Sole Cropping System (2020-21 & 2021-22)



**Fig. 2:** Cost of Cultivation (Rs/ha) and B:C ratio of Interaction of Planting Spacing of Mahaneem and Intercrop Okra Varieties under Agri-Silviculture System and Okra Sole Cropping System (2020-21 & 2021-2022)