

## **Effect of Bionoculants on Total Chlorophyll Content, Yield of Soybean and Fertility Status of a Vertisols**

### **ABSTRACT**

The field experiment was conducted at the Research Farm, Department of Soil Science & Agricultural Chemistry, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh (INDIA), during *kharif* season 2019-20. The study was aimed to findout the effect of bioinoculants on total chlorophyll content, soybean yield and fertility status of a Vertisols. There were 15 treatments comprised of different beneficial microbial consortia in possible combinations applied as seed treatments with three replications in a randomized block design (RBD). Besides these, two control plots were maintained as fertilized un-inoculated control (FUI) and unfertilized un-inoculated control (UFUI). The recommended dose of fertilizers for soybean is 20 N : 80 P<sub>2</sub>O<sub>5</sub> : 20 K<sub>2</sub>O kg ha<sup>-1</sup>. The results of present study showed that the significant enhancement were noticed by the application of consortia NPK+EM+PGPR in total chlorophyll contents of soybean leaves at 25, 45 and 65 DAS over control. Similarly, significant increases in the soybean yield (seed and stover) were also found by the NPK+EM+PGPR consortia over FUI (control). Further, results revealed that the application of consortia NPK+EM+PGPR were improved the organic carbon and available NPK in a Vertisols. Therefore, it may be concluded that the NPK + EM + PGPR consortia was superior for sustainable soybean productivity and soil fertility.

**Keywords:** Bioinoculants, Chlorophyll content, PGPR, EM, Soybean yield, Vertisols.

## INTRODUCTION

Soybean (*Glycine max* L.) is an important leguminous oil seed crop, which contains 40-45% protein and 18-20% oil. The top two soybean growing states in India are Madhya Pradesh and Maharashtra with 45% and 40% shares, respectively. In Madhya Pradesh the soybean cultivation spreads over 5.2 M ha with total annual production of 6.7 M tones and productivity of 1285 kg ha<sup>-1</sup> [1, 2]. Soybean rhizosphere harbors vast proportions of soil microorganisms, whose activities largely determine the biological condition of the soil and influence the plant growth right from seed germination to maturity [3, 4]. Different microbial consortia i.e. *Pseudomonas* as PGPR is the most efficient and effective strain with significant remarks on isolates of *P. fluorescens* and *P. putida* increasing growth and yield of different crops, especially legumes. *Rhizobium* (diazotroph) is a symbiotic N<sub>2</sub>-fixer with roots of legume crops [5, 6]. It colonizes the roots of specific legumes to form tumor like growths called root nodules, which acts as the factories of ammonia production. *Bacillus sp.* produces soluble exudates organic acids which is recognized as a major mechanism responsible for the release of phosphates from the hydroxyl apatites. *Frateuria aurantia* is a potassium solubilizer which increases the potassium uptake by the plant. Further, the isolates of constitutional microorganisms of EM (Effective Microbial) culture individually have already been evidenced beneficial but their consortium could be more valuable to enhance the supply of nutrients through anti-phytopathogenicity, solubilization, induced phytoresistance and phytostimulator.

The availability of nutrients in the soil for plant utilization is affected not only by the inherent soil characteristics but also by the fertilizer use and practices followed in a cropping system. It has been observed that a major part of the applied nutrient gets fixed and only a small part of it becomes available to the crop plants. However, the different microbial consortia i.e. *Rhizobium*, *Bacillus sp.* etc. offer great promise to the crops enabling the inoculated plants for more uptake of nutrients from the soil [7]. Therefore, the present investigation was carried to find out the effect of various bioinoculants on chlorophyll content, yield of soybean and fertility status of a Vertisols.

## Materials and Methods

The present investigation was conducted during kharif 2019-20 under All India Network Project on Biofertilizers, to find out the effect of bioinoculants on

chlorophyll content in leaves, yield of soybean and fertility status of a Vertisol at the Research Farm, Department of Soil Science, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh (INDIA). It is situated at 23°10'N latitude and 79°57' E longitude at 393 meters above the mean sea level.

The soil of the experimental site is belonging to Kheri series of fine montmorillonitic hyperthermic family of *Typic Haplusterts* (Vertisol) with pH of 7.15, electrical conductivity 0.24 dS m<sup>-1</sup> (1 : 2.5 soil : water ratio) and organic carbon 5.2 g kg<sup>-1</sup>. The soil available N, P and K were 226, 15.8 and 282 kg ha<sup>-1</sup>, respectively. The experiment comprised 15 treatments with three replications having 45 plots laid out under randomized block design (RBD). The treatments of different liquid biofertilizers either solo and/or consortia were applied on soybean (cv. JS 20 69) as seed treatment / basal application as per the appropriate recommendations. The biofertilizers used were diazotroph (*Rhizobium*), PSB - Phosphate Solubilizing Bacteria (*Bacillus* sp.), KSB - Potash Solubilizing Bacteria (*Fraturia aurentia*), PGPR- Plant Growth Promoting Rhizobacteria (*Pseudomonas fluorescens*) and EM- Effective microbial culture/consortium (six bacteria, two fungus and one actinomycetes) culture. The different treatment combinations are presented in Table 1.

**Table 1: Details of different treatment combinations**

Treatment Combinations			
T <sub>1</sub>	<i>Rhizobium</i>	T <sub>9</sub>	NPK + PGPR
T <sub>2</sub>	NPK consortium	T <sub>10</sub>	PK + EM
T <sub>3</sub>	EM culture	T <sub>11</sub>	PK + PGPR
T <sub>4</sub>	PGPR	T <sub>12</sub>	NPK + EM + PGPR
T <sub>5</sub>	PK Consortium	T <sub>13</sub>	PK + EM + PGPR
T <sub>6</sub>	<i>Rhizobium</i> + EM	T <sub>14</sub>	FUI
T <sub>7</sub>	<i>Rhizobium</i> + PGPR	T <sub>15</sub>	UFUI
T <sub>8</sub>	NPK + EM		

The recommended NPK dose for soybean, based on initial soil test, was 20:80:20 (N : P<sub>2</sub>O<sub>5</sub> : K<sub>2</sub>O kg ha<sup>-1</sup>). The sources of N, P and K used were urea, single super phosphate and muriate of potash. Besides these, two types of control plots were maintained as fertilized uninoculated control (FUI) and unfertilized uninoculated control (UFUI) to measure the comparative effects of different microbial inoculants.

#### ***Chlorophyll Content:-***

Leaf chlorophyll (a, b and total) content was estimated by acetone extraction method in fresh plant leaves at 25, 45 and 65 DAS by spectrophotometer at 663 nm and

645 nm wavelength for chlorophyll a and b, respectively. The content of chlorophyll a, b and total were calculated by using following formula:

$$\begin{aligned} \text{Chlorophyll a (mg g}^{-1} \text{ fresh leaves)} &= \frac{12.7 \times (A663) - 2.69 \times (A645) \times V}{a \times 1000 \times W} \\ \text{Chlorophyll b (mg g}^{-1} \text{ fresh leaves)} &= \frac{22.9 \times (A645) - 4.68 \times (A663) \times V}{a \times 1000 \times W} \\ \text{Total Chlorophyll (mg g}^{-1} \text{ fresh leaves)} &= \text{Chlorophyll a} + \text{Chlorophyll b} \end{aligned}$$

Where is, A= Optical density; a = Length of light path in cell (1 cm); W = Weight of sample (1 gm) and V = Volume of solution

#### ***Seed and stover yields:-***

The soybean crop was harvested and bundles were made plot wise, then allowed to dry in the plot for 2-3 days and weighed. After threshing, plot wise yields for straw and seeds were recorded.

#### ***Soil Fertility Status:-***

In the present study, surface 0 -15 cm soil samples were collected from different treatment plots after harvest of soybean crop during 2019-20. Composite soil samples were air-dried at room temperature, pulverized, and sieved through a 2.0 mm sieve. The soil samples were analyzed for different basic soil properties viz. soil pH, electrical conductivity (EC), soil organic carbon and soil available nutrients (NPK) by the standard laboratory procedures.

#### ***Statistical analysis:-***

The data generated on leghemoglobin content in soybean nodules and microbial populations in rhizospheric soil were statistically analyzed to draw suitable inference as per standard method [8].

## **RESULT AND DISCUSSION**

### **Leaf chlorophyll content in leaves of soybean at different stages of growth**

The data related to total chlorophyll content at 25, 45 and 65 DAS of soybean are illustrated in Table 2. It is evident from the results that the maximum total chlorophyll content at 25 DAS was estimated with the microbial consortium of NPK+EM+PGPR for 3.53 mg g<sup>-1</sup> leaf which was 24.9% increment over that of FUI (2.84 mg g<sup>-1</sup> leaf), followed by PK+EM+PGPR, NPK+PGPR and Rhizo+EM for the chlorophyll content of 3.42, 3.38 and 3.31 mg g<sup>-1</sup> leaf, respectively with respective response of 20.6, 19.2 and 16.7 over that of FUI. Similarly, NPK+ EM+PGPR treatment was attributed to

maximum total chlorophyll content in soybean at 45 DAS and 65 DAS for 3.62 and 2.86 mg g<sup>-1</sup> of leaf with increments of 29 and 39%, respectively over that of FUI. The rhizobial inoculation significantly increased the total leaf chlorophyll content by 70%, 62% and 90% measured at 2, 4, and 8 weeks after planting, respectively. An increase in the leaf Fe concentration as a result of the PGPR treatments enhanced the chlorophyll concentration and prevented leaf chlorosis [9, 10, 11].

### **Yields of soybean (seed and stover)**

The data pertaining to the seed yield production of soybean are presented in Table 3 and observed that the consortium of NPK+EM+PGPR achieved significantly maximum seed yield of 2295 kg ha<sup>-1</sup> with relative increment of 39% over FUI (1647 kg ha<sup>-1</sup>), followed by PK+ EM+PGPR, EM culture, NPK+PGPR, NPK consortium, Rhizo+EM, NPK+EM, PK+EM, PK+PGPR, Rhizo+PGPR and *Rhizobium* for grain yield of 2234, 2200, 2157, 2151, 2150, 2217, 2113, 2090, 2033 and 2029 kg ha<sup>-1</sup> representing the response increment of 36, 34, 31, 31, 31, 28, 28, 27, 23 and 23%, respectively. The significant increase in grain yield was recorded due to EM application in farmyard manure as well as in soil amended with the recommended dose of NPK fertilizers in mung bean. It could be due to provides the respective nutrients to the crop along with that the EM culture organisms (acts as phytostimulators) enhanced plant growth and productivity by fixing atmospheric nitrogen. Additionally, the release of trace elements, secreted antioxidant, bioactive compounds (vitamins, hormones and enzymes), in addition to that PGPR acts as plant growth promoters [12, 13 14].

The treatment NPK+EM+PGPR recorded the highest stover yield (Table 3) of soybean (4894 kg ha<sup>-1</sup>) and 62% more response as compared to that of FUI (3026 kg ha<sup>-1</sup>). This was followed by the effects of treatment combinations of PK+EM+PGPR, EM, NPK+PGPR, NPK consortium, PK+EM, NPK+EM, Rhizo+EM, PK+PGPR, Rhizo+PGPR and *Rhizobium* for stover yield of 4654, 4628, 4504, 4450, 4324, 4308, 4291, 4264, 4235 and 4157 kg ha<sup>-1</sup>, respectively corresponding to 54, 53, 49, 47, 43, 42, 42, 41, 40 and 37% increment, respectively over that of FUI. The increase in stover yield might be attributed to the increased availability of N, P and K in soil which resulted in higher growth and development and finally yields [15, 16].

### **Basic soil properties at harvest of soybean**

The data of basic soil properties at harvest of soybean are presented in Table 4. The result revealed that the soil pH ranged between 7.09 -7.15. There were no significant difference found amongst the different treatment. The data also indicated that

the organic carbon content in soil found to increase with different microbial consortia application thereby, lower organic carbon content was found in FUI treatment UFUI treatments [17, 18, 19].

Further, the data on available nutrients content(NPK) of surface soil (0-15 cm depth) showed that the consortium of NPK+EM+PGPR significantly increased the soil available N content ( $288 \text{ kg N ha}^{-1}$ ) with response of 41.6% to that of FUI ( $203 \text{ kg N ha}^{-1}$ ), followed by PK+EM+PGPR, NPK+EM, NPK consortium, NPK+PGPR, Rhizo+EM, EM culture and *Rhizobium* for N content in soil of 287, 274, 271, 266, 260, 259 and  $256 \text{ kg N ha}^{-1}$ , respectively and with 41.4, 35.0, 33.3, 30.7, 27.7, 27.2 and 26.1% response increment, respectively over that of FUI. It has been known to stimulate and enhance plant growth directly as they can improve and mobilize the nutrients such as nitrogen and phosphorous in available forms [20, 21].

Results further revealed that the highest available P content of  $22.4 \text{ kg ha}^{-1}$  was recorded with the application of NPK+EM+PGPR along with 55.30% more response over that of FUI ( $14.4 \text{ kg ha}^{-1}$ ). This was followed by the effects of treatment combination, PK consortium, PK+EM+PGPR, EM culture, NPK+EM, PK+PGPR, PK+EM and NPK+PGPR for P content in soil  $22.4, 22.3, 22.0, 21.5, 21.5, 21.2$  and  $20.7 \text{ kg ha}^{-1}$ , respectively with the respective response of 55.1, 54.2, 52.7, 49.0, 48.9, 46.9 and 43.4% over that of FUI. This might be attributed to many genera of bacteria such as *Bacillus*, *Pseudomonas* and several others have been reported to solubilize varying quantities of phosphorus depending on the efficiency of the strains [22, 23].

It is evident from the results that the maximum available potassium content in soil of soybean at harvest was recorded with NPK+EM+PGPR of  $313 \text{ kg ha}^{-1}$  by 31% increase relative to FUI ( $239 \text{ kg ha}^{-1}$ ), followed by the performance of PK consortium, NPK consortium, EM culture, NPK+EM and PK+EM with values of 301, 295, 294, 293 and  $290 \text{ kg K ha}^{-1}$ , respectively and the respective response of 25.8, 23.4, 22.8, 22.5 and 21.8 % over that of FUI. The findings of Supanjani *et al.* [24] reported that the co-inoculation of KSB and PSB has increased P and K availability. The inoculation of *B. mucilaginosus* (KSB) increased available K over uninoculated control due to mechanism of potassium mobilization by *B. mucilaginosus* [25].

## CONCLUSION

It was concluded that the NPK+EM+PGPR treatment recorded maximum total chlorophyll content in leaves and yield of soybean followed by PK+EM+PGPR

treatments at 25, 45 and 65 DAS. The consortium NPK+EM+PGPR increased the available nutrients (NPK) content 41.6% N, 55.3% P and 31.0% K, respectively overfertilized uninoculated control (FUI). Further, the microbes of EM culture can also enhance crop productivity and sustained soil fertility.

## References

1. SOPA. The soybean Processors Association of India. Email: [xxxx@sopa.org](mailto:xxxx@sopa.org); 2018.
2. Sawarkar S.D., Thakur Risikesh and Khamparia R.S. (2010). Impact of Long Term Continuous Use of Inorganic and Organic Nutrients on Micronutrients Uptake by Soybean in Vertisol. *Journal of Soils and Crops*, **20** (2): 207 – 210.
3. Awasthi R, Tewari R, Nayyar H. Synergy between plants and P solubilizing microbes in soils: effects on growth and physiology of crops. *International Research Journal of Microbiology*. 2011; 2(12): 484-503.
4. Khandagle, A.; Dwivedi, B.S.; Dwivedi, A.K.; Panwar, S.; Thakur, R.K. (2020). Nitrogen fractions under long-term fertilizer and manure applications in soybean – wheat rotation in a Vertisol. *Journal of the Indian Society of Soil Science*, **68**, 186-193.
5. Kushwaha, S.; Sawarkar, S.D.; Thakur, R.; Khamparia, N.K.; Singh, M. (2017). Impact of Long-Term Nutrient Management on Soil N dynamics under Soybean – Wheat Cropping Sequence on a Vertisol. *Journal of the Indian Society of Soil Science*, **65**, 274-282.
6. Patel Gajendra, Dwivedi BS, Dwivedi AK, Thakur Risikesh and Singh Muneshwar (2018). Long-term Effect of Nutrient Management on Soil Biochemical Properties in a Vertisol under Soybean–Wheat Cropping Sequence. *Journal of the Indian Society of Soil Science*, **66** (2): 215-221.
7. Thakur R.K., N.K. Bisen, A.K. Shrivastava, S.K. Rai and S. Sarvade (2023). Impact of Integrated Nutrient Management on Crop Productivity and Soil Fertility under Rice (*Oryza Sativa*) – Chickpea (*Cicer Arietinum*) Cropping System in Chhattisgarh Plain Agro-Climatic Zone. *Indian Journal of Agronomy*, **68**(1): 9-13.
8. Panse, V.G. and S.V. Sukhatme (1970). Statistical methods for Agril. Workers. ICAR Publication.
9. Tairo EV and Ndakidemi PA. 2013. *Bradyrhizobium japonicum* inoculation and phosphorus supplementation on growth and chlorophyll accumulation in soybean (*Glycine max* L.). *American Journal of Plant Sciences*, 2013.

10. Khatik S.K., Thakur Risikesh and Sharma G.D. (2006). Lead: The Heavy Metal in Soil, Water and Plant Environment. *Journal of Industrial Pollution Control*, 22(2): 233–244.
11. Dwivedi, B.S., Rawat, A.K., Dixit, B.K. and Thakur, R.K. (2016) Effect of inputs integration on yield, uptake and economics of kodo millet (*Paspalum scrobiculatum* L). *Economic Affairs*, 61(3): 519-526.
12. Tiwari, R.; Dwivedi, B.S.; Sharma, Y.M.; Thakur, R.; Sharma, A.; Nagwanshi, A. (2023). Soil Properties and Soybean Yield as Influenced by Long Term Fertilizer and Organic Manure Application in a Vertisol under Soybean-Wheat Cropping Sequence. *Legume Research*. DOI: 10.18805/LR-5111.
13. Meshram MK, Dwivedi BS, Naik KR, Thakur RK and Keram KS (2018) Impact of organic and inorganic sources of nutrients on yield, nutrient uptake, soil fertility and economic performance of rice in a Typic Haplustert. *Journal of Soils and Crops*, 28(1) : 31-36.
14. Yaduwanshi B, Sahu RK, Mitra NG, Amule FC and Jakhar S. 2019. Effect of Microbial Consortia on Growth, Nodulation, Yield and Nutrient Uptake of Soybean in Vertisol of Central India. *Int. J. Curr. Microbiol. App. Sci*, 8(9), pp.2649-2659.
15. Keram K.S., Sharma B.L., Sharma G.D. and Thakur R.K. (2014). Impact of zinc application on its translocation into various plant parts of wheat in a Vertisol. *The Bioscan* 9(2): 491-495.
16. Thakur Risikesh, Sharma G.D., Dwivedi B.S. and Khatik S.K. (2007). Chromium: As a Pollutant. *J. of Industrial Pollution Control*, 23(2): 197-203.
17. Thakur Risikesh, Sawarkar S.D., Kauraw D.L. and Singh Muneshwar (2010). Effect of Inorganic and Organic Sources on Nutrients Availability in a Verisol. *Agropedology*, 20 (1): 53-59.
18. Thakur Risikesh and Sawarkar S.D. (2009). Influence of long term continuous application of nutrients and spatial distribution of sulphur on soybean-wheat cropping sequence. *Journal of Soils and Crops* 19, 225 – 228.
19. Sharma Y.M., Jatav R.C., Sharma, G.D. and Thakur Risikesh (2013). Status of Micronutrients in Mixed Red and Black Soils of Rewa District of Madhya Pradesh, India. *Asian Journal of Chemistry*, 25 (6) : 3109-3112.
20. Bairwa Jalendra, B.S. Dwivedi, Anay Rawat, R.K. Thakur and Neeta Mahawar (2021). Long-term effect of nutrient management on soil microbial properties and nitrogen fixation in a Vertisol under soybean–wheat cropping sequence. *Journal of the Indian Society of Soil Science*, 69(2): 171-178.

21. Thakur Risikesh, S. Sarvade and B.S. Dwivedi (2022). Heavy Metals: Soil Contamination and Its Remediation. *AATCC Review*, 10(02): 59-76.
22. Dubey Lokesh, Dwivedi BS, Dwivedi AK and Thakur RK (2016) Effect of long term application of fertilizers and manure on profile distribution of various phosphorus fractions in Vertisol. *Green Farming*, 7(2): 365-370.
23. Pathariya, Priyanka, Dwivedi, B.S, Dwivedi, A.K., Thakur, R.K., Singh Muneshwar and Sarvade, S. (2022) Potassium Balance under Soybean–wheat Cropping System in a 44 Year Old Long Term Fertilizer Experiment on a Vertisol, *Communications in Soil Science and Plant Analysis*, 53(2): 214-226.
24. Supanjani HH, Jung JS and Lee KD. 2006. Rock phosphate-potassium and rock-solubilising bacteria as alternative, sustainable fertilizers. *Agron Sustain Dev*, 26(4), pp.233-240.
25. Sugumaran P and Janarthanam B.2007. Solubilization of potassium containing minerals by bacteria and their effect on plant growth. *World Journal of Agricultural Sciences*, 3(3), pp.350-355.

**Table 2: Effect of bioinoculants on total chlorophyll content in leaves of soybean at different growth stages**

Treatment	Total chlorophyll content (mg g <sup>-1</sup> leaf)		
	25 DAS	45 DAS	65 DAS
<i>Rhizobium</i>	2.77	2.83	2.58
NPK consortium	3.10	3.12	2.87
EM culture	3.11	3.09	2.84
PGPR	2.80	2.79	2.54
PK consortium	2.79	2.90	2.65
<i>Rhizobium</i> + EM	3.31	3.26	3.01
<i>Rhizobium</i> + PGPR	2.97	3.16	2.91
NPK +EM	3.18	3.20	2.95
NPK+PGPR	3.38	3.40	3.15
PK+EM	3.20	3.28	3.03
PK+PGPR	3.23	3.13	2.88
NPK+EM+PGPR	3.53	3.62	3.37
PK+EM+PGPR	3.42	3.42	3.17
FUI	2.84	2.81	2.56
UFUI	2.65	2.67	2.42
<b>CD (<i>p</i>=0.05)</b>	<b>0.25</b>	<b>0.14</b>	<b>0.14</b>

**Table3: Effect of bioinoculants on yields (seed and stover) of soybean at harvest**

Treatment	Yield (kg ha <sup>-1</sup> )	
	Seed	Stover
<i>Rhizobium</i>	2029	4157
NPK consortium	2151	4450
EM culture	2200	4628
PGPR	1905	3436
PK consortium	1872	3258
<i>Rhizobium</i> + EM	2150	4291
<i>Rhizobium</i> + PGPR	2033	4235
NPK +EM	2117	4308
NPK+PGPR	2157	4504
PK+EM	2113	4324
PK+PGPR	2090	4264
NPK+EM+PGPR	2295	4894
PK+EM+PGPR	2234	4654
FUI	1647	3026
UFUI	1127	2787
<b>CD (<i>p</i>=0.05)</b>	<b>346</b>	<b>1050</b>

**Table 4: Effect of bioinoculants on soil pH, organic carbon and available nutrient (N, P and K) contents in soil (0- 15 cm depth) of soybean at harvest**

Treatment	Soil pH	OC (g kg <sup>-1</sup> )	Available nutrients (kg ha <sup>-1</sup> )		
			N	P	K
<i>Rhizobium</i>	7.10	5.4	256	18.3	250
NPK consortium	7.12	5.7	271	17.9	295
EM culture	7.14	5.5	259	22.0	294
PGPR	7.11	5.7	227	15.8	250
PK consortium	7.15	5.5	208	22.4	301
<i>Rhizobium</i> + EM	7.12	5.8	260	20.5	281
<i>Rhizobium</i> + PGPR	7.14	5.9	250	16.1	248
NPK +EM	7.13	5.8	274	21.5	293
NPK+PGPR	7.14	6.1	266	20.7	285
PK+EM	7.14	5.9	247	21.2	290
PK+PGPR	7.10	5.8	247	21.5	265
NPK+EM+PGPR	7.09	6.2	288	23.1	313
PK+EM+PGPR	7.15	6.0	279	22.2	291
FUI	7.09	5.1	203	14.4	239
UFUI	7.10	5.2	187	13.2	231
<b>CD (<i>p</i>=0.05)</b>	<b>NS</b>	<b>0.58</b>	<b>49.1</b>	<b>4.60</b>	<b>48.7</b>