

On Hyperbolic Generalized Woodall Numbers

Abstract. In this study, we introduce the generalized hyperbolic Woodall numbers. As special cases, we study with hyperbolic Woodall, hyperbolic modified Woodall, hyperbolic Cullen numbers and hyperbolic modified Cullen numbers. We present Binet's formulas, generating functions and the summation formulas for these numbers. Besides, we give Catalan's and Cassini's identities and present matrices related to these sequences.

2010 Mathematics Subject Classification. 11B39, 11B83.

Keywords. Woodall numbers, Cullen numbers, hyperbolic numbers, hyperbolic Woodall numbers, hyperbolic Cullen numbers.

1. Introduction

In this section, we give some information which we need about the definition and properties of Woodall numbers.

1.1. Woodall Numbers. The generalized Woodall sequence $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1, W_2, 5, -8, 4)\}_{n \geq 0}$ is defined by the third-order recurrence relations

$$W_n = 5W_{n-1} - 8W_{n-2} + 4W_{n-3} \quad (1.1)$$

with the initial values W_0, W_1, W_2 not all being zero. The sequence $\{W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$W_{-n} = 2W_{-(n-1)} - \frac{5}{4}W_{-(n-2)} + \frac{1}{4}W_{-(n-3)}$$

for $n = 1, 2, 3, \dots$. Therefore, recurrence (1.1) holds for all integer n .

In the following theorem, we give Binet formula of generalized Woodall numbers.

THEOREM 1. [53, Theorem 1.1] Binet formula of generalized Woodall numbers can be given as

$$W_n = (A_1 + A_2 n) \times 2^n + A_3$$

where

$$\begin{aligned} A_1 &= -W_2 + 4W_1 - 3W_0, \\ A_2 &= \frac{W_2 - 3W_1 + 2W_0}{2}, \\ A_3 &= W_2 - 4W_1 + 4W_0, \end{aligned}$$

that is,

$$W_n = ((-W_2 + 4W_1 - 3W_0) + \frac{W_2 - 3W_1 + 2W_0}{2} n) \times 2^n + (W_2 - 4W_1 + 4W_0). \quad (1.2)$$

Here, α, β and γ are the roots of the cubic equation

$$x^3 - 5x^2 + 8x - 4 = (x - 2)^2(x - 1) = 0,$$

where $\alpha = \beta = 2$, $\gamma = 1$.

Now, the first few generalized Woodall numbers with positive subscript and negative subscript are given in the following Table 1.

Table 1. A few generalized Woodall numbers

n	W_n	W_{-n}
0	W_0	W_0
1	W_1	$\frac{1}{4}(8W_0 - 5W_1 + W_2)$
2	W_2	$\frac{1}{4}(11W_0 - 9W_1 + 2W_2)$
3	$4W_0 - 8W_1 + 5W_2$	$\frac{1}{16}(52W_0 - 47W_1 + 11W_2)$
4	$20W_0 - 36W_1 + 17W_2$	$\frac{1}{16}(57W_0 - 54W_1 + 13W_2)$
5	$68W_0 - 116W_1 + 49W_2$	$\frac{1}{64}(240W_0 - 233W_1 + 57W_2)$

Now, we define four specific cases of the sequence $\{W_n\}$.

The Woodall numbers $\{R_n\}$, sometimes called Riesel numbers, and also called Cullen numbers of the second kind, are numbers of the form

$$R_n = n \times 2^n - 1.$$

The first few Woodall numbers are:

$$1, 7, 23, 63, 159, 383, 895, 2047, 4607, 10239, 22527, 49151, 106495, 229375, 491519, 1048575, \dots$$

(sequence A003261 in the OEIS [47]). Woodall numbers were first studied by Allan J. C. Cunningham and H. J. Woodall in [13] in 1917, inspired by James Cullen's earlier study of the similarly-defined Cullen numbers.

The Cullen numbers $\{C_n\}$ are numbers of the form

$$C_n = n \times 2^n + 1.$$

The first few Cullen numbers are:

$$1, 3, 9, 25, 65, 161, 385, 897, 2049, 4609, 10241, 22529, 49153, 106497, 229377, 491521, \dots$$

(sequence A002064 in the OEIS). Woodall and Cullen sequences have been studied by many authors and more detail can be found in the extensive literature dedicated to these sequences, see [5,6,13,24,26,29,32,37,38,39,40].

Note that $\{R_n\}$ and $\{C_n\}$ hold the following relations:

$$R_n = 4R_{n-1} - 4R_{n-2} - 1,$$

$$C_n = 4C_{n-1} - 4C_{n-2} + 1.$$

Note also that the sequences $\{R_n\}$ and $\{C_n\}$ satisfy the following third order linear recurrences:

$$R_n = 5R_{n-1} - 8R_{n-2} + 4R_{n-3}, \quad R_0 = -1, R_1 = 1, R_2 = 7,$$

$$C_n = 5C_{n-1} - 8C_{n-2} + 4C_{n-3}, \quad C_0 = 1, C_1 = 3, C_2 = 9.$$

If we set $G_0 = 0, G_1 = 1, G_2 = 5$ then $\{G_n\}$ is the well-known modified Woodall sequence, if we set $H_0 = 3, H_1 = 5, H_2 = 9$ then $\{H_n\}$ is the well-known modified Cullen sequence. In other words, modified Woodall sequence $\{G_n\}_{n \geq 0}$ and modified Cullen sequence $\{H_n\}_{n \geq 0}$ are defined by the third-order recurrence relations

$$G_n = 5G_{n-1} - 8G_{n-2} + 4G_{n-3}, \quad G_0 = 0, G_1 = 1, G_2 = 5, \quad (1.3)$$

$$H_n = 5H_{n-1} - 8H_{n-2} + 4H_{n-3}, \quad H_0 = 3, H_1 = 5, H_2 = 9, \quad (1.4)$$

The sequences $\{G_n\}_{n \geq 0}, \{H_n\}_{n \geq 0}, \{R_n\}_{n \geq 0}$ and $\{C_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$G_{-n} = 2G_{-(n-1)} - \frac{5}{4}G_{-(n-2)} + \frac{1}{4}G_{-(n-3)},$$

$$H_{-n} = 2H_{-(n-1)} - \frac{5}{4}H_{-(n-2)} + \frac{1}{4}H_{-(n-3)},$$

$$R_{-n} = 2R_{-(n-1)} - \frac{5}{4}R_{-(n-2)} + \frac{1}{4}R_{-(n-3)},$$

$$C_{-n} = 2C_{-(n-1)} - \frac{5}{4}C_{-(n-2)} + \frac{1}{4}C_{-(n-3)},$$

for $n = 1, 2, 3, \dots$ respectively. Therefore, recurrences (1.3) and (1.4) hold for all integer n .

Using the initial conditions in (1.2), Binet's formula of modified Woodall, modified Cullen, Woodall and Cullen sequences are

$$G_n = (n-1)2^n + 1,$$

$$H_n = 2^{n+1} + 1,$$

$$R_n = n \times 2^n - 1,$$

$$C_n = n \times 2^n + 1.$$

Now, we give the generating function and the Cassini identity for generalized Woodall numbers.

The generating function for generalized Woodall numbers is

$$\sum_{n=0}^{\infty} W_n x^n = \frac{W_0 + (W_1 - 5W_0)x + (W_2 - 5W_1 + 8W_0)x^2}{1 - 5x + 8x^2 - 4x^3}. \quad (1.5)$$

The Cassini identity for generalized Woodall numbers is

$$\begin{aligned} W_{n+1}W_{n-1} - W_n^2 &= \frac{1}{4}2^n(A + B2^n + Cn). \\ A &= 4W_1^2 + W_2^2 - 4W_0W_1 + 4W_0W_2 - 5W_1W_2. \\ B &= -4W_0^2 - 9W_1^2 - W_2^2 + 12W_0W_1 - 4W_0W_2 + 6W_1W_2. \\ C &= 8W_0^2 + 12W_1^2 + W_2^2 - 20W_0W_1 + 6W_0W_2 - 7W_1W_2. \end{aligned}$$

For further information about generalized Woodall numbers, see [53].

Next, we give some information about special numbers. In 1989, I. Kantor is worked the hypercomplex numbers systems, [31]. This numbers systems are extensions of real numbers. Some commutative some of hypercomplex number systems are defined below.

Complex numbers are

$$\mathbb{C} = \{z = a + ib : a, b \in \mathbb{R}, i^2 = -1\},$$

hyperbolic (double, split-complex) numbers [48] are

$$\mathbb{H} = \{h = a + jb : a, b \in \mathbb{R}, j^2 = 1, j \neq \pm 1\}$$

and dual numbers [20] are

$$\mathbb{D} = \{d = a + \varepsilon b : a, b \in \mathbb{R}, \varepsilon^2 = 0, \varepsilon \neq 0\}.$$

One of the non-commutative examples of hypercomplex number systems are quaternions, [28],

$$\mathbb{H}_{\mathbb{Q}} = \{q = a_0 + ia_1 + ja_2 + ka_3 : a_0, a_1, a_2, a_3 \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\},$$

octonions [3] and sedenions [51]. The algebras \mathbb{C} (complex numbers), $\mathbb{H}_{\mathbb{Q}}$ (quaternions), \mathbb{O} (octonions) and \mathbb{S} (sedenions) are real algebras obtained from the real numbers \mathbb{R} by a doubling procedure called the Cayley-Dickson Process. This doubling process can be extended beyond the sedenions to form what are known as the 2^n -ions (see for example [7], [30], [41]).

Quaternions were invented by Irish mathematician W. R. Hamilton (1805-1865) [28] as an extension to the complex numbers. Hyperbolic numbers with complex coefficients are introduced by J. Cockle in 1848, [12].

Now, we will give some information related to hyperbolic numbers. **We present hyperbolic numbers as follows:**

$$\mathbb{H} = \{h = a + jb : a, b \in \mathbb{R}, j^2 = 1, j \neq \pm 1\}.$$

The base elements $\{1, j\}$ of hyperbolic numbers satisfy the following properties (commutative multiplications):

$$1.j = j, \quad j^2 = j.j = 1$$

where j symbolizes the hyperbolic unit ($j^2 = 1$).

The multiplication of two hyperbolic numbers $m = a_0 + ja_1$ and $n = b_0 + jb_1$ is

$$mn = a_0b_0 + a_1b_1 + j(a_0b_1 + a_1b_0).$$

Sum of two hyperbolic numbers $m = a_0 + ja_1$ and $n = b_0 + jb_1$ is

$$m + n = a_0 + b_0 + j(a_1 + b_1).$$

Now, we give details about hyperbolic and some information related to hyperbolic sequences from the literature.

- Richter, [44] worked On Hyperbolic Complex Numbers.
- Gürses, Şentürk and Yüce, [25] studied A Study on Dual-Generalized Complex and Hyperbolic-Generalized Complex numbers.
- Cockle [12] worked the Hyperbolic numbers with complex coefficients.
- Aydın, [1] worked hyperbolic Fibonacci numbers given by

$$\tilde{F}_n = F_n + hF_{n+1}, \quad (h^2 = 1)$$

where Fibonacci numbers, respectively, given by $F_n = F_{n-1} + F_{n-2}$ with the initial condition $F_1 = F_2 = 1$, ($n \geq 3$).

- Dikmen, [17] worked hyperbolic Jacobsthal numbers given by

$$\hat{J}_n = J_n + hJ_{n+1}, \quad (h^2 = 1)$$

where Jacobsthal numbers, respectively, given by $J_n = J_{n-1} + 2J_{n-2}$, $J_0 = 0$, $J_1 = 1$.

- Taş, [74] worked on hyperbolic Jacobsthal-Lucas sequence given by

$$HJ_n = J_n + hJ_{n+1}, \quad (h^2 = 1)$$

where Jacobsthal-Lucas numbers, respectively, given by $J_{n+2} = J_{n+1} + 2J_n$, with the initial condition $J_0 = 2$, $J_1 = 1$.

- Soykan and Taşdemir, [57] worked on hyperbolic generalized Jacobsthal numbers given by

$$\tilde{V}_n = V_n + hV_{n+1}, \quad (h^2 = 1)$$

where generalized Jacobsthal numbers are given by $V_n = V_{n-1} + 2V_{n-2}$, $V_0 = a$, $V_1 = b$ ($n \geq 2$) with the initial values V_0, V_1 not all being zero.

- Dişkaya, Menken, Catarino, [19] worked on hyperbolic Leonardo and hyperbolic Francois quaternions given by

$$\begin{aligned} H\mathcal{L}_n &= \mathcal{L}_n e_0 + \mathcal{L}_{n+1} e_1 + \mathcal{L}_{n+2} e_2 + \mathcal{L}_{n+3} e_3, \\ H\mathcal{F}_n &= \mathcal{F}_n e_0 + \mathcal{F}_{n+1} e_1 + \mathcal{F}_{n+2} e_2 + \mathcal{F}_{n+3} e_3 \end{aligned}$$

where Francois and Leonardo numbers, respectively, given by $\mathcal{F}_n = \mathcal{F}_{n-1} + \mathcal{F}_{n-2} + 1$, with the initial condition $\mathcal{F}_0 = 2$, $\mathcal{F}_1 = 1$ and $\mathcal{L}_{n+2} = \mathcal{L}_{n+1} + \mathcal{L}_n$, with the initial condition $\mathcal{L}_0 = 1$, $\mathcal{L}_1 = 1$.

- Dikmen and Altınoy, [18] worked on third order hyperbolic Jacobsthal numbers given by

$$\begin{aligned} \widehat{J}_n^{(3)} &= J_n^{(3)} + hJ_{n+1}^{(3)}, \\ \widehat{j}_n^{(3)} &= j_n^{(3)} + hj_{n+1}^{(3)} \end{aligned}$$

where Jacobsthal numbers, respectively, given by $J_n^{(3)} = J_{n-1}^{(3)} + J_{n-2}^{(3)} + 2J_{n-3}^{(3)}$, $J_0^{(3)} = 0$, $J_1^{(3)} = 1$, $J_2^{(3)} = 1$, $j_n^{(3)} = j_{n-1}^{(3)} + j_{n-2}^{(3)} + 2j_{n-3}^{(3)}$, $j_0^{(3)} = 2$, $j_1^{(3)} = 1$, $j_2^{(3)} = 5$.

Next section, we present the hyperbolic generalized Woodall numbers and their generating functions and Binet's formulas.

2. Hyperbolic Generalized Woodall Numbers

In this section, we define hyperbolic generalized Woodall numbers and present generating functions and Binet's formulas for **these numbers**.

We now define hyperbolic generalized Woodall numbers over \mathbb{H} . The n th hyperbolic generalized Woodall number is

$$\mathcal{H}W_n = W_n + jW_{n+1}. \quad (2.1)$$

with the initial values $\mathcal{H}W_0$, $\mathcal{H}W_1$, $\mathcal{H}W_2$. (2.1) can be written to negative subscripts by defining,

$$\mathcal{H}W_{-n} = W_{-n} + jW_{-n+1}.$$

so identity (2.1) holds for all integers n .

The special cases of the n th dual hyperbolic generalized Woodall numbers are given as

$$\begin{aligned} \mathcal{H}G_n &= G_n + jG_{n+1}, \\ \mathcal{H}H_n &= H_n + jH_{n+1}, \\ \mathcal{H}R_n &= R_n + jR_{n+1}, \\ \mathcal{H}C_n &= C_n + jC_{n+1}. \end{aligned}$$

Hence, for $n \geq 0$, the following identity is true.

$$\mathcal{H}W_n = 5\mathcal{H}W_{n-1} - 8\mathcal{H}W_{n-2} + 4\mathcal{H}W_{n-3}. \quad (2.2)$$

The sequence $\{\mathcal{H}W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$\mathcal{H}W_{-n} = -2\mathcal{H}W_{-(n-1)} - \frac{5}{4}\mathcal{H}W_{-(n-2)} + \frac{1}{4}\mathcal{H}W_{-(n-3)}.$$

for $n = 1, 2, 3, \dots$ respectively. Therefore, recurrence (2.2) holds for all integer n .

The few hyperbolic generalized Woodall numbers with positive subscript and negative subscript are given in the following Table 2.

Table 2. A few hyperbolic generalized Woodall numbers.

n	$\mathcal{H}W_n$	$\mathcal{H}W_{-n}$
0	$\mathcal{H}W_0$	$\mathcal{H}W_0$
1	$\mathcal{H}W_1$	$\frac{1}{4}(8\mathcal{H}W_0 - 5\mathcal{H}W_1 + \mathcal{H}W_2)$
2	$\mathcal{H}W_2$	$\frac{1}{4}(11\mathcal{H}W_0 - 9\mathcal{H}W_1 + 2\mathcal{H}W_2)$
3	$4\mathcal{H}W_0 - 8\mathcal{H}W_1 + 5\mathcal{H}W_2$	$\frac{1}{16}(52\mathcal{H}W_0 - 47\mathcal{H}W_1 + 11\mathcal{H}W_2)$
4	$20\mathcal{H}W_0 - 36\mathcal{H}W_1 + 17\mathcal{H}W_2$	$\frac{1}{16}(57\mathcal{H}W_0 - 54\mathcal{H}W_1 + 13\mathcal{H}W_2)$
5	$68\mathcal{H}W_0 - 116\mathcal{H}W_1 + 49\mathcal{H}W_2$	$\frac{1}{64}(240\mathcal{H}W_0 - 233\mathcal{H}W_1 + 57\mathcal{H}W_2)$

Note that

$$\mathcal{H}W_0 = W_0 + jW_1,$$

$$\mathcal{H}W_1 = W_1 + jW_2,$$

$$\mathcal{H}W_2 = W_2 + jW_3 = W_2 + j(4W_0 - 8W_1 + 5W_2).$$

For hyperbolic modified Woodall numbers (taking $W_n = G_n$, $G_0 = 0$, $G_1 = 1$, $G_2 = 5$), we get

$$\mathcal{H}G_0 = G_0 + jG_1 = j,$$

$$\mathcal{H}G_1 = G_1 + jG_2 = 1 + 5j,$$

$$\mathcal{H}G_2 = G_2 + jG_3 = 5 + 17j$$

and for hyperbolic modified Cullen numbers (taking $W_n = H_n$, $H_0 = 3$, $H_1 = 5$, $H_2 = 9$), we get

$$\mathcal{H}H_0 = H_0 + jH_1 = 3 + 5j,$$

$$\mathcal{H}H_1 = H_1 + jH_2 = 5 + 9j,$$

$$\mathcal{H}H_2 = H_2 + jH_3 = 9 + 17j$$

and for hyperbolic Woodall numbers (taking $W_n = R_n$, $R_0 = -1$, $R_1 = 1$, $R_2 = 7$), we get

$$\mathcal{H}R_0 = R_0 + jR_1 = -1 + j,$$

$$\mathcal{H}R_1 = R_1 + jR_2 = 1 + 7j,$$

$$\mathcal{H}R_2 = R_2 + jR_3 = 7 + 23j$$

and for hyperbolic Cullen numbers (taking $W_n = C_n$, $C_0 = 1$, $C_1 = 3$, $C_2 = 9$), we get

$$\mathcal{H}C_0 = C_0 + jC_1 = 1 + 3j,$$

$$\mathcal{H}C_1 = C_1 + jC_2 = 3 + 9j,$$

$$\mathcal{H}C_2 = C_2 + jC_3 = 9 + 25j.$$

A few hyperbolic modified Woodall numbers, hyperbolic modified Cullen numbers, hyperbolic Woodall numbers and hyperbolic Cullen numbers with positive subscript and negative subscript are given in Table 3, Table 4, Table 5 and Table 6.

Table 3. Hyperbolic modified Woodall numbers

n	$\mathcal{H}G_n$	$\mathcal{H}G_{-n}$
0	j	j
1	$1 + 5j$	0
2	$5 + 17j$	$\frac{1}{4}$
3	$17 + 49j$	$\frac{1}{2} + \frac{1}{4}j$
4	$49 + 129j$	$\frac{11}{16} + \frac{1}{2}j$
5	$129 + 321j$	$\frac{13}{16} + \frac{11}{16}j$

Table 5. Hyperbolic Woodall numbers

n	$\mathcal{H}R_n$	$\mathcal{H}R_{-n}$
0	$-1 + j$	$-1 + j$
1	$1 + 7j$	$-\frac{3}{2} - j$
2	$7 + 23j$	$-\frac{3}{2} - \frac{3}{2}j$
3	$23 + 63j$	$-\frac{11}{8} - \frac{3}{2}j$
4	$63 + 159j$	$-\frac{5}{4} - \frac{11}{8}j$
5	$159 + 383j$	$-\frac{37}{32} - \frac{5}{4}j$

Table 4. Hyperbolic modified Cullen numbers

n	$\mathcal{H}H_n$	$\mathcal{H}H_{-n}$
0	$3 + 5j$	$3 + 5j$
1	$5 + 9j$	$2 + 3j$
2	$9 + 17j$	$\frac{3}{2} + 3j$
3	$17 + 33j$	$\frac{5}{4} + \frac{3}{2}j$
4	$33 + 65j$	$\frac{9}{8} + \frac{3}{2}j$
5	$65 + 129j$	$\frac{17}{16} + \frac{9}{8}j$

Table 6. Hyperbolic Cullen numbers

n	$\mathcal{H}C_n$	$\mathcal{H}C_{-n}$
0	$1 + 3j$	$1 + 3j$
1	$3 + 9j$	$\frac{1}{2} + j$
2	$9 + 25j$	$\frac{1}{2} + j$
3	$25 + 65j$	$\frac{5}{8} + \frac{1}{2}j$
4	$65 + 161j$	$\frac{3}{4} + \frac{1}{2}j$
5	$161 + 385j$	$\frac{27}{32} + \frac{3}{4}j$

Now, we will state Binet's formula for the hyperbolic generalized Woodall numbers and in the rest of the paper, we fix the following notations:

$$\widehat{\alpha} = 1 + 2j,$$

$$\widehat{\beta} = 2j,$$

$$\widehat{\gamma} = 1 + j.$$

Note that we have the following identities:

$$\begin{aligned}
\widehat{\alpha}^2 &= 5 + 4j, \\
\widehat{\beta}^2 &= 4, \\
\widehat{\gamma}^2 &= 2 + 2j, \\
\widehat{\alpha}\widehat{\beta} &= 4 + 2j, \\
\widehat{\alpha}\widehat{\gamma} &= 3 + 3j, \\
\widehat{\beta}\widehat{\gamma} &= 2 + 2j, \\
\widehat{\alpha}\widehat{\beta}\widehat{\gamma} &= 6 + 6j.
\end{aligned}$$

2.1. Binet's Formula. Now, we present Binet's formula in the following theorem.

THEOREM 2. (*Binet's Formula*) For any integer n , the n th hyperbolic generalized Woodall number is

$$\mathcal{H}W_n = (A_1\widehat{\alpha} + A_2\widehat{\beta} + A_2n\widehat{\alpha})2^n + A_3\widehat{\gamma}. \quad (2.3)$$

Proof. Using Binet's formula given below

$$W_n = (A_1 + A_2n)2^n + A_3,$$

we obtain

$$\begin{aligned}
\mathcal{H}W_n &= W_n + jW_{n+1} \\
&= (A_1 + A_2n)2^n + A_3 + j((A_1 + A_2(n+1))2^{n+1} + A_3) \\
&= A_12^n + A_2n2^n + A_3 \\
&\quad + jA_12^{n+1} + jA_2n2^{n+1} + jA_22^{n+1} + jA_3 \\
&= A_12^n(1 + 2j) + A_2n2^n(1 + 2j) + A_22^n(2j) + A_3(1 + j) \\
&= A_12^n\widehat{\alpha} + A_2n2^n\widehat{\alpha} + A_22^n\widehat{\beta} + A_3\widehat{\gamma} \\
&= (A_1\widehat{\alpha} + A_2\widehat{\beta} + A_2n\widehat{\alpha})2^n + A_3\widehat{\gamma}.
\end{aligned}$$

This proves (2.3). \square

As special cases, for any integer n , the Binet's Formula of n th hyperbolic modified Woodall number, hyperbolic modified Cullen number, hyperbolic Woodall number and hyperbolic Cullen number are

- $\mathcal{H}G_n = (-\widehat{\alpha} + \widehat{\beta} + n\widehat{\alpha})2^n + \widehat{\gamma}$,
 $\mathcal{H}G_n = 1 + (n-1)2^n + j(1 + n2^{n+1})$.
- $\mathcal{H}H_n = (2\widehat{\alpha})2^n + \widehat{\gamma}$,
 $\mathcal{H}H_n = 1 + 2^{n+1} + j(1 + 2^{n+2})$.

- $\mathcal{H}R_n = (\widehat{\beta} + n\widehat{\alpha})2^n - \widehat{\gamma}$,
 $\mathcal{H}R_n = -1 + n2^n + j(-1 + 2^{n+1} + n2^{n+1})$.
- $\mathcal{H}C_n = (\widehat{\beta} + n\widehat{\alpha})2^n + \widehat{\gamma}$,
 $\mathcal{H}C_n = 1 + n2^n + j(1 + 2^{n+1} + n2^{n+1})$.

Next, we present generating function **of the hyperbolic generalized Woodall numbers**.

2.2. Generating Function.

THEOREM 3. *The generating function for the hyperbolic generalized Woodall numbers is*

$$\sum_{n=0}^{\infty} \mathcal{H}W_n x^n = \frac{\mathcal{H}W_0 + (\mathcal{H}W_1 - 5\mathcal{H}W_0)x + (\mathcal{H}W_2 - 5\mathcal{H}W_1 + 8\mathcal{H}W_0)x^2}{1 - 5x + 8x^2 - 4x^3}. \quad (2.4)$$

Proof. Let

$$g(x) = \sum_{n=0}^{\infty} \mathcal{H}W_n x^n$$

be generating function of the hyperbolic generalized Woodall numbers. Then, using the definition of the hyperbolic generalized Woodall numbers, and subtracting $xg(x)$, $x^2g(x)$ and $x^3g(x)$ from $g(x)$, we obtain (note the shift in the index n in the third line)

$$\begin{aligned} (1 - 5x + 8x^2 - 4x^3)g(x) &= \sum_{n=0}^{\infty} \mathcal{H}W_n x^n - 5x \sum_{n=0}^{\infty} \mathcal{H}W_n x^n + 8x^2 \sum_{n=0}^{\infty} \mathcal{H}W_n x^n - 4x^3 \sum_{n=0}^{\infty} \mathcal{H}W_n x^n \\ &= \sum_{n=0}^{\infty} \mathcal{H}W_n x^n - 5 \sum_{n=0}^{\infty} \mathcal{H}W_n x^{n+1} + 8 \sum_{n=0}^{\infty} \mathcal{H}W_n x^{n+2} - 4 \sum_{n=0}^{\infty} \mathcal{H}W_n x^{n+3} \\ &= \sum_{n=0}^{\infty} \mathcal{H}W_n x^n - 5 \sum_{n=1}^{\infty} \mathcal{H}W_{n-1} x^n + 8 \sum_{n=2}^{\infty} \mathcal{H}W_{n-2} x^n - 4 \sum_{n=3}^{\infty} \mathcal{H}W_{n-3} x^n \\ &= (\mathcal{H}W_0 + \mathcal{H}W_1 x + \mathcal{H}W_2 x^2) - 5(\mathcal{H}W_0 x + \mathcal{H}W_1 x^2) + 8\mathcal{H}W_0 x^2 \\ &\quad + \sum_{n=3}^{\infty} (\mathcal{H}W_n - 5\mathcal{H}W_{n-1} + 8\mathcal{H}W_{n-2} - 4\mathcal{H}W_{n-3}) x^n \\ &= \mathcal{H}W_0 + (\mathcal{H}W_1 - 5\mathcal{H}W_0)x + (\mathcal{H}W_2 - 5\mathcal{H}W_1 + 8\mathcal{H}W_0)x^2. \end{aligned}$$

Note that, we **use** the recurrence relation $\mathcal{H}W_n = 5\mathcal{H}W_{n-1} - 8\mathcal{H}W_{n-2} + 4\mathcal{H}W_{n-3}$. Rearranging above equation, we get

$$g(x) = \frac{\mathcal{H}W_0 + (\mathcal{H}W_1 - 5\mathcal{H}W_0)x + (\mathcal{H}W_2 - 5\mathcal{H}W_1 + 8\mathcal{H}W_0)x^2}{1 - 5x + 8x^2 - 4x^3}.$$

The proof is finished. \square

As special cases, the generating functions for the hyperbolic modified Woodall, hyperbolic modified Cullen, hyperbolic Woodall and hyperbolic Cullen numbers are

$$\begin{aligned}\sum_{n=0}^{\infty} \mathcal{H}G_n x^n &= \frac{j+x}{1-5x+8x^2-4x^3}, \\ \sum_{n=0}^{\infty} \mathcal{H}H_n x^n &= \frac{5j+3+(-16j-10)x+(12j+8)x^2}{1-5x+8x^2-4x^3}, \\ \sum_{n=0}^{\infty} \mathcal{H}R_n x^n &= \frac{-1+j+(2j+6)x+(-4j-6)x^2}{1-5x+8x^2-4x^3}\end{aligned}$$

and

$$\sum_{n=0}^{\infty} \mathcal{H}C_n x^n = \frac{3j+1+(-6j-2)x+(4j+2)x^2}{1-5x+8x^2-4x^3}$$

respectively.

Now, we obtained the Binet formula using the generating function.

2.3. Obtaining Binet's Formula From Generating Function. We obtain the Binet's formula of hyperbolic generalized Woodall number $\{\mathcal{H}W_n\}$ by the use of generating function for $\mathcal{H}W_n$.

THEOREM 4. *(The Binet's formula of hyperbolic generalized Woodall numbers)*

$$\mathcal{H}W_n = (A_1 \hat{\alpha} + A_2 \hat{\beta} + A_2 n \hat{\alpha}) 2^n + A_3 \hat{\gamma}. \quad (2.5)$$

Proof. Let

$$\sum_{n=0}^{\infty} \mathcal{H}W_n x^n = \frac{\mathcal{H}W_0 + (\mathcal{H}W_1 - 5\mathcal{H}W_0)x + (\mathcal{H}W_2 - 5\mathcal{H}W_1 + 8\mathcal{H}W_0)x^2}{1-5x+8x^2-4x^3}.$$

Then, we write

$$\frac{\mathcal{H}W_0 + (\mathcal{H}W_1 - 5\mathcal{H}W_0)x + (\mathcal{H}W_2 - 5\mathcal{H}W_1 + 8\mathcal{H}W_0)x^2}{(1-x)(1-2x)^2} = \frac{d_1}{(1-x)} + \frac{d_2}{(1-2x)} + \frac{d_3}{(1-2x)^2}. \quad (2.6)$$

So

$$\begin{aligned}\mathcal{H}W_0 + (\mathcal{H}W_1 - 5\mathcal{H}W_0)x + (\mathcal{H}W_2 - 5\mathcal{H}W_1 + 8\mathcal{H}W_0)x^2 &= (d_1 + d_2 + d_3) + (-4d_1 - 3d_2 - d_3)x \\ &\quad + (4d_1 + 2d_2)x^2.\end{aligned}$$

We get

$$\begin{aligned}\mathcal{H}W_0 &= d_1 + d_2 + d_3, \\ \mathcal{H}W_1 - 5\mathcal{H}W_0 &= -4d_1 - 3d_2 - d_3, \\ \mathcal{H}W_2 - 5\mathcal{H}W_1 + 8\mathcal{H}W_0 &= 4d_1 + 2d_2.\end{aligned}$$

If we solve these simultaneous equation,

$$\begin{aligned} d_1 &= 4\mathcal{H}W_0 - 4\mathcal{H}W_1 + \mathcal{H}W_2, \\ d_2 &= -4\mathcal{H}W_0 + \frac{11}{2}\mathcal{H}W_1 - \frac{3}{2}\mathcal{H}W_2, \\ d_3 &= \mathcal{H}W_0 - \frac{3}{2}\mathcal{H}W_1 + \frac{1}{2}\mathcal{H}W_2. \end{aligned}$$

Thus (2.6) can be written as

$$\begin{aligned} \sum_{n=0}^{\infty} \mathcal{H}W_n x^n &= d_1 \frac{1}{(1-x)} + d_2 \frac{1}{(1-2x)} + d_3 \frac{1}{(2x-1)^2}, \\ &= d_1 \sum_{n=0}^{\infty} x^n + d_2 \sum_{n=0}^{\infty} 2^n x^n + d_3 \sum_{n=0}^{\infty} 2^n (n+1) x^n, \\ &= \sum_{n=0}^{\infty} (d_1 + d_2 2^n + d_3 2^n (n+1)) x^n, \\ &= \sum_{n=0}^{\infty} (4\mathcal{H}W_0 - 4\mathcal{H}W_1 + \mathcal{H}W_2 + (-4\mathcal{H}W_0 + \frac{11}{2}\mathcal{H}W_1 - \frac{3}{2}\mathcal{H}W_2) 2^n \\ &\quad + (\mathcal{H}W_0 - \frac{3}{2}\mathcal{H}W_1 + \frac{1}{2}\mathcal{H}W_2) 2^n (n+1)) x^n, \\ &= \sum_{n=0}^{\infty} (4\mathcal{H}W_0 - 4\mathcal{H}W_1 + \mathcal{H}W_2 + (-4\mathcal{H}W_0 + \frac{11}{2}\mathcal{H}W_1 - \frac{3}{2}\mathcal{H}W_2) 2^n \\ &\quad + (\mathcal{H}W_0 - \frac{3}{2}\mathcal{H}W_1 + \frac{1}{2}\mathcal{H}W_2) 2^n + (\mathcal{H}W_0 - \frac{3}{2}\mathcal{H}W_1 + \frac{1}{2}\mathcal{H}W_2) 2^n n) x^n, \\ &= \sum_{n=0}^{\infty} (4\mathcal{H}W_0 - 4\mathcal{H}W_1 + \mathcal{H}W_2 + (\mathcal{H}W_0 - \frac{3}{2}\mathcal{H}W_1 + \frac{1}{2}\mathcal{H}W_2) n 2^n \\ &\quad + (-3\mathcal{H}W_0 + 4\mathcal{H}W_1 - \mathcal{H}W_2) 2^n) x^n, \\ &= \sum_{n=0}^{\infty} ((-3\mathcal{H}W_0 + 4\mathcal{H}W_1 - \mathcal{H}W_2) + (\mathcal{H}W_0 - \frac{3}{2}\mathcal{H}W_1 + \frac{1}{2}\mathcal{H}W_2) n) 2^n \\ &\quad + 4\mathcal{H}W_0 - 4\mathcal{H}W_1 + \mathcal{H}W_2) x^n. \end{aligned}$$

This gives

$$\mathcal{H}W_n = (\mathcal{H}A_1 + \mathcal{H}A_2 n) 2^n + \mathcal{H}A_3$$

where

$$\begin{aligned} \mathcal{H}A_1 &= -3\mathcal{H}W_0 + 4\mathcal{H}W_1 - \mathcal{H}W_2, \\ \mathcal{H}A_2 &= \mathcal{H}W_0 - \frac{3}{2}\mathcal{H}W_1 + \frac{1}{2}\mathcal{H}W_2, \\ \mathcal{H}A_3 &= 4\mathcal{H}W_0 - 4\mathcal{H}W_1 + \mathcal{H}W_2. \end{aligned}$$

Note that the following equalities are true:

$$\begin{aligned} A_1 \hat{\alpha} + A_2 \hat{\beta} &= (-W_2 + 4W_1 - 3W_0)(1 + 2j) + \left(\frac{W_2 - 3W_1 + 2W_0}{2}\right)(2j) \\ &= -3W_0 + 4W_1 - W_2 + j(-4W_0 + 5W_1 - W_2). \end{aligned}$$

$$\begin{aligned}
A_2\widehat{\alpha} &= \frac{W_2 - 3W_1 + 2W_0}{2}(1 + 2j) \\
&= W_0 - \frac{3}{2}W_1 + \frac{1}{2}W_2 + j(2W_0 - 3W_1 + W_2).
\end{aligned}$$

$$A_3\widehat{\gamma} = W_2 - 4W_1 + 4W_0 + j(W_2 - 4W_1 + 4W_0).$$

Therefore, we can write the following equalition:

$$\mathcal{H}W_n = (A_1\widehat{\alpha} + A_2\widehat{\beta} + A_2n\widehat{\alpha})2^n + A_3\widehat{\gamma}.$$

The proof is finished. \square

Next, using Theorem 4, we present the Binet's formulas of hyperbolic modified Woodall, hyperbolic modified Cullen, hyperbolic Woodall and hyperbolic Cullen numbers.

3. Some Identities For Hyperbolic Generalized Woodall Numbers

We now present a few special identities for the hyperbolic generalized Woodall sequence $\{\mathcal{H}W_n\}$. The following theorem presents the Simpson's identity for the hyperbolic generalized Woodall numbers.

THEOREM 5. (*Simpson's formula for hyperbolic generalized Woodall sequence*) For all integers n we have

$$\begin{vmatrix} \mathcal{H}W_{n+2} & \mathcal{H}W_{n+1} & \mathcal{H}W_n \\ \mathcal{H}W_{n+1} & \mathcal{H}W_n & \mathcal{H}W_{n-1} \\ \mathcal{H}W_n & \mathcal{H}W_{n-1} & \mathcal{H}W_{n-2} \end{vmatrix} = 4^n \begin{vmatrix} \mathcal{H}W_2 & \mathcal{H}W_1 & \mathcal{H}W_0 \\ \mathcal{H}W_1 & \mathcal{H}W_0 & \mathcal{H}W_{-1} \\ \mathcal{H}W_0 & \mathcal{H}W_{-1} & \mathcal{H}W_{-2} \end{vmatrix}.$$

Proof. For the proof we use mathematical induction. For $n = 0$ identity is true. **First, we proof the identity for $n \geq 0$.** Now, we obtain **the identity** is true for $n = k$. Hence, we write the following identity

$$\begin{vmatrix} \mathcal{H}W_{k+2} & \mathcal{H}W_{k+1} & \mathcal{H}W_k \\ \mathcal{H}W_{k+1} & \mathcal{H}W_k & \mathcal{H}W_{k-1} \\ \mathcal{H}W_k & \mathcal{H}W_{k-1} & \mathcal{H}W_{k-2} \end{vmatrix} = 4^k \begin{vmatrix} \mathcal{H}W_2 & \mathcal{H}W_1 & \mathcal{H}W_0 \\ \mathcal{H}W_1 & \mathcal{H}W_0 & \mathcal{H}W_{-1} \\ \mathcal{H}W_0 & \mathcal{H}W_{-1} & \mathcal{H}W_{-2} \end{vmatrix}.$$

For $n = k + 1$, we get

$$\begin{aligned}
\begin{vmatrix} \mathcal{H}W_{k+3} & \mathcal{H}W_{k+2} & \mathcal{H}W_{k+1} \\ \mathcal{H}W_{k+2} & \mathcal{H}W_{k+1} & \mathcal{H}W_k \\ \mathcal{H}W_{k+1} & \mathcal{H}W_k & \mathcal{H}W_{k-1} \end{vmatrix} &= \begin{vmatrix} 5\mathcal{H}W_{k+2} - 8\mathcal{H}W_{k+1} + 4\mathcal{H}W_k & \mathcal{H}W_{k+2} & \mathcal{H}W_{k+1} \\ 5\mathcal{H}W_{k+1} - 8\mathcal{H}W_k + 4\mathcal{H}W_{k-1} & \mathcal{H}W_{k+1} & \mathcal{H}W_k \\ 5\mathcal{H}W_k - 8\mathcal{H}W_{k-1} + 4\mathcal{H}W_{k-2} & \mathcal{H}W_k & \mathcal{H}W_{k-1} \end{vmatrix} \\
&= 5 \begin{vmatrix} \mathcal{H}W_{k+2} & \mathcal{H}W_{k+2} & \mathcal{H}W_{k+1} \\ \mathcal{H}W_{k+1} & \mathcal{H}W_{k+1} & \mathcal{H}W_k \\ \mathcal{H}W_k & \mathcal{H}W_k & \mathcal{H}W_{k-1} \end{vmatrix} - 8 \begin{vmatrix} \mathcal{H}W_{k+1} & \mathcal{H}W_{k+2} & \mathcal{H}W_{k+1} \\ \mathcal{H}W_k & \mathcal{H}W_{k+1} & \mathcal{H}W_k \\ \mathcal{H}W_{k-1} & \mathcal{H}W_k & \mathcal{H}W_{k-1} \end{vmatrix} \\
&\quad + 4 \begin{vmatrix} \mathcal{H}W_k & \mathcal{H}W_{k+2} & \mathcal{H}W_{k+1} \\ \mathcal{H}W_{k-1} & \mathcal{H}W_{k+1} & \mathcal{H}W_k \\ \mathcal{H}W_{k-2} & \mathcal{H}W_k & \mathcal{H}W_{k-1} \end{vmatrix} \\
&= 4 \begin{vmatrix} \mathcal{H}W_{k+2} & \mathcal{H}W_{k+1} & \mathcal{H}W_k \\ \mathcal{H}W_{k+1} & \mathcal{H}W_k & \mathcal{H}W_{k-1} \\ \mathcal{H}W_k & \mathcal{H}W_{k-1} & \mathcal{H}W_{k-2} \end{vmatrix} = 4^{k+1} \begin{vmatrix} \mathcal{H}W_2 & \mathcal{H}W_1 & \mathcal{H}W_0 \\ \mathcal{H}W_1 & \mathcal{H}W_0 & \mathcal{H}W_{-1} \\ \mathcal{H}W_0 & \mathcal{H}W_{-1} & \mathcal{H}W_{-2} \end{vmatrix}.
\end{aligned}$$

The other case can be done similarly. Thus, the proof is finished. \square

From previous theorem, we get following corollary.

COROLLARY 6. (*Simpson's formula for hyperbolic generalized Woodall sequence's special cases*)

$$\begin{aligned}
\text{(a): } & \begin{vmatrix} \mathcal{H}G_{k+2} & \mathcal{H}G_{k+1} & \mathcal{H}G_k \\ \mathcal{H}G_{k+1} & \mathcal{H}G_k & \mathcal{H}G_{k-1} \\ \mathcal{H}G_k & \mathcal{H}G_{k-1} & \mathcal{H}G_{k-2} \end{vmatrix} = -4^{n-1}(9 + 9j). \\
\text{(b): } & \begin{vmatrix} \mathcal{H}H_{k+2} & \mathcal{H}H_{k+1} & \mathcal{H}H_k \\ \mathcal{H}H_{k+1} & \mathcal{H}H_k & \mathcal{H}H_{k-1} \\ \mathcal{H}H_k & \mathcal{H}H_{k-1} & \mathcal{H}H_{k-2} \end{vmatrix} = 0. \\
\text{(c): } & \begin{vmatrix} \mathcal{H}R_{k+2} & \mathcal{H}R_{k+1} & \mathcal{H}R_k \\ \mathcal{H}R_{k+1} & \mathcal{H}R_k & \mathcal{H}R_{k-1} \\ \mathcal{H}R_k & \mathcal{H}R_{k-1} & \mathcal{H}R_{k-2} \end{vmatrix} = 4^{n-1}(9 + 9j). \\
\text{(d): } & \begin{vmatrix} \mathcal{H}C_{k+2} & \mathcal{H}C_{k+1} & \mathcal{H}C_k \\ \mathcal{H}C_{k+1} & \mathcal{H}C_k & \mathcal{H}C_{k-1} \\ \mathcal{H}C_k & \mathcal{H}C_{k-1} & \mathcal{H}C_{k-2} \end{vmatrix} = -4^{n-1}(9 + 9j).
\end{aligned}$$

THEOREM 7. (*Catalan's identity*) For all integers n and m , the following identity holds

$$\mathcal{H}W_{n+m}\mathcal{H}W_{n-m} - \mathcal{H}W_n^2 = 2^{n-m}(-2^{m+n}m^2\hat{\alpha}^2A_2^2 + A_2A_3(-2^{m+1}\hat{\beta}\hat{\gamma} + \hat{\beta}\hat{\gamma} + 2^{2m}\hat{\beta}\hat{\gamma} - m\hat{\alpha}\hat{\gamma} + n\hat{\alpha}\hat{\gamma} - 2^{m+1}n\hat{\alpha}\hat{\gamma} + 2^{2m}m\hat{\alpha}\hat{\gamma} + 2^{2m}n\hat{\alpha}\hat{\gamma})) + A_1A_3(\hat{\alpha}\hat{\gamma} - 2^{m+1}\hat{\alpha}\hat{\gamma} + 2^{2m}\hat{\alpha}\hat{\gamma}).$$

Proof. Using the Binet's formula $\mathcal{H}W_n = (A_1\hat{\alpha} + A_2\hat{\beta} + A_2n\hat{\alpha})2^n + A_3\hat{\gamma}$, we get the required identity. \square

As special cases of the above theorem, we give Catalan's identity of hyperbolic modified Woodall, hyperbolic modified Cullen, hyperbolic Woodall and hyperbolic Cullen numbers. **Firstly, we present Catalan's identity of hyperbolic Woodall numbers.**

COROLLARY 8. (*Catalan's identity for the hyperbolic modified Woodall numbers*) For all integers n and m , the following identity holds

$$\begin{aligned} \mathcal{H}G_{n+m}\mathcal{H}G_{n-m} - \mathcal{H}G_n^2 &= -2^{n-m}(\widehat{\alpha}\widehat{\gamma} - \widehat{\beta}\widehat{\gamma} + 2^{2m}\widehat{\alpha}\widehat{\gamma} - 2^{2m}\widehat{\beta}\widehat{\gamma} - 2^{m+1}\widehat{\alpha}\widehat{\gamma} + 2^{m+1}\widehat{\beta}\widehat{\gamma} \\ &\quad + m\widehat{\alpha}\widehat{\gamma} - n\widehat{\alpha}\widehat{\gamma} + 2^{m+n}m^2\widehat{\alpha}^2 - 2^{2m}m\widehat{\alpha}\widehat{\gamma} - 2^{2m}n\widehat{\alpha}\widehat{\gamma} + 2^{m+1}n\widehat{\alpha}\widehat{\gamma}). \end{aligned}$$

Proof. Take $W_n = G_n$ in Theorem 7. \square

Secondly, we give Catalan's identity of hyperbolic modified Cullen numbers.

COROLLARY 9. (*Catalan's identity for the hyperbolic modified Cullen numbers*) For all integers n and m , the following identity holds

$$\mathcal{H}H_{n+m}\mathcal{H}H_{n-m} - \mathcal{H}H_n^2 = 2^{n-m}(2\widehat{\alpha}\widehat{\gamma} + 2 \times 2^{2m}\widehat{\alpha}\widehat{\gamma} - 2 \times 2^{m+1}\widehat{\alpha}\widehat{\gamma}).$$

Proof. Take $W_n = H_n$ in Theorem 7. \square

Thirdly, we give Catalan's identity of hyperbolic Woodall numbers.

COROLLARY 10. (*Catalan's identity for the hyperbolic Woodall numbers*) For all integers n and m , the following identity holds

$$\begin{aligned} \mathcal{H}R_{n+m}\mathcal{H}R_{n-m} - \mathcal{H}R_n^2 &= -2^{n-m}(\widehat{\beta}\widehat{\gamma} + 2^{2m}\widehat{\beta}\widehat{\gamma} - 2^{m+1}\widehat{\beta}\widehat{\gamma} - m\widehat{\alpha}\widehat{\gamma} + n\widehat{\alpha}\widehat{\gamma} + 2^{m+n}m^2\widehat{\alpha}^2 \\ &\quad + 2^{2m}m\widehat{\alpha}\widehat{\gamma} + 2^{2m}n\widehat{\alpha}\widehat{\gamma} - 2^{m+1}n\widehat{\alpha}\widehat{\gamma}). \end{aligned}$$

Proof. Take $W_n = R_n$ in Theorem 7. \square

Fourthly, we give Catalan's identity of hyperbolic Cullen numbers.

COROLLARY 11. (*Catalan's identity for the hyperbolic Cullen numbers*) For all integers n and m , the following identity holds

$$\begin{aligned} \mathcal{H}C_{n+m}\mathcal{H}C_{n-m} - \mathcal{H}C_n^2 &= 2^{n-m}(\widehat{\beta}\widehat{\gamma} + 2^{2m}\widehat{\beta}\widehat{\gamma} - 2^{m+1}\widehat{\beta}\widehat{\gamma} - m\widehat{\alpha}\widehat{\gamma} + n\widehat{\alpha}\widehat{\gamma} - 2^{m+n}m^2\widehat{\alpha}^2 \\ &\quad + 2^{2m}m\widehat{\alpha}\widehat{\gamma} + 2^{2m}n\widehat{\alpha}\widehat{\gamma} - 2^{m+1}n\widehat{\alpha}\widehat{\gamma}). \end{aligned}$$

Proof. Take $W_n = C_n$ in Theorem 7. \square

Note that for $m = 1$ in Catalan's identity, we get the Cassini's identity for the hyperbolic generalized Woodall sequence.

COROLLARY 12. (*Cassini's identity*) For all integers n , the following identity holds

$$\mathcal{H}W_{n+1}\mathcal{H}W_{n-1} - \mathcal{H}W_n^2 = 2^{n-1}(A_2A_3(3\widehat{\alpha}\widehat{\gamma} + \widehat{\beta}\widehat{\gamma} + n\widehat{\alpha}\widehat{\gamma}) - 2^{n+1}A_2^2\widehat{\alpha}^2 + A_1A_3\widehat{\alpha}\widehat{\gamma}).$$

As special cases of Cassini's identity, we give Cassini's identity of hyperbolic modified Woodall, hyperbolic modified Cullen, hyperbolic Woodall and hyperbolic Cullen numbers. Firstly, we present Cassini's identity of hyperbolic modified Woodall numbers.

Firstly, we give Cassini's identity of hyperbolic modified Woodall numbers.

COROLLARY 13. (*Cassini's identity of hyperbolic modified Woodall numbers*) For all integers n , the following identity holds

$$\mathcal{H}G_{n+1}\mathcal{H}G_{n-1} - \mathcal{H}G_n^2 = 2^{n-1}(2\widehat{\alpha}\widehat{\gamma} + \widehat{\beta}\widehat{\gamma} - 2^{n+1}\widehat{\alpha}^2 + n\widehat{\alpha}\widehat{\gamma}).$$

Secondly, we give Cassini's identity of hyperbolic modified Cullen numbers.

COROLLARY 14. (*Cassini's identity of hyperbolic modified Cullen numbers*) For all integers n , the following identity holds

$$\mathcal{H}H_{n+1}\mathcal{H}H_{n-1} - \mathcal{H}H_n^2 = 2^n\widehat{\alpha}\widehat{\gamma}.$$

Fourth, we give Cassini's identity of hyperbolic Woodall numbers.

COROLLARY 15. (*Cassini's identity of hyperbolic Woodall numbers*) For all integers n , the following identity holds

$$\mathcal{H}R_{n+1}\mathcal{H}R_{n-1} - \mathcal{H}R_n^2 = -2^{n-1}(3\widehat{\alpha}\widehat{\gamma} + \widehat{\beta}\widehat{\gamma} + 2^{n+1}\widehat{\alpha}^2 + n\widehat{\alpha}\widehat{\gamma}).$$

Third, we give Cassini's identity of hyperbolic Cullen numbers.

COROLLARY 16. (*Cassini's identity of hyperbolic Cullen numbers*) For all integers n , the following identity holds

$$\mathcal{H}C_{n+1}\mathcal{H}C_{n-1} - \mathcal{H}C_n^2 = 2^{n-1}(3\widehat{\alpha}\widehat{\gamma} + \widehat{\beta}\widehat{\gamma} - 2^{n+1}\widehat{\alpha}^2 + n\widehat{\alpha}\widehat{\gamma}).$$

THEOREM 17. For all integers m, n , G_n is woodall numbers, the following identity is true:

$$\mathcal{H}W_{n+m} = \mathcal{H}W_n G_{m+1} + \mathcal{H}W_{n-1}(-8G_m + 4G_{m-1}) + 4\mathcal{H}W_{n-2}G_m.$$

Proof. The identity (17) can be proved by mathematical induction on m . First of all, we assume that $m \geq 0$ and $n \geq 0$. If $m = 0$ we get

$$\mathcal{H}W_n = \mathcal{H}W_n G_1 + \mathcal{H}W_{n-1}(-8G_0 + 4G_{-1}) + 4\mathcal{H}W_{n-2}G_0$$

which is true by seeing that $G_{-1} = 0$, $G_{-2} = \frac{1}{4}$, $G_{-3} = \frac{1}{2}$. We assume that the identity given holds for $m = k$. For $m = k + 1$, we get

$$\begin{aligned}
\mathcal{H}W_{(k+1)+n} &= 5\mathcal{H}W_{n+k} - 8\mathcal{H}W_{n+k-1} + 4\mathcal{H}W_{n+k-2} \\
&= 5(\mathcal{H}W_n G_{k+1} + \mathcal{H}W_{n-1}(-8G_k + 4G_{k-1}) + 4\mathcal{H}W_{n-2}G_k) \\
&\quad - 8(\mathcal{H}W_n G_k + \mathcal{H}W_{n-1}(-8G_{k-1} + 4G_{k-2}) + 4\mathcal{H}W_{n-2}G_{k-1}) \\
&\quad + 4(\mathcal{H}W_n G_{k-1} + \mathcal{H}W_{n-1}(-8G_{k-2} + 4G_{k-3}) + 4\mathcal{H}W_{n-2}G_{k-2}) \\
&= \mathcal{H}W_n(5G_{k+1} - 8G_k + 4G_{k-1}) + \mathcal{H}W_{n-1}(-8(5G_k - 8G_{k-1} + 4G_{k-2}) \\
&\quad + 4(5G_{k-1} - 8G_{k-2} + 4G_{k-3})) + 4\mathcal{H}W_{n-2}(5G_k - 8G_{k-1} + 4G_{k-2}) \\
&= \mathcal{H}W_n G_{k+2} + \mathcal{H}W_{n-1}(-8G_{k+1} + 4G_k) + 4\mathcal{H}W_{n-2}G_{k+1} \\
&= \mathcal{H}W_n G_{(k+1)+1} + \mathcal{H}W_{n-1}(-8G_{(k+1)} + 4G_{(k+1)-1}) + 4\mathcal{H}W_{n-2}G_{(k+1)}.
\end{aligned}$$

Consequently, by mathematical induction on m , this proves (17). Similarly, we can show for the other cases.

So the proof is finished. \square

4. Linear Sums For Hyperbolic Generalized Woodall Numbers

In this section, we give the summation formulas of the hyperbolic generalized Woodall numbers with positive and negativ subscripts. Now, we present the summation formulas of the generalized Woodall numbers.

PROPOSITION 18. *For the generalized Woodall numbers, we have the following formulas:*

- $\sum_{k=0}^n W_k = \frac{1}{2}W_2(2n - 2^{n+1}(n-1) + 2^{n+2}(n-2) + 6) - \frac{1}{2}W_1(8n - 2^{n+1}(3n-5) + 2^{n+2}(3n-8) + 22) + W_0(4n - 2^{n+1}(n-2) + 2^{n+2}(n-3) + 9).$
- $\sum_{k=0}^n W_{k+1} = \frac{1}{2}W_2(2n + 2^{n+3}(n-1) - 2^{n+2}n + 8) - \frac{1}{2}W_1(8n - 2^{n+2}(3n-2) + 2^{n+3}(3n-5) + 30) + W_0(4n - 2^{n+2}(n-1) + 2^{n+3}(n-2) + 12).$
- $\sum_{k=0}^n W_{k+2} = \frac{1}{2}W_2(2n - 2^{n+3}(n+1) + 2^{n+4}n + 10) + W_0(4n + 2^{n+4}(n-1) - 2^{n+3}n + 16) - \frac{1}{2}W_1(8n - 2^{n+3}(3n+1) + 2^{n+4}(3n-2) + 40).$
- $\sum_{k=0}^n W_{k+3} = W_0(4n - 2^{n+4}(n+1) + 2^{n+5}n + 20) - \frac{1}{2}W_1(8n + 2^{n+5}(3n+1) - 2^{n+4}(3n+4) + 48) + \frac{1}{2}W_2(2n - 2^{n+4}(n+2) + 2^{n+5}(n+1) + 10).$

Proof. For the proof, see Soykan [52]. \square

PROPOSITION 19. *For the generalized Woodall numbers, we have the following formulas:*

- $\sum_{k=0}^n W_{2k} = \frac{1}{9}W_0(36n - 2^{2n+2}(2n-1) + 2^{2n+4}(2n-3) + 53) - \frac{1}{18}W_1(72n - 2^{2n+2}(6n-2) + 2^{2n+4}(6n-8) + 120) + \frac{1}{18}W_2(18n + 2^{2n+4}(2n-2) - 2 \times 2^{2n+2}n + 32).$
- $\sum_{k=0}^n W_{2k+1} = \frac{1}{18}W_2(18n - 2^{2n+3}(2n+1) + 2^{2n+5}(2n-1) + 40) - \frac{1}{18}W_1(72n - 2^{2n+3}(6n+1) + 2^{2n+5}(6n-5) + 150) + \frac{1}{9}W_0(36n + 2^{2n+5}(2n-2) - 2 \times 2^{2n+3}n + 64).$

- $\sum_{k=0}^n W_{2k+2} = \frac{1}{9}W_0(36n - 2^{2n+4}(2n+1) + 2^{2n+6}(2n-1) + 80) - \frac{1}{18}W_1(72n - 2^{2n+4}(6n+4) + 2^{2n+6}(6n-2) + 192) + \frac{1}{18}W_2(18n - 2^{2n+4}(2n+2) + 2 \times 2^{2n+6}n + 50)$.
- $\sum_{k=0}^n W_{2k+3} = \frac{1}{18}W_2((18n - 2^{2n+5}(2n+3) + 2^{2n+7}(2n+1) + 58) - \frac{1}{18}W_1(72n + 2^{2n+7}(6n+1) - 2^{2n+5}(6n+7) + 240) + \frac{1}{9}W_0(36n - 2^{2n+5}(2n+2) + 2 \times 2^{2n+7}n + 100)$.
- $\sum_{k=0}^n W_{2k+4} = \frac{1}{18}W_2(18n - 2^{2n+6}(2n+4) + 2^{2n+8}(2n+2) + 50) + \frac{1}{9}W_0(36n - 2^{2n+6}(2n+3) + 2^{2n+8}(2n+1) + 116) - \frac{1}{18}W_1(72n + 2^{2n+8}(6n+4) - 2^{2n+6}(6n+10) + 264)$.

Proof. For the proof, see Soykan [52]. \square

PROPOSITION 20. *For the generalized Woodall numbers, we have the following formulas:*

- $\sum_{k=0}^n W_{-k} = 4W_0(n + \frac{1}{2^{n+1}}(n+4) - \frac{1}{2^{n+2}}(n+3) - 1) + 2W_1(\frac{1}{2^{n+2}}(3n+8) - 2n - \frac{1}{2^{n+1}}(3n+11) + \frac{7}{2}) + 2W_2(\frac{1}{2}n + \frac{1}{2^{n+1}}(n+3) - \frac{1}{2^{n+2}}(n+2) - 1)$.
- $\sum_{k=0}^n W_{-k+1} = 2W_2(\frac{1}{2}n + \frac{1}{2^n}(n+2) - \frac{1}{2^{n+1}}(n+1) - \frac{3}{2}) + 4W_0(n + \frac{1}{2^n}(n+3) - \frac{1}{2^{n+1}}(n+2) - 2) + 2W_1(\frac{1}{2^{n+1}}(3n+5) - 2n - \frac{1}{2^n}(3n+8) + 6)$.
- $\sum_{k=0}^n W_{-k+2} = 2W_2(\frac{1}{2}n + 2^{1-n}(n+1) - \frac{1}{2^n}n - \frac{3}{2}) + 4W_0(n - \frac{1}{2^n}(n+1) + 2^{1-n}(n+2) - 3) - 2W_1(2n + 2^{1-n}(3n+5) - \frac{1}{2^n}(3n+2) - 8)$.
- $\sum_{k=0}^n W_{-k+3} = 2W_2(\frac{1}{2}n + 2^{2-n}n - 2^{1-n}(n-1) + \frac{1}{2}) + 2W_1(2^{1-n}(3n-1) - 2n - 2^{2-n}(3n+2) + 6) + 4W_0(n - 2^{1-n}n + 2^{2-n}(n+1) - 3)$.

Proof. For the proof, see Soykan [52]. \square

PROPOSITION 21. *For the generalized Woodall numbers, we have the following formulas:*

- $\sum_{k=0}^n W_{-2k} = \frac{8}{9}W_1(\frac{1}{2^{2n+4}}(6n+8) - \frac{9}{2}n - \frac{1}{2^{2n+2}}(6n+14) + 3) + \frac{16}{9}W_0(\frac{9}{4}n + \frac{1}{2^{2n+2}}(2n+5) - \frac{1}{2^{2n+4}}(2n+3) - \frac{1}{2}) + \frac{8}{9}W_2(\frac{9}{8}n + \frac{1}{2^{2n+2}}(2n+4) - \frac{1}{2^{2n+4}}(2n+2) - \frac{7}{8})$.
- $\sum_{k=0}^n W_{-2k+1} = \frac{8}{9}W_1(\frac{1}{2^{2n+3}}(6n+5) - \frac{9}{2}n - \frac{1}{2^{2n+1}}(6n+11) + 6) + \frac{16}{9}W_0(\frac{9}{4}n + \frac{1}{2^{2n+1}}(2n+4) - \frac{1}{2^{2n+3}}(2n+2) - \frac{7}{4}) + \frac{8}{9}W_2(\frac{9}{8}n + \frac{1}{2^{2n+1}}(2n+3) - \frac{1}{2^{2n+3}}(2n+1) - \frac{11}{8})$.
- $\sum_{k=0}^n W_{-2k+2} = \frac{8}{9}W_2(\frac{9}{8}n - \frac{2}{2^{2n+2}}n + \frac{1}{2^{2n}}(2n+2) - \frac{7}{8}) - \frac{16}{9}W_0(\frac{1}{2^{2n+2}}(2n+1) - \frac{9}{4}n - \frac{1}{2^{2n}}(2n+3) + \frac{11}{4}) + \frac{8}{9}W_1(\frac{1}{2^{2n+2}}(6n+2) - \frac{9}{2}n - \frac{1}{2^{2n}}(6n+8) + \frac{15}{2})$.
- $\sum_{k=0}^n W_{-2k+3} = \frac{8}{9}W_1(\frac{1}{2^{2n+1}}(6n-1) - \frac{9}{2}n - 2^{1-2n}(6n+5) + \frac{3}{2}) + \frac{8}{9}W_2(\frac{9}{8}n - \frac{1}{2^{2n+1}}(2n-1) + 2^{1-2n}(2n+1) + \frac{25}{8}) + \frac{16}{9}W_0(\frac{9}{4}n + 2^{1-2n}(2n+2) - \frac{2}{2^{2n+1}}n - \frac{7}{4})$.
- $\sum_{k=0}^n W_{-2k+4} = \frac{8}{9}W_2(\frac{9}{8}n + 2 \times 2^{2-2n}n - \frac{1}{2^{2n}}(2n-2) + \frac{137}{8}) + \frac{16}{9}W_0(\frac{9}{4}n + 2^{2-2n}(2n+1) - \frac{1}{2^{2n}}(2n-1) + \frac{25}{4}) - \frac{8}{9}W_1(\frac{9}{2}n + 2^{2-2n}(6n+2) - \frac{1}{2^{2n}}(6n-4) + \frac{57}{2})$.

Proof. For the proof, see Soykan [52]. \square

Now, we present the formulas which give the summation of the hyperbolic generalized Woodall numbers.

THEOREM 22. *For $n \geq 0$, hyperbolic generalized Woodall numbers have the following formulas:*

$$(a): \sum_{k=0}^n \mathcal{H}W_k = (3+n-3 \times 2^n + 2^n n + 4j + jn - 2^{n+2}j + 2^{n+1}jn)W_2 + (-11-4n+11 \times 2^n - 3 \times 2^n n - 15j - 4jn + 2^{n+4}j - 3 \times 2^{n+1}jn)W_1 + (9+4n-2^{n+3} + 2^{n+1}n + 12j + 4jn - 3 \times 2^{n+2}j + 2^{n+2}jn)W_0.$$

$$\begin{aligned}
\text{(b): } \sum_{k=0}^n \mathcal{H}W_{2k} &= \left(\frac{16}{9} + n - \frac{1}{9}2^{2n+4} + \frac{1}{3}2^{2n+2}n + \frac{20}{9}j + jn - \frac{5}{9}2^{2n+2}j + \frac{1}{3}2^{2n+3}jn\right)W_2 + \left(-\frac{20}{3} - 4n + \frac{5}{3}2^{2n+2} - 2^{2n+2}n - \frac{25}{3}j + \frac{7}{3}2^{2n+2}j - 4jn - 2^{2n+3}jn\right)W_1 + \left(\frac{53}{9} - \frac{11}{9}2^{2n+2} + 4n + \frac{1}{3}2^{2n+3}n + \frac{64}{9}j - \frac{1}{9}2^{2n+6}j + 4jn + \frac{1}{3}2^{2n+4}jn\right)W_0. \\
\text{(c): } \sum_{k=0}^n \mathcal{H}W_{2k+1} &= \left(\frac{20}{9} - \frac{5}{9}2^{2n+2} + n + \frac{1}{3}2^{2n+3}n + \frac{25}{9}j - \frac{1}{9}2^{2n+4}j + jn + \frac{1}{3}2^{2n+4}jn\right)W_2 + \left(-\frac{25}{3} + \frac{7}{3}2^{2n+2} - 4n - 2^{2n+3}n + \frac{1}{3}2^{2n+5}j - \frac{32}{3}j - 4jn - 2^{2n+4}jn\right)W_1 + \left(\frac{64}{9} - \frac{1}{9}2^{2n+6} + 4n + \frac{1}{3}2^{2n+4}n + \frac{80}{9}j - \frac{5}{9}2^{2n+4}j + 4jn + \frac{1}{3}2^{2n+5}jn\right)W_0.
\end{aligned}$$

Proof. Proof can be obtained by using Proposition 21.

(a): We can derive the following using the formulas in Proposition 18.

$$\sum_{k=0}^n \mathcal{H}W_k = \sum_{k=0}^n W_k + j \sum_{k=0}^n W_{k+1}.$$

$$\begin{aligned}
&\sum_{k=0}^n \mathcal{H}W_k \\
&= \frac{1}{2}W_2(2n - 2^{n+1}(n-1) + 2^{n+2}(n-2) + 6) - \frac{1}{2}W_1(8n - 2^{n+1}(3n-5) + 2^{n+2}(3n-8) + 22) + W_0(4n - 2^{n+1}(n-2) + 2^{n+2}(n-3) + 9) \\
&\quad + j\left(\frac{1}{2}W_2(2n + 2^{n+3}(n-1) - 2^{n+2}n + 8) - \frac{1}{2}W_1(8n - 2^{n+2}(3n-2) + 2^{n+3}(3n-5) + 30) + W_0(4n - 2^{n+2}(n-1) + 2^{n+3}(n-2) + 12)\right). \\
&\sum_{k=0}^n \mathcal{H}W_k \\
&= (3 + n - 3 \times 2^n + 2^n n + 4j + jn - 2^{n+2}j + 2^{n+1}jn)W_2 \\
&\quad + (-11 - 4n + 11 \times 2^n - 3 \times 2^n n - 15j - 4jn + 2^{n+4}j - 3 \times 2^{n+1}jn)W_1 \\
&\quad + (9 + 4n - 2^{n+3} + 2^{n+1}n + 12j + 4jn - 3 \times 2^{n+2}j + 2^{n+2}jn)W_0.
\end{aligned}$$

The proof is finished. \square

(b): We can derive the following using the formulas in Proposition 19.

$$\sum_{k=0}^n \mathcal{H}W_{2k} = \sum_{k=0}^n W_{2k} + j \sum_{k=0}^n W_{2k+1}.$$

$$\begin{aligned}
&\sum_{k=0}^n \mathcal{H}W_{2k} \\
&= \frac{1}{9}W_0(36n - 2^{2n+2}(2n-1) + 2^{2n+4}(2n-3) + 53) - \frac{1}{18}W_1(72n - 2^{2n+2}(6n-2) + 2^{2n+4}(6n-8) + 120) + \frac{1}{18}W_2(18n + 2^{2n+4}(2n-2) - 2 \times 2^{2n+2}n + 32) \\
&\quad + j\left(\frac{1}{18}W_2(18n - 2^{2n+3}(2n+1) + 2^{2n+5}(2n-1) + 40) - \frac{1}{18}W_1(72n - 2^{2n+3}(6n+1) + 2^{2n+5}(6n-5) + 150) + \frac{1}{9}W_0(36n + 2^{2n+5}(2n-2) - 2 \times 2^{2n+3}n + 64)\right).
\end{aligned}$$

$$\begin{aligned}
& \sum_{k=0}^n \mathcal{H}W_{2k} \\
&= \left(\frac{16}{9} + n - \frac{1}{9}2^{2n+4} + \frac{1}{3}2^{2n+2}n + \frac{20}{9}j + jn - \frac{5}{9}2^{2n+2}j + \frac{1}{3}2^{2n+3}jn \right) W_2 \\
&\quad + \left(-\frac{20}{3} - 4n + \frac{5}{3}2^{2n+2} - 2^{2n+2}n - \frac{25}{3}j + \frac{7}{3}2^{2n+2}j - 4jn - 2^{2n+3}jn \right) W_1 \\
&\quad + \left(\frac{53}{9} - \frac{11}{9}2^{2n+2} + 4n + \frac{1}{3}2^{2n+3}n + \frac{64}{9}j - \frac{1}{9}2^{2n+6}j + 4jn + \frac{1}{3}2^{2n+4}jn \right) W_0.
\end{aligned}$$

The proof is completed. \square

(c): We can derive the following using the formulas in Proposition 21.

$$\sum_{k=0}^n \mathcal{H}W_{2k+1} = \sum_{k=0}^n W_{2k+1} + j \sum_{k=0}^n W_{2k+2}.$$

$$\begin{aligned}
& \sum_{k=0}^n \mathcal{H}W_{2k+1} \\
&= \frac{1}{18} W_2(18n - 2^{2n+3}(2n+1) + 2^{2n+5}(2n-1) + 40) - \frac{1}{18} W_1(72n - 2^{2n+3}(6n+1) \\
&\quad + 2^{2n+5}(6n-5) + 150) + \frac{1}{9} W_0(36n + 2^{2n+5}(2n-2) - 2 \times 2^{2n+3}n + 64) \\
&\quad + j \left(\frac{1}{9} W_0(36n - 2^{2n+4}(2n+1) + 2^{2n+6}(2n-1) + 80) - \frac{1}{18} W_1(72n - 2^{2n+4}(6n+4) \right. \\
&\quad \left. + 2^{2n+6}(6n-2) + 192) + \frac{1}{18} W_2(18n - 2^{2n+4}(2n+2) + 2 \times 2^{2n+6}n + 50) \right).
\end{aligned}$$

$$\begin{aligned}
& \sum_{k=0}^n \mathcal{H}W_{2k+1} \\
&= \left(\frac{20}{9} - \frac{5}{9}2^{2n+2} + n + \frac{1}{3}2^{2n+3}n + \frac{25}{9}j - \frac{1}{9}2^{2n+4}j + jn + \frac{1}{3}2^{2n+4}jn \right) W_2 \\
&\quad + \left(-\frac{25}{3} + \frac{7}{3}2^{2n+2} - 4n - 2^{2n+3}n + \frac{1}{3}2^{2n+5}j - \frac{32}{3}j - 4jn - 2^{2n+4}jn \right) W_1 \\
&\quad + \left(\frac{64}{9} - \frac{1}{9}2^{2n+6} + 4n + \frac{1}{3}2^{2n+4}n + \frac{80}{9}j - \frac{5}{9}2^{2n+4}j + 4jn + \frac{1}{3}2^{2n+5}jn \right) W_0.
\end{aligned}$$

The proof is finished. \square

As a first special case of the above theorem, we have the following summation formulas for hyperbolic Woodall numbers:

COROLLARY 23. For $n \geq 0$, hyperbolic modified Woodall numbers have the following properties:

- (a): $\sum_{k=0}^n \mathcal{H}G_k = 4 + n + 2^{n+1}n - 2^{n+2} + j(5 - 5 \times 2^{n+2} + n + 2^{n+4} + 2^{n+2}n)$.
- (b): $\sum_{k=0}^n \mathcal{H}G_{2k} = \frac{20}{9} + n + \frac{2}{3}2^{2n+2}n + \frac{5}{3}2^{2n+2} - \frac{5}{9}2^{2n+4} + j(\frac{25}{9} - \frac{4}{9}2^{2n+2} + n + \frac{2}{3}2^{2n+3}n)$.
- (c): $\sum_{k=0}^n \mathcal{H}G_{2k+1} = \frac{25}{9} + n + \frac{2}{3}2^{2n+3}n - \frac{4}{9}2^{2n+2} + j(\frac{29}{9} - \frac{5}{9}2^{2n+4} + \frac{1}{3}2^{2n+5} + n + \frac{2}{3}2^{2n+4}n)$.

As a second special case of the above theorem, we have the following summation formulas for hyperbolic modified Cullen numbers:

COROLLARY 24. For $n \geq 0$, hyperbolic modified Cullen numbers have the following properties:

- (a): $\sum_{k=0}^n \mathcal{H}H_k = -1 + n - 6 \times 2^n n - 3 \times 2^{n+3} + 3 \times 2^{n+1} n + 28 \times 2^n + j(-3 - 18 \times 2^{n+2} + 5 \times 2^{n+4} + n - 6 \times 2^{n+1} n + 3 \times 2^{n+2} n)$.
- (b): $\sum_{k=0}^n \mathcal{H}H_{2k} = \frac{1}{3} + n - 2^{2n+3} n + 2^{2n+3} n + \frac{14}{3} 2^{2n+2} - 2^{2n+4} + j(-\frac{1}{3} + \frac{20}{3} 2^{2n+2} - \frac{1}{3} 2^{2n+6} + n - 2^{2n+4} n + 2^{2n+4} n)$.
- (c): $\sum_{k=0}^n \mathcal{H}H_{2k+1} = -\frac{1}{3} + n - 2 \times 2^{2n+3} n + 2^{2n+4} n + \frac{20}{3} 2^{2n+2} - \frac{1}{3} 2^{2n+6} + j(-\frac{5}{3} - \frac{8}{3} 2^{2n+4} + \frac{5}{3} 2^{2n+5} + n - 2^{2n+5} n + 2^{2n+5} n)$.

As a third special case of the above theorem, we have the following summation formulas for hyperbolic Woodall numbers:

COROLLARY 25. *For $n \geq 0$, hyperbolic Woodall numbers have the following properties:*

- (a): $\sum_{k=0}^n \mathcal{H}R_k = 1 - n + 4 \times 2^n n + 2^{n+3} - 2^{n+1} n - 10 \times 2^n + j(1 - 2^{n+4} + 2^{n+4} - n + 2^{n+3} n - 2^{n+2} n)$.
- (b): $\sum_{k=0}^n \mathcal{H}R_{2k} = -\frac{1}{9} - n + \frac{4}{3} 2^{2n+2} n - \frac{1}{3} 2^{2n+3} n + \frac{26}{9} 2^{2n+2} - \frac{7}{9} 2^{2n+4} + j(\frac{1}{9} - n - \frac{14}{9} 2^{2n+2} + \frac{1}{9} 2^{2n+6} + \frac{4}{3} 2^{2n+3} n - \frac{1}{3} 2^{2n+4} n)$.
- (c): $\sum_{k=0}^n \mathcal{H}R_{2k+1} = \frac{1}{9} - n + \frac{4}{3} 2^{2n+3} n - \frac{1}{3} 2^{2n+4} n - \frac{14}{9} 2^{2n+2} + \frac{1}{9} 2^{2n+6} + j(-\frac{1}{9} + \frac{1}{3} 2^{2n+5} - \frac{2}{9} 2^{2n+4} - n + \frac{4}{3} 2^{2n+4} n - \frac{1}{3} 2^{2n+5} n)$.

As a fourth special case of the above theorem, we have the following summation formulas for hyperbolic Cullen numbers:

COROLLARY 26. *For $n \geq 0$, hyperbolic Cullen numbers have the following properties:*

- (a): $\sum_{k=0}^n \mathcal{H}C_k = 3 + n - 2^{n+3} + 2^{n+1} n + 6 \times 2^n + j(3 + n + 2^{n+2} n)$.
- (b): $\sum_{k=0}^n \mathcal{H}C_{2k} = \frac{17}{9} + n + \frac{1}{3} 2^{2n+3} n - \frac{2}{9} 2^{2n+2} + j(\frac{19}{9} + n + \frac{1}{9} 2^{2n+3} + \frac{1}{3} 2^{2n+4} n)$.
- (c): $\sum_{k=0}^n \mathcal{H}C_{2k+1} = \frac{19}{9} + n + \frac{1}{3} 2^{2n+4} n + \frac{1}{9} 2^{2n+3} + j(\frac{17}{9} + \frac{4}{9} 2^{2n+4} + n + \frac{1}{3} 2^{2n+5} n)$.

We now introduce the formulas that allow us to find the sum of hyperbolic generalized Woodall numbers with negative subscripts in the following theorem.

THEOREM 27. *For $n \geq 0$, hyperbolic generalized Woodall numbers have the following formulas:*

- (a): $\sum_{k=0}^n \mathcal{H}W_{-k} = (-2 + \frac{2}{2^n} - 3j + n + \frac{3}{2^n} j + \frac{1}{2 \times 2^n} n + jn + \frac{1}{2^n} jn)W_2 + (7 - \frac{7}{2^n} + 12j - 4n - \frac{11}{2^n} j - \frac{3}{2 \times 2^n} n - 4jn - \frac{3}{2^n} jn)W_1 + (-4 + \frac{5}{2^n} - 8j + 4n + \frac{8}{2^n} j + \frac{1}{2^n} n + 4jn + \frac{2}{2^n} jn)W_0$.
- (b): $\sum_{k=0}^n \mathcal{H}W_{-2k} = (-\frac{7}{9} + \frac{7}{9 \times 2^{2n}} - \frac{11}{9} j + n + \frac{11}{9 \times 2^{2n}} j + \frac{1}{3 \times 2^{2n}} n + jn + \frac{2}{3 \times 2^{2n}} jn)W_2 + (\frac{8}{3} - \frac{8}{3 \times 2^{2n}} + \frac{16}{3} j - 4n - \frac{13}{3 \times 2^{2n}} j - \frac{1}{2^{2n}} n - 4jn - \frac{2}{2^{2n}} jn)W_1 + (-\frac{8}{9} + \frac{17}{9 \times 2^{2n}} - \frac{28}{9} j + 4n + \frac{28}{9 \times 2^{2n}} j + \frac{2}{3 \times 2^{2n}} n + 4jn + \frac{4}{3 \times 2^{2n}} jn)W_0$.
- (c): $\sum_{k=0}^n \mathcal{H}W_{-2k+1} = (-\frac{11}{9} + \frac{11}{9 \times 2^{2n}} - \frac{7}{9} j + n + \frac{16}{9 \times 2^{2n}} j + \frac{2}{3 \times 2^{2n}} n + jn + \frac{4}{3 \times 2^{2n}} jn)W_2 + (\frac{16}{3} - \frac{13}{3 \times 2^{2n}} + \frac{20}{3} j - 4n - \frac{20}{3 \times 2^{2n}} j - \frac{2}{2^{2n}} n - 4jn - \frac{4}{2^{2n}} jn)W_1 + (-\frac{28}{9} + \frac{28}{9 \times 2^{2n}} - \frac{44}{9} j + 4n + \frac{44}{9 \times 2^{2n}} j + \frac{4}{3 \times 2^{2n}} n + 4jn + \frac{8}{3 \times 2^{2n}} jn)W_0$.

Proof. It can be obtained by using Proposition 20.

- (a): We can derive the following using the formulas in Proposition 20.

$$\sum_{k=0}^n \mathcal{H}W_{-k} = \sum_{k=0}^n W_{-k} + j \sum_{k=0}^n W_{-k+1}.$$

$$\begin{aligned}
\sum_{k=0}^n \mathcal{H}W_{-k} &= 4W_0(n + \frac{1}{2^{n+1}}(n+4) - \frac{1}{2^{n+2}}(n+3) - 1) + 2W_1(\frac{1}{2^{n+2}}(3n+8) - 2n - \frac{1}{2^{n+1}}(3n+11) + \frac{7}{2}) \\
&\quad + 2W_2(\frac{1}{2}n + \frac{1}{2^{n+1}}(n+3) - \frac{1}{2^{n+2}}(n+2) - 1) \\
&\quad + j(2W_2(\frac{1}{2}n + \frac{1}{2^n}(n+2) - \frac{1}{2^{n+1}}(n+1) - \frac{3}{2}) + 4W_0(n + \frac{1}{2^n}(n+3) - \frac{1}{2^{n+1}}(n+2) - 2) \\
&\quad + 2W_1(\frac{1}{2^{n+1}}(3n+5) - 2n - \frac{1}{2^n}(3n+8) + 6)).
\end{aligned}$$

$$\begin{aligned}
\sum_{k=0}^n \mathcal{H}W_{-k} &= (-2 + \frac{2}{2^n} - 3j + n + \frac{3}{2^n}j + \frac{1}{2 \times 2^n}n + jn + \frac{1}{2^n}jn)W_2 \\
&\quad + (7 - \frac{7}{2^n} + 12j - 4n - \frac{11}{2^n}j - \frac{3}{2 \times 2^n}n - 4jn - \frac{3}{2^n}jn)W_1 \\
&\quad + (-4 + \frac{5}{2^n} - 8j + 4n + \frac{8}{2^n}j + \frac{1}{2^n}n + 4jn + \frac{2}{2^n}jn)W_0.
\end{aligned}$$

This proves (a). We can prove (b) and (c) similarly way using Proposition 21. \square

As a first special case of the above theorem, we have the following summation formulas for hyperbolic modified Woodall numbers:

COROLLARY 28. *For $n \geq 0$, hyperbolic modified Woodall numbers have the following properties:*

- (a): $\sum_{k=0}^n \mathcal{H}G_{-k} = -3 + n + \frac{n+3}{2^n} + j(-3 + n + \frac{2n+4}{2^n})$.
- (b): $\sum_{k=0}^n \mathcal{H}G_{-2k} = -\frac{11}{9} + n + \frac{11+6n}{9 \times 2^{2n}} + j(-\frac{7}{9} + n + \frac{16+12n}{9 \times 2^{2n}})$.
- (c): $\sum_{k=0}^n \mathcal{H}G_{-2k+1} = -\frac{7}{9} + n + \frac{16+12n}{9 \times 2^{2n}} + j(\frac{25}{9} + n + \frac{20+24n}{9 \times 2^{2n}})$.

As a second special case of the above theorem, we have the following summation formulas for hyperbolic modified Cullen numbers:

COROLLARY 29. *For $n \geq 0$, hyperbolic modified Cullen numbers have the following properties:*

- (a): $\sum_{k=0}^n \mathcal{H}H_{-k} = 5 + n - \frac{2}{2^n} + j(9 - \frac{4}{2^n} + n)$.
- (b): $\sum_{k=0}^n \mathcal{H}H_{-2k} = \frac{11}{3} + n - \frac{2}{3 \times 2^{2n}} + j(\frac{19}{3} - \frac{4}{3 \times 2^{2n}} + n)$.
- (c): $\sum_{k=0}^n \mathcal{H}H_{-2k+1} = \frac{19}{3} + n - \frac{4}{3 \times 2^{2n}} + j(\frac{35}{3} - \frac{8}{3 \times 2^{2n}} + n)$.

As a third special case of the above theorem, we have the following summation formulas for hyperbolic Woodall numbers:

COROLLARY 30. *For $n \geq 0$, hyperbolic Woodall numbers have the following properties:*

- (a): $\sum_{k=0}^n \mathcal{H}R_{-k} = -3 - n + \frac{2+n}{2^n} + j(-1 - n + \frac{2+2n}{2^n})$.
- (b): $\sum_{k=0}^n \mathcal{H}R_{-2k} = -\frac{17}{9} - n + \frac{8+6n}{9 \times 2^{2n}} + j(-\frac{1}{9} - n + \frac{10+12n}{9 \times 2^{2n}})$.
- (c): $\sum_{k=0}^n \mathcal{H}R_{-2k+1} = -\frac{1}{9} - n + \frac{10+12n}{9 \times 2^{2n}} + j(\frac{55}{9} - n + \frac{8+24n}{9 \times 2^{2n}})$.

As a fourth special case of the above theorem, we have the following summation formulas for hyperbolic Cullen numbers:

COROLLARY 31. For $n \geq 0$, hyperbolic Cullen numbers have the following properties:

- (a): $\sum_{k=0}^n \mathcal{H}C_{-k} = -1 + n + \frac{2+n}{2^n} + j(1 + \frac{2+2n}{2^n} + n)$.
- (b): $\sum_{k=0}^n \mathcal{H}C_{-2k} = \frac{1}{9} + n + \frac{8+6n}{9 \times 2^{2n}} + j(\frac{17}{9} + \frac{10+12n}{9 \times 2^{2n}} + n)$.
- (c): $\sum_{k=0}^n \mathcal{H}C_{-2k+1} = \frac{17}{9} + n + \frac{10+12n}{9 \times 2^{2n}} + j(\frac{73}{9} + \frac{8+24n}{9 \times 2^{2n}} + n)$.

5. Matrices related with Hyperbolic Generalized Woodall Numbers

In this section, we present matrices related with hyperbolic generalized Woodall numbers.

Now, $\{G_n\}$ defined by the third-order recurrence relation as follows

$$G_n = 5G_{n-1} - 8G_{n-2} + 4G_{n-3} \text{ with the initial conditions } G_0 = 0, G_1 = 1, G_2 = 5.$$

We present the square matrix A of order 3 as

$$A = \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

such that $\det A = 1$. Then, we give the following Lemma.

LEMMA 32. For all integers n the following identity is true.

$$\begin{pmatrix} \mathcal{H}W_{n+2} \\ \mathcal{H}W_{n+1} \\ \mathcal{H}W_n \end{pmatrix} = \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^n \begin{pmatrix} \mathcal{H}W_2 \\ \mathcal{H}W_1 \\ \mathcal{H}W_0 \end{pmatrix}.$$

Proof. First, we suppose that $n \geq 0$. Lemma (32) can be given by mathematical induction on n . If $n = 0$ we get

$$\begin{pmatrix} \mathcal{H}W_2 \\ \mathcal{H}W_1 \\ \mathcal{H}W_0 \end{pmatrix} = \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^0 \begin{pmatrix} \mathcal{H}W_2 \\ \mathcal{H}W_1 \\ \mathcal{H}W_0 \end{pmatrix}$$

which is true. We assume that the identity given holds for $n = k$. Thus the following identity is true.

$$\begin{pmatrix} \mathcal{H}W_{k+2} \\ \mathcal{H}W_{k+1} \\ \mathcal{H}W_k \end{pmatrix} = \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} \mathcal{H}W_2 \\ \mathcal{H}W_1 \\ \mathcal{H}W_0 \end{pmatrix}$$

For $n = k + 1$, we get

$$\begin{aligned}
\begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^{k+1} \begin{pmatrix} \mathcal{H}W_2 \\ \mathcal{H}W_1 \\ \mathcal{H}W_0 \end{pmatrix} &= \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} \mathcal{H}W_2 \\ \mathcal{H}W_1 \\ \mathcal{H}W_0 \end{pmatrix} \\
&= \begin{pmatrix} 5 & -8 & 4 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} \mathcal{H}W_{k+2} \\ \mathcal{H}W_{k+1} \\ \mathcal{H}W_k \end{pmatrix} \\
&= \begin{pmatrix} 5\mathcal{H}W_{k+2} - 8\mathcal{H}W_{k+1} + 4\mathcal{H}W_k \\ \mathcal{H}W_{k+2} \\ \mathcal{H}W_{k+1} \end{pmatrix} \\
&= \begin{pmatrix} \mathcal{H}W_{k+3} \\ \mathcal{H}W_{k+2} \\ \mathcal{H}W_{k+1} \end{pmatrix}.
\end{aligned}$$

If we suppose that $n < 0$ the proof can be done similarly. Consequently, by mathematical induction on n , the proof is completed. \square

Note that

$$A^n = \begin{pmatrix} G_{n+1} & -8G_n + 4G_{n-1} & 4G_n \\ G_n & -8G_{n-1} + 4G_{n-2} & 4G_{n-1} \\ G_{n-1} & -8G_{n-2} + 4G_{n-3} & 4G_{n-2} \end{pmatrix}.$$

For the proof see [56].

THEOREM 33. *If we define the matrices $N_{\mathcal{H}W}$ and $E_{\mathcal{H}W}$ as follow,*

$$N_{\mathcal{H}W} = \begin{pmatrix} \mathcal{H}W_2 & \mathcal{H}W_1 & \mathcal{H}W_0 \\ \mathcal{H}W_1 & \mathcal{H}W_0 & \mathcal{H}W_{-1} \\ \mathcal{H}W_0 & \mathcal{H}W_{-1} & \mathcal{H}W_{-2} \end{pmatrix}, \quad E_{\mathcal{H}W} = \begin{pmatrix} \mathcal{H}W_{n+2} & \mathcal{H}W_{n+1} & \mathcal{H}W_n \\ \mathcal{H}W_{n+1} & \mathcal{H}W_n & \mathcal{H}W_{n-1} \\ \mathcal{H}W_n & \mathcal{H}W_{n-1} & \mathcal{H}W_{n-2} \end{pmatrix}.$$

then the following identity is true:

$$A^n N_{\mathcal{H}W} = E_{\mathcal{H}W}.$$

Proof. We can use the following identities for the proof.

$$\begin{aligned}
A^n N_{\mathcal{H}W} &= \begin{pmatrix} G_{n+1} & -8G_n + 4G_{n-1} & 4G_n \\ G_n & -8G_{n-1} + 4G_{n-2} & 4G_{n-1} \\ G_{n-1} & -8G_{n-2} + 4G_{n-3} & 4G_{n-2} \end{pmatrix} \begin{pmatrix} \mathcal{H}W_2 & \mathcal{H}W_1 & \mathcal{H}W_0 \\ \mathcal{H}W_1 & \mathcal{H}W_0 & \mathcal{H}W_{-1} \\ \mathcal{H}W_0 & \mathcal{H}W_{-1} & \mathcal{H}W_{-2} \end{pmatrix}, \\
&= \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{pmatrix}
\end{aligned}$$

where

$$\begin{aligned}
b_{11} &= \mathcal{HW}_2 G_{n+1} + \mathcal{HW}_1 (-8G_n + 4G_{n-1}) + \mathcal{HW}_0 4G_n, \\
b_{12} &= \mathcal{HW}_1 G_{n+1} + \mathcal{HW}_0 (-8G_n + 4G_{n-1}) + \mathcal{HW}_{-1} 4G_n, \\
b_{13} &= \mathcal{HW}_0 G_{n+1} + \mathcal{HW}_{-1} (-8G_n + 4G_{n-1}) + \mathcal{HW}_{-2} 4G_n, \\
b_{21} &= \mathcal{HW}_2 G_n + \mathcal{HW}_1 (-8G_n + 4G_{n-1}) + \mathcal{HW}_0 4G_{n-1}, \\
b_{22} &= \mathcal{HW}_1 G_n + \mathcal{HW}_0 (-8G_n + 4G_{n-1}) + \mathcal{HW}_{-1} 4G_{n-1}, \\
b_{23} &= \mathcal{HW}_0 G_n + \mathcal{HW}_{-1} (-8G_n + 4G_{n-1}) + \mathcal{HW}_{-2} 4G_{n-1}, \\
b_{31} &= \mathcal{HW}_2 G_{n-1} + \mathcal{HW}_1 (-8G_n + 4G_{n-1}) + \mathcal{HW}_0 4G_{n-2}, \\
b_{32} &= \mathcal{HW}_1 G_{n-1} + \mathcal{HW}_0 (-8G_n + 4G_{n-1}) + \mathcal{HW}_{-1} 4G_{n-2}, \\
b_{33} &= \mathcal{HW}_0 G_{n-1} + \mathcal{HW}_{-1} (-8G_n + 4G_{n-1}) + \mathcal{HW}_{-2} 4G_{n-2},
\end{aligned}$$

Using the Theorem (17) the proof is done. \square

From Theorem (33), we can write the following corollary.

COROLLARY 34. *We have the following identity.*

(a): *If we define $N_{\mathcal{HG}}$ and $E_{\mathcal{HG}}$ as follows,*

$$N_{\mathcal{HG}} = \begin{pmatrix} \mathcal{HG}_2 & \mathcal{HG}_1 & \mathcal{HG}_0 \\ \mathcal{HG}_1 & \mathcal{HG}_0 & \mathcal{HG}_{-1} \\ \mathcal{HG}_0 & \mathcal{HG}_{-1} & \mathcal{HG}_{-2} \end{pmatrix}, \quad E_{\mathcal{HG}} = \begin{pmatrix} \mathcal{HG}_{n+2} & \mathcal{HG}_{n+1} & \mathcal{HG}_n \\ \mathcal{HG}_{n+1} & \mathcal{HG}_n & \mathcal{HG}_{n-1} \\ \mathcal{HG}_n & \mathcal{HG}_{n-1} & \mathcal{HG}_{n-2} \end{pmatrix},$$

then we get

$$A^n N_{\mathcal{HG}} = E_{\mathcal{HG}}.$$

(b): *If we define $N_{\mathcal{HH}}$ and $E_{\mathcal{HH}}$ as follows,*

$$N_{\mathcal{HH}} = \begin{pmatrix} \mathcal{HH}_2 & \mathcal{HH}_1 & \mathcal{HH}_0 \\ \mathcal{HH}_1 & \mathcal{HH}_0 & \mathcal{HH}_{-1} \\ \mathcal{HH}_0 & \mathcal{HH}_{-1} & \mathcal{HH}_{-2} \end{pmatrix}, \quad E_{\mathcal{HH}} = \begin{pmatrix} \mathcal{HH}_{n+2} & \mathcal{HH}_{n+1} & \mathcal{HH}_n \\ \mathcal{HH}_{n+1} & \mathcal{HH}_n & \mathcal{HH}_{n-1} \\ \mathcal{HH}_n & \mathcal{HH}_{n-1} & \mathcal{HH}_{n-2} \end{pmatrix},$$

then we get

$$A^n N_{\mathcal{HH}} = E_{\mathcal{HH}}.$$

(c): *If we define $N_{\mathcal{HR}}$ and $E_{\mathcal{HR}}$ as follows,*

$$N_{\mathcal{HR}} = \begin{pmatrix} \mathcal{HR}_2 & \mathcal{HR}_1 & \mathcal{HR}_0 \\ \mathcal{HR}_1 & \mathcal{HR}_0 & \mathcal{HR}_{-1} \\ \mathcal{HR}_0 & \mathcal{HR}_{-1} & \mathcal{HR}_{-2} \end{pmatrix}, \quad E_{\mathcal{HR}} = \begin{pmatrix} \mathcal{HR}_{n+2} & \mathcal{HR}_{n+1} & \mathcal{HR}_n \\ \mathcal{HR}_{n+1} & \mathcal{HR}_n & \mathcal{HR}_{n-1} \\ \mathcal{HR}_n & \mathcal{HR}_{n-1} & \mathcal{HR}_{n-2} \end{pmatrix},$$

then we get

$$A^n N_{\mathcal{HR}} = E_{\mathcal{HR}}.$$

(d): If we define $N_{\mathcal{HC}}$ and $E_{\mathcal{HC}}$ as follows,

$$N_{\mathcal{HC}} = \begin{pmatrix} \mathcal{HC}_2 & \mathcal{HC}_1 & \mathcal{HC}_0 \\ \mathcal{HC}_1 & \mathcal{HC}_0 & \mathcal{HC}_{-1} \\ \mathcal{HC}_0 & \mathcal{HC}_{-1} & \mathcal{HC}_{-2} \end{pmatrix}, \quad E_{\mathcal{HC}} = \begin{pmatrix} \mathcal{HC}_{n+2} & \mathcal{HC}_{n+1} & \mathcal{HC}_n \\ \mathcal{HC}_{n+1} & \mathcal{HC}_n & \mathcal{HC}_{n-1} \\ \mathcal{HC}_n & \mathcal{HC}_{n-1} & \mathcal{HC}_{n-2} \end{pmatrix},$$

then we get

$$A^n N_{\mathcal{HC}} = E_{\mathcal{HC}}.$$

6. Conclusion

In the literature, there have been so many studies of the sequences of numbers and the sequences of numbers were widely used in many research areas, such as physics, engineering, architecture, nature and art. In this study we introduce hyperbolic generalized Woodall sequence and focused on four special cases such as hyperbolic modified Woodall numbers, hyperbolic modified Cullen numbers, hyperbolic Woodall numbers and hyperbolic Cullen numbers.

■ In section 1, we present some important information related to generalized Woodall numbers such as recurrence relation, Binet's formula and generating function. Moreover we give some information about hyperbolic numbers and some examples studied in the literature.

■ In section 2, we define hyperbolic generalized Woodall numbers and four special cases such as hyperbolic modified Woodall numbers, hyperbolic modified Cullen numbers, hyperbolic Woodall numbers and hyperbolic Cullen numbers. In addition, we introduce Binet's formula and generating function of hyperbolic generalized Woodall numbers and four special cases.

■ In section 3, we define some identities related to hyperbolic generalized Woodall sequence such as hyperbolic modified Woodall numbers, hyperbolic modified Cullen numbers, hyperbolic Woodall numbers and hyperbolic Cullen numbers. e.g Simpson's formula, Catalan's identity and Cassani's identity.

■ In section 4, we define linear sum formulas related to hyperbolic generalized Woodall sequence and four special cases hyperbolic modified Woodall numbers, hyperbolic modified Cullen numbers, hyperbolic Woodall numbers and hyperbolic Cullen numbers.

■ In section 5, we define matrix formulation to hyperbolic generalized Woodall sequence.

Linear recurrence relations (sequences) have many applications. Next, we list applications of sequences which are linear recurrence relations.

First, we present some applications of second order sequences.

- For the applications of Gaussian Fibonacci and Gaussian Lucas numbers to Pauli Fibonacci and Pauli Lucas quaternions, see [2].
- For the application of Pell Numbers to the solutions of three-dimensional difference equation systems, see [9].
- For the application of Jacobsthal numbers to special matrices, see [77].
- For the application of generalized k-order Fibonacci numbers to hybrid quaternions, see [27].

- For the applications of Fibonacci and Lucas numbers to Split Complex Bi-Periodic numbers, see [78].
- For the applications of generalized bivariate Fibonacci and Lucas polynomials to matrix polynomials, see [79].
- For the applications of generalized Fibonacci numbers to binomial sums, see [75].
- For the application of generalized Jacobsthal numbers to hyperbolic numbers, see [57].
- For the application of generalized Fibonacci numbers to dual hyperbolic numbers, see [58].
- For the application of Laplace transform and various matrix operations to the characteristic polynomial of the Fibonacci numbers, see [16].
- For the application of Generalized Fibonacci Matrices to Cryptography, see [43].
- For the application of higher order Jacobsthal numbers to quaternions, see [42].
- For the application of Fibonacci and Lucas Identities to Toeplitz-Hessenberg matrices, see [21].
- For the applications of Fibonacci numbers to lacunary statistical convergence, see [8].
- For the applications of Fibonacci numbers to lacunary statistical convergence in intuitionistic fuzzy normed linear spaces, see [33].
- For the applications of Fibonacci numbers to ideal convergence on intuitionistic fuzzy normed linear spaces, see [34].
- For the applications of k -Fibonacci and k -Lucas numbers to spinors, see [35].
- For the application of dual-generalized complex Fibonacci and Lucas numbers to Quaternions, see [73].
- For the application of special cases of Horadam numbers to Neutrosophic analysis see [23].
- For the application of Hyperbolic Fibonacci numbers to Quaternions, see [14].

We now present some applications of third order sequences.

- For the applications of third order Jacobsthal numbers and Tribonacci numbers to quaternions, see [11] and [10], respectively.
- For the application of Tribonacci numbers to special matrices, see [80].
- For the applications of Padovan numbers and Tribonacci numbers to coding theory, see [45] and [4], respectively.
- For the application of Pell-Padovan numbers to groups, see [15].
- For the application of adjusted Jacobsthal-Padovan numbers to the exact solutions of some difference equations, see [22].
- For the application of Gaussian Tribonacci numbers to various graphs, see [72].
- For the application of third-order Jacobsthal numbers to hyperbolic numbers, see [18].
- For the application of Narayan numbers to finite groups see [36].
- For the application of generalized Guglielmo numbers to Gaussian numbers, see [54].

- For the application of generalized Woodall numbers to Gaussian numbers, see [55].
- For the application of generalized third-order Jacobsthal sequence to binomial transform, see [59].
- For the application of generalized Generalized Padovan numbers to Binomial Transform, see [60].
- For the application of generalized Tribonacci numbers to Gaussian numbers, see [61].
- For the application of generalized Tribonacci numbers to Sedenions, see [62].
- For the application of Tribonacci and Tribonacci-Lucas numbers to matrices, see [63].
- For the application of generalized Tribonacci numbers to circulant matrix, see [64].
- For the application of Tribonacci and Tribonacci-Lucas numbers to hybrinomials, see [76].
- For the application of hyperbolic Leonardo and hyperbolic Francois numbers to quaternions, see [19].

Next, we now list some applications of fourth order sequences.

- For the application of Tetranacci and Tetranacci-Lucas numbers to quaternions, see [65].
- For the application of generalized Tetranacci numbers to Gaussian numbers, see [66].
- For the application of Tetranacci and Tetranacci-Lucas numbers to matrices, see [67].
- For the application of generalized Tetranacci numbers to binomial transform, see [68].

We now present some applications of fifth order sequences.

- For the application of Pentanacci numbers to matrices, see [46].
- For the application of generalized Pentanacci numbers to quaternions, see [49].
- For the application of generalized Pentanacci numbers to binomial transform, see [50].

We now present some applications of second order sequences of polynomials.

- For the application of generalized Fibonacci Polynomials to the summation formulas, see [70].
- For some applications of generalized Fibonacci Polynomials, see [71].

We now present some applications of third order sequences of polynomials.

- For some applications of generalized Tribonacci Polynomials, see [69].

References

- [1] Aydın, F. T., Hyperbolic Fibonacci Sequence, *Universal Journal of Mathematics and Applications*, 2 (2), 59-64, 2019.
- [2] Azak, A.Z., Pauli Gaussian Fibonacci and Pauli Gaussian Lucas Quaternions. *Mathematics*, 2022, 10, 4655. <https://doi.org/10.3390/math10244655>.
- [3] Baez, J., The octonions, *Bull. Amer. Math. Soc.* 39(2), 145-205, 2002.
- [4] Basu, M., Das, M., Tribonacci Matrices and a New Coding Theory, *Discrete Mathematics, Algorithms and Applications*, 6 (1), 1450008, (17 pages), 2014.
- [5] Berrizbeitia, P., Fernandes, J. G., . González, M. J, Luca, F., Huguet, V. J. M., On Cullen numbers which are Both Riesel and Sierpiński numbers, *Journal of Number Theory*, 132, 2836-2841, 2012.
- [6] Bilu, Y., Marques, D., Togbe, A., Generalized Cullen numbers in linear recurrence sequences, *Journal of Number Theory*, 202, 412-425, 2019.

- [7] Biss, D.K., Dugger, D., Isaksen, D.C., Large annihilators in Cayley-Dickson algebras, *Communication in Algebra*, 36 (2), 632-664, 2008.
- [8] Bilgin, N.G., Fibonacci Lacunary Statistical Convergence of Order γ in IFNLS, *International Journal of Advances in Applied Mathematics and Mechanics*, 8(4), 28-36, 2021.
- [9] Büyük, H., Taşkara, N., On The Solutions of Three-Dimensional Difference Equation Systems Via Pell Numbers, *European Journal of Science and Technology*, Special Issue 34, 433-440, 2022.
- [10] Cerda-Morales, G., On a Generalization of Tribonacci Quaternions, *Mediterranean Journal of Mathematics* 14:239, 1-12, 2017.
- [11] Cerda-Morales, G., Identities for Third Order Jacobsthal Quaternions, *Advances in Applied Clifford Algebras* 27(2), 1043–1053, 2017.
- [12] Cockle, J., On a New Imaginary in Algebra, *Philosophical magazine, London-Dublin-Edinburgh*, 3(34), 37-47, 1849.
- [13] Cunningham, A. J. C., Woodall, H. J., Factorisation of $Q = (2^q \mp q)$ and $(q \cdot 2^q \mp 1)$, *Messenger of Mathematics*, 47, 1–38, 1917.
- [14] Daşdemir, A., On Recursive Hyperbolic Fibonacci Quaternions, *Communications in Advanced Mathematical Sciences*, 4(4), 198-207, 2021. DOI:10.33434/cams.997824.
- [15] Deveci, Ö., Shannon, A.G., Pell–Padovan-Circulant Sequences and Their Applications, *Notes on Number Theory and Discrete Mathematics*, 23(3), 100–114, 2017.
- [16] Deveci, Ö., Shannon, A.G., On Recurrence Results From Matrix Transforms, *Notes on Number Theory and Discrete Mathematics*, 28(4), 589–592, 2022. DOI: 10.7546/nntdm.2022.28.4.589-592.
- [17] Dikmen, C. M., Hyperbolic Jacobsthal Numbers, *Asian Research Journal of Mathematics*, 15(4), 1-9, 2019.
- [18] Dikmen, C. M., Altınoy, M., On Third Order Hyperbolic Jacobsthal Numbers, *Konuralp Journal of Mathematics*, 10 (1), 118-126, 2022.
- [19] Dişkaya, O., Menken, H., Catarino, P. M. M. C., On the Hyperbolic Leonardo and Hyperbolic Francois Quaternions, *Journal of New Theory*, 42, 74-85, 2023.
- [20] Fjelstad, P., Gal, S.G., n-dimensional Hyperbolic Complex Numbers, *Advances in Applied Clifford Algebras*, 8(1), 47-68, 1998.
- [21] Goy, T., Shattuck, M., Fibonacci and Lucas Identities from Toeplitz-Hessenberg Matrices, *Appl. Appl. Math*, 14(2), 699–715, 2019.
- [22] Göcen, M., The Exact Solutions of Some Difference Equations Associated with Adjusted Jacobsthal-Padovan Numbers, *Kırklareli University Journal of Engineering and Science* 8(1), 1-14, 2022. DOI: 10.34186/klujes.1078836.
- [23] Gökbaşı, H., Topal, S., Smarandache, F., Neutrosophic Number Sequences: An Introductory Study, *International Journal of Neutrosophic Science (IJNS)*, 20(01), 27-48, 2023. <https://doi.org/10.54216/IJNS.200103>.
- [24] Grantham, J., Graves, H., The abc conjecture implies that only finitely many s-Cullen numbers are repunits, <http://arxiv.org/abs/2009.04052v3>, *MathNT*, 2021.
- [25] Gurses, N., Senturk, G. Y., Yuce, S., A Study on Dual-Generalized Complex and Hyperbolic-Generalized Complex Numbers, *Journal of Science*, 34(1), 180-194, 2021.
- [26] Guy, R., *Unsolved Problems in Number Theory* (2nd ed.), Springer-Verlag, New York, 1994.
- [27] Gül, K., Generalized k-Order Fibonacci Hybrid Quaternions, *Erzincan University Journal of Science and Technology*, 15(2), 670-683, 2022. DOI: 10.18185/erzifbed.1132164.
- [28] Hamilton, W.R., *Elements of Quaternions*, Chelsea Publishing Company, New York, 1969.
- [29] Hooley, C., *Applications of the sieve methods to the theory of numbers*, Cambridge University Press, Cambridge, 1976.
- [30] Imaeda, K., M. Imaeda, *Sedenions: algebra and analysis*, *Applied Mathematics and Computation*, 115, 77-88, 2000.
- [31] Kantor, I., Solodovnikov, A., *Hypercomplex Numbers*, Springer-Verlag, New York, 1989.

- [32] Keller, W., New Cullen primes, *Math. Comput.* 64, 1733-1741, 1995.
- [33] Kişi, Ö., Tuzcuoglu, I., Fibonacci Lacunary Statistical Convergence in Intuitionistic Fuzzy Normed Linear Spaces, *Journal of Progressive Research in Mathematics* 16(3), 3001-3007, 2020.
- [34] Kişi, Ö., Debnath, P., Fibonacci Ideal Convergence on Intuitionistic Fuzzy Normed Linear Spaces, *Fuzzy Information and Engineering*, 1-13, 2022. <https://doi.org/10.1080/16168658.2022.2160226>.
- [35] Kumari, M., Prasad, K., Frontczak, R., On the k -Fibonacci and k -Lucas Spinors, *Notes on Number Theory and Discrete Mathematics*, 29(2), 322-335, 2023. DOI: 10.7546/nntdm.2023.29.2.322-335.
- [36] Kuloğlu, B., Özkan, E., Shannon, A.G., The Narayana Sequence in Finite Groups, *Fibonacci Quarterly*. 60(5), 212-221, 2022.
- [37] Luca, F., Stanica, P., Cullen numbers in binary recurrent sequences, FT. Howard (ed.), *Applications of Fibonacci Numbers : Proceedings of the Tenth International Research Conference on Fibonacci Numbers and their Applications*, Kluwer Academic Publishers, 9, 167-175, 2004.
- [38] Marques, D., On Generalized Cullen and Woodall numbers that are also Fibonacci numbers, *Journal of Integer Sequences*, 17, Article 14.9.4, 2014.
- [39] Marques, D., Fibonacci s -Cullen and s -Woodall numbers, *Journal of Integer Sequences*, 18, Article 15.1.4, 2015.
- [40] Meher, N. K., Rout, S. S., Cullen numbers in sums of terms of recurrence sequence, <http://arxiv.org/abs/2010.10014v1>, *MathNT*, 2020.
- [41] Moreno, G., The zero divisors of the Cayley-Dickson algebras over the real numbers, *Bol. Soc. Mat. Mexicana* 3(4), 13-28, 1998.
- [42] Özkan, E., Uysal, M., On Quaternions with Higher Order Jacobsthal Numbers Components, *Gazi University Journal of Science*, 36(1), 336-347, 2023. DOI: 10.35378/gujs.1002454.
- [43] Prasad, K., Mahato, H., Cryptography Using Generalized Fibonacci Matrices with Affine-Hill Cipher, *Journal of Discrete Mathematical Sciences & Cryptography*, 25(8-A), 2341-2352, 2022. DOI : 10.1080/09720529.2020.1838744.
- [44] Richter, W.D., On Hyperbolic Complex Numbers, *Appl. Sci.*, 12, 5844, 2022.
- [45] Shtayat, J., Al-Kateeb, A., An Encoding-Decoding algorithm based on Padovan numbers, arXiv:1907.02007, 2019.
- [46] Sivakumar, B., James, V., A Notes on Matrix Sequence of Pentanacci Numbers and Pentanacci Cubes, *Communications in Mathematics and Applications*, 13(2), 603-611, 2022. DOI: 10.26713/cma.v13i2.1725.
- [47] Sloane, N.J.A., The on-line encyclopedia of integer sequences, <http://oeis.org/>.
- [48] Sobczyk, G., The Hyperbolic Number Plane, *The College Mathematics Journal*, 26(4), 268-280, 1995.
- [49] Soykan, Y., Özmen, N., Göcen, M., On Generalized Pentanacci Quaternions, *Tbilisi Mathematical Journal*, 13(4), 169-181, 2020.
- [50] Soykan, Y., Binomial Transform of the Generalized Pentanacci Sequence, *Asian Research Journal of Current Science*, 3(1), 209-231, 2021.
- [51] Soykan, Y., Tribonacci and Tribonacci-Lucas Sedenions. *Mathematics* 7(1), 74, 2019.
- [52] Y. Soykan, On Summing Formulas For Generalized Fibonacci and Gaussian Generalized Fibonacci Numbers, *Advances in Research*, 20(2), 1-15, 2019.
- [53] Soykan, Y., Generalized Woodall Numbers: An Investigation of Properties of Woodall and Cullen Numbers via Their Third Order Linear Recurrence Relations, *Universal Journal of Mathematics and Applications*, 5 : 2, 69-81, 2022.
- [54] Soykan, Y., Yılmaz, B., Gaussian Generalized Guglielmo Numbers, *Asian Journal of Advanced Research and Reports*, 17 (12), 1-18, 2023.
- [55] Soykan, Y., Eren, O., Gaussian Generalized Woodall Numbers, *Archives of Current Research International*, 23 (8), 48-68, 2023.
- [56] Soykan, Y., A Study On Generalized (r,s,t) -Numbers, *MathLAB Journal*, 7, 101-129, 2020.

- [57] Soykan, Y., Taşdemir, E., A study on hyperbolic numbers with generalized Jacobsthal numbers components, *Int. J. Nonlinear Anal. Appl.*, 2, 1965–1981, 2022.
- [58] Soykan, Y., On Dual Hyperbolic Generalized Fibonacci Numbers, *Indian J Pure Appl Math*, 2021. <https://doi.org/10.1007/s13226-021-00128-2>.
- [59] Soykan, Y., Taşdemir, E., Göcen, M., Binomial Transform of the Generalized Third-Order Jacobsthal Sequence, *Asian-European Journal of Mathematics*, 15(12), 2022. <https://doi.org/10.1142/S1793557122502242>.
- [60] Soykan, Y., Taşdemir, E., Okumuş, İ., A Study on Binomial Transform of the Generalized Padovan Sequence, *Journal of Science and Arts*, 22(1), 63-90, 2022. <https://doi.org/10.46939/J.Sci.Arts-22.1-a06>.
- [61] Soykan, Y., Taşdemir, E., Okumuş, İ., Göcen, M., Gaussian Generalized Tribonacci Numbers, *Journal of Progressive Research in Mathematics(JPRM)*, 14 (2), 2373-2387, 2018.
- [62] Soykan, Y., Okumuş, İ., Taşdemir, E., On Generalized Tribonacci Sedenions, *Sarajevo Journal of Mathematics*, 16(1), 103-122, 2020. ISSN 2233-1964, DOI: 10.5644/SJM.16.01.08.
- [63] Soykan, Y., Matrix Sequences of Tribonacci and Tribonacci-Lucas Numbers, *Communications in Mathematics and Applications*, 11(2), 281-295, 2020. DOI: 10.26713/cma.v11i2.1102 .
- [64] Soykan, Y., Explicit Euclidean Norm, Eigenvalues, Spectral Norm and Determinant of Circulant Matrix with the Generalized Tribonacci Numbers, *Earthline Journal of Mathematical Sciences*, 6(1), 131-151, 2021. <https://doi.org/10.34198/ejms.6121.131151>.
- [65] Soykan, Y., Tetranacci and Tetranacci-Lucas Quaternions, *Asian Research Journal of Mathematics*, 15(1): 1-24, 2019; Article no.ARJOM.50749.
- [66] Soykan, Y., Gaussian Generalized Tetranacci Numbers, *Journal of Advances in Mathematics and Computer Science*, 31(3): 1-21, Article no.JAMCS.48063, 2019.
- [67] Soykan, Y., Matrix Sequences of Tetranacci and Tetranacci-Lucas Numbers, *Int. J. Adv. Appl. Math. and Mech.* 7(2), 57-69, 2019, (ISSN: 2347-2529).
- [68] Soykan, Y., On Binomial Transform of the Generalized Tetranacci Sequence, *International Journal of Advances in Applied Mathematics and Mechanics*, 9(2), 8-27, 2021.
- [69] Soykan, Y., Generalized Tribonacci Polynomials, *Earthline Journal of Mathematical Sciences*, 13(1), 1-120, 2023. <https://doi.org/10.34198/ejms.13123.1120>.
- [70] Soykan, Y., A Study on Generalized Fibonacci Polynomials: Sum Formulas, *International Journal of Advances in Applied Mathematics and Mechanics*, 10(1), 39-118, 2022. (ISSN: 2347-2529).
- [71] Soykan, Y., On Generalized Fibonacci Polynomials: Horadam Polynomials, *Earthline Journal of Mathematical Sciences*, 11(1), 23-114, 2023. E-ISSN: 2581-8147. <https://doi.org/10.34198/ejms.11123.23114>.
- [72] Sunitha, K., Sheriba, M., Gaussian Tribonacci R-Graceful Labeling of Some Tree Related Graphs, *Ratio Mathematica*, 44, 188-196, 2022.
- [73] Şentürk, G.Y., Gürses, N., Yüce, S., Construction of Dual-Generalized Complex Fibonacci and Lucas Quaternions, *Carpathian Math. Publ.* 2022, 14 (2), 406-418, 2022. doi:10.15330/cmp.14.2.406-418.
- [74] Taş, S., On Hyperbolic Jacobsthal-Lucas Sequence, *Fundamental Journal of Mathematics and Applications*, 5 (1), 16-20, 2022.
- [75] Ulutaş, Y.T., Toy, D., Some Equalities and Binomial Sums about the Generalized Fibonacci Number u_n , *Notes on Number Theory and Discrete Mathematics*, 28(2), 252–260, 2022. DOI: 10.7546/nntdm.2022.28.2.252-260.
- [76] Taşyurdu, Y., Polat, Y.E., Tribonacci and Tribonacci-Lucas Hybrinomials, *Journal of Mathematics Research*; 13(5), 2021.
- [77] Vasanthi, S., Sivakumar, B., Jacobsthal Matrices and their Properties. *Indian Journal of Science and Technology* 15(5): 207-215, 2022, <https://doi.org/10.17485/IJST/v15i5.1948>.

- [78] Yılmaz, N., Split Complex Bi-Periodic Fibonacci and Lucas Numbers, *Commun.Fac.Sci.Univ.Ank.Ser. A1 Math. Stat.* 71(1), 153–164, 2022. DOI:10.31801/cfsuasmas.704435.
- [79] Yılmaz, N., The Generalized Bivariate Fibonacci and Lucas Matrix Polynomials, *Mathematica Montisnigri*, Vol LIII, 33-44, 2022. DOI: 10.20948/mathmontis-2022-53-5.
- [80] Yılmaz, N., Taskara, N., Tribonacci and Tribonacci-Lucas Numbers via the Determinants of Special Matrices, *Applied Mathematical Sciences*, 8(39), 1947-1955, 2014.