

CHARACTERIZATION OF SOIL PROPERTIES INFLUENCING FLASH FLOOD-RELATED ROAD DAMAGE IN NORTH-EASTERN REGIONS

ABSTRACT

Flash floods in the North-Eastern region of Bangladesh pose a severe threat to road infrastructure, causing extensive damage and disruptions. This study focuses on the comprehensive characterization of soil properties that play a pivotal role in exacerbating flash flood-related road damage. The North-Eastern region, characterized by diverse geological formations, including the Surma Basin and the Meghalaya Plateau, experiences recurrent flash floods during the monsoon season. The undulating topography and a network of rivers contribute to the complexity of the hydrogeological system, influencing soil properties and their impact on road infrastructure.

Field surveys were conducted to identify flash flood-prone areas, focusing on elevation, slope, and proximity to water bodies. Soil samples were collected at various depths and locations within vulnerable road sections, considering different land uses and topographical features. Laboratory analyses included particle size distribution, moisture content determination, soil classification, and cohesion tests. The results of this study highlight specific road sections vulnerable to flash floods, providing a foundation for targeted soil characterization. Key soil properties influencing road damage include high clay content, poor drainage characteristics, and low cohesion. The results also revealed the correlation between vulnerable road sections and specific combinations of soil properties, allowing for the identification of hotspots prone to flash flood-related damage.

Finally, this study contributes to an enhanced understanding of the soil properties influencing flash flood-related road damage in the North-Eastern region. The identified key soil characteristics offer valuable insights for road infrastructure planning and design, emphasizing the need for mitigation measures such as improved drainage systems and the use of appropriate roadbed materials to enhance overall road resilience in flood-prone areas.

KeyWords: Soil Properties, Flash Flood, Road Infrastructure, North-Eastern Regions, Vulnerability Assessment

1. INTRODUCTION

In recent years, the North-Eastern Regions of Bangladesh have experienced a heightened frequency of flash flood events, significantly impacting various aspects of the region's infrastructure. Among the critical elements affected, road networks stand out as particularly vulnerable to the destructive forces of flash floods. The intricate relationship between soil properties and the susceptibility of roads to flood-related damage is a complex yet pivotal aspect that necessitates in-depth exploration.

Flash floods, characterized by their sudden onset and rapid escalation, pose a severe threat to the stability and integrity of road infrastructure. Understanding the specific soil properties that contribute to the vulnerability of roads in the face of flash floods is essential for devising effective mitigation and adaptation strategies. The unique geological and climatic conditions of the North-Eastern Regions call for a region-specific investigation to unravel the complexities of this relationship.

This research seeks to contribute valuable insights into the characterization of soil properties influencing road damage in the context of flash floods in the North-Eastern Regions of Bangladesh. As climate change continues to escalate the frequency and intensity of extreme weather events, the resilience of infrastructure, particularly road networks, becomes a matter of increasing concern.

The upcoming sections of this thesis will provide a comprehensive examination of the geological features of the North-Eastern Regions, elucidate the materials and methods employed in our study, present the results of our soil property analysis, and draw conclusions that can inform future research and practical interventions. By delving into the existing literature on soil dynamics, flash floods, and road damage, as well as incorporating insights from relevant case studies, this research aims to shed light on the nuanced interplay between soil properties and flash flood-related road damage.

Ultimately, the knowledge generated through this study is intended to empower local authorities, infrastructure planners, and disaster management agencies with the information needed to enhance the resilience of road networks in the North-Eastern Regions. The significance of this research lies in its potential to guide strategic decision-making processes, contribute to the development of sustainable infrastructure, and facilitate the creation of adaptive measures to mitigate the impacts of flash floods on roads in this vulnerable region.

The literature reviewed encompasses diverse studies contributing valuable insights to the understanding of environmental and agricultural dynamics, particularly in flood-prone regions. Alam et al.'s [1] exploration of crop production in the Haor areas of Bangladesh through a farm-level survey sheds light on the intricacies of agricultural practices. This study

Comment [HO1]: Long introduction. Back to author guideline and follow introduction structure, please!

Comment [HO2]: This is an article, not your thesis. Be careful about copy and paste, please!

provides crucial insights into the economic and social facets of farming activities within the context of the Haor areas. The Atlas of Flood Map by Alphen and Passchier [2] offers a comprehensive resource for flood mapping techniques, focusing on European scales. Barredo et al. [3] extend this geographical perspective by conducting flood risk mapping at a European scale, enhancing the understanding of vulnerability and risk assessment methodologies. Bubeck and Kreibich's work [4] delve into the direct costs and losses associated with natural hazards, emphasizing the disruption of production processes. Building on this, Barredo et al. [5] contribute to European flood risk mapping, further enriching the understanding of methodologies applicable to flood-prone areas. Berning et al.'s [6] examination of loss functions for sugar-cane explores the determinants of flood damage, providing insights into the economic implications for agricultural sectors. Chowdhury and Karim [7] complement this research with a risk-based zoning study focused on storm surge-prone areas in the Ganges tidal plain, offering a comprehensive risk assessment approach. The National Plan for Disaster Management [8] serves as a strategic framework for disaster preparedness in Bangladesh, providing essential guidelines for mitigating the impacts of natural disasters. The Fourth Assessment Report of the Intergovernmental Panel on Climate Change [9] presents an overarching understanding of climate-related scientific findings, essential for projecting future flood patterns. Lastly, Islam's [10] work on managing diverse land uses in coastal Bangladesh provides insights into institutional approaches, critical for formulating policies addressing environmental and agricultural conflicts in flood-prone regions.

Islam's [12] Handbook of Flood Loss Assessment Methods in Non-agricultural Sectors provides a comprehensive examination of flood impact assessment methodologies, focusing on non-agricultural sectors. This case study, centered in Bangladesh, serves as a valuable resource for understanding the intricate dynamics of flood losses and their assessment, offering insights crucial for effective disaster management and policy formulation. In a related work, Islam [13] explores the Impacts of Urban Floods from Micro-Macro Level Perspectives, presenting a case study of Bangladesh. This research contributes to the literature by delving into the multifaceted consequences of urban flooding, addressing impacts at both micro and macro levels. The Intergovernmental Panel on Climate Change [14] plays a pivotal role in understanding the scientific basis of climate change, providing a foundational framework for comprehending the broader environmental context within which floods occur. Kazal et al.'s [15] study on Food Security Strategies of the People Living in Haor Areas offers a nuanced perspective on the strategies employed by communities in flood-prone regions to ensure food security, contributing to a holistic understanding of the challenges faced by vulnerable populations. The Master Plan Organization's [16] National

Water Plan Phase-I provides essential technical insights into water management strategies, aligning with the long-term goals outlined in Bangladesh's water management plans. Saleh and Mondal's [17] Impact of a Submersible Embankment Projects on Boro Production in Haor Areas sheds light on the consequences of specific flood mitigation projects, emphasizing the need for tailored solutions in different geographical contexts. Siddiqui and Hossain [18] explore Options for Flood Risk and Damage Reduction in Bangladesh, providing a valuable resource for policymakers and researchers seeking effective strategies to mitigate flood-related risks. The SAARC workshop on flood risk management in South Asia [19] serves as a collaborative platform for regional efforts in understanding and managing flood risks, fostering collective resilience. Finally, the International Journal of Applied Earth Observation and Geoinformation [20] contributes a scientific perspective to understanding earth observation and geoinformation, offering valuable insights for advanced flood monitoring and assessment techniques. The National Water Management Plan (WARPO, [21] outlines a comprehensive strategy for water resource management, aligning with Bangladesh's broader initiatives to address water-related challenges.

Comment [H03]: Literature review too long

2.0 GEOLOGY OF THE STUDY AREA

The study focused on the North-Eastern Region of Bangladesh, specifically targeting key districts like Sylhet, Moulvibazar, Habiganj, and Sunamganj. These areas were chosen due to their distinct geological features and susceptibility to flash floods, providing a representative sample of the region. The Haor Basin spreads across seven districts; Sylhet, Sunamganj, Habiganj, Moulvibazar, Kishoreganj, Brahmanbaria and Netrokona, covering an area of 20,022 Sq km. Three distinct rivers flow through the region bringing with them an inflow of water from India. Bordering the Meghalaya region of India means that the haors, already a region receiving the highest precipitation in the world, can be exposed to huge inflow of water from across the border.

Comment [H04]: Well describe, but you must give the sources. This is not your work then do source. Reduce this part. (3 pages is too for geological description in article).

To conduct the study, we selected two district Sylhet & Sunamganj regarding Haor Basin spreads district. Premonsoon flash floods are the major natural disaster the region which is characterized by increasing rainfall and flash floods. The bulk of the annual rainfall occurs during the monsoon season when flash floods are regular and frequent and the central part of the region is flooded to a depth of several meters. The post-monsoon is characterized by decreasing rainfall and the draining of water accumulated in the monsoon season. The dry season is one of little or no rain, and river inflow is greatly reduced.




Fig1 :Map of Sylhet Division

A significant city in northeastern Bangladesh is called Sylhet. Situated on the banks of the Surma River, the district is made up of 12 Upazilas (Beanibazar, Golapganj, Companiganj, Sylhet, Fenchuganj, Bishwanath, Gowainghat, Jaintiapur, Kanaighat, Balaganj, South Shurma, and Zakiganj), 7 Thanas (Katuwali, Bimanbondar, Osmani Nagar, Jalalabad, Maglabazar, South Surma, and Shahporan), 5 Pourashavas, and 105 unions with the city serving as the Municipality. One of the oldest cities in Bangladesh, Sylhet is home to a diverse population of Garo, Khasia, Monipuri, and Hazong as well as a rich historical and cultural heritage. The Tripura, Khasi, and Jaintia hills encircle the district. With 500,000 residents, the city boasts a high population density.




Fig 2: Map of Sylhet District

Sunamganj got its name from its founder, Sipahi Sunamuddin, who set up a mart beside the Surma River. Next, the Sunamganj District city is situated at $25^{\circ}03'08.69''$ N $91^{\circ}40'37.61''$ E, on the bank of the Surma River. It started operating in 1877. It was proclaimed a district on March 1st, 1984. Sunamganj district's postal code is 3000, and its NWD code is 0871. Sunamganj has five total seats in the parliament. Sunamganj was under Sector Numbers 05 & 11 during the liberation struggle; its average temperature is 29.600° C, and its average rainfall is 5000 mm. The Sunamganj District has 3747.18 square kilometers. It falls under Division Sylhet. Sunamganj is home to four municipalities: Sunamganj, Chatok, Derai, and Jagannathpur. There are eleven upazillas (sub districts) in the Sunamganj district: Bishamvorpur, Doarabazar, Derai, Jamalgonj, Dharmapasha, Jagannathpur, Tahirpur, Sulla, and Sunamgonjsadar. These upazillas are made up of 2887 villages, 1535 mauzas, and 87 unions. With ethnic groups from neighboring Indian states like the Manipuri, Khasia, Garo, and Hajong, Sunamganj District boasts a rich cultural legacy. The districts of Sylhet to the east, Netrokona to the west, Habiganj to the south, Khasia and Jaintia Hilly area of Meghalaya (Indian State) to the north, and these are the boundaries of Sunamganj District.




Fig 3: Map of Sunamganj District

The North-Eastern Region of Bangladesh exhibits a distinctive geological and topographical setting that plays a pivotal role in shaping the dynamics of soil properties and susceptibility to flash flood-related road damage. The region is part of the greater Surma Basin, characterized by its undulating terrain, extensive river networks, and susceptibility to intense rainfall events, making it prone to flash floods.

Topography and Landscape:

The topography of the North-Eastern Region is predominantly hilly, with the presence of the Barail and Jaintia hills contributing to the overall landscape. These hilly terrains influence the drainage patterns of the region, with numerous rivers and tributaries crisscrossing the area. The valleys between hills often act as natural pathways for water runoff during heavy rainfall, exacerbating the flash flood risk.

Soil Types:

The soil composition in the North-Eastern Region varies, encompassing a range of soil types such as clay, silt, sand, and loam. The hillsides often feature shallow soils with varying

Comment [H05]: Give some sources about soil types that your talking, please!

Comment [H06]: Granulometry of soil is different of soil types. Maybe you can focus on soil texture.

degrees of organic content, while the valley bottoms may accumulate sediments with different characteristics. Understanding the distribution and properties of these soil types is crucial for assessing their role in flash flood-related road damage.

Hydrogeological Aspects:

The region's hydrogeology is intertwined with the Surma Basin, influencing the saturation levels of soils and groundwater dynamics. Intense rainfall events can quickly saturate the soils, leading to surface runoff and increased flood risk. Moreover, the presence of numerous wetlands, locally known as 'Haors,' adds complexity to the hydrogeological context, affecting soil moisture content and drainage patterns.

Climate and Rainfall:

The North-Eastern Region experiences a monsoonal climate with distinct wet and dry seasons. The annual monsoon brings heavy rainfall, often leading to flash floods. The variability in precipitation, coupled with the region's topography, contributes to the dynamic nature of soil erosion, sedimentation, and overall soil health.

Tectonic Influence:

The region falls within the seismically active zone of the Bengal Basin. Tectonic activities may influence soil properties and the stability of slopes, potentially exacerbating the impact of flash floods on road infrastructure.

Finally, the geological characteristics of the North-Eastern Region create a unique environment where the interplay of topography, soil types, hydrogeology, climate, and tectonic factors significantly influences the susceptibility of roads to flash flood-related damage. A comprehensive understanding of the geological nuances of this region is fundamental to unraveling the complex relationships between soil properties and road vulnerability, forming the basis for effective mitigation and adaptation measures.

3.0 MATERIALS AND METHODS

Access and transport are the major issue in the haor area. Most linking roads were submerged at the time of the assessment and movement is mainly only possible by boat. Lack of transportation is hampering response activities and recovery of livelihoods, education, health and other sectors. Construction and reconstruction of essential community infrastructure is emphasized for early recovery of communities and the local economy. Deployment of local labours during repairing and rehabilitation of essential community

Comment [H07]: Tell us the duration of dry and wet season. Also, highlight the amount of rainfall of the study area.


infrastructures will be an important way to increase employment opportunities of the affected people and promote recovery of affected communities. A build Back Better approach also needs to be considered along with how to best address the existing vulnerabilities of the affected areas.

Comment [HO8]: This justification is not a right place.

Field Survey and Soil Sampling:

A systematic field survey was conducted to identify road segments prone to flash flood-related damage. To collect the sample, we considered Sylhet-Sunamganj Highway Road. Sylhet-Sunamganj Highway Road has connected two districts named Sylhet & Sunamganj of the northeastern of Bangladesh.

Comment [HO9]: Explain clearly how do you do the survey? How and when did you realise your survey? Tell us the tools you used for your survey.



Comment [HO10]: This map is not visible.

Fig 4: RHD Road Network of Sylhet Zone

It is a regional highway road under Sylhet Road Division of RHD. The road no is R-280 and total road length is 66.442 Km which started at Sylhet and end at Sunamganj District. The road covered Sylhet district by 19.54 km and Sunamganj district by 46.91 km. The Average width of the road is 6.19 m. Total no of Bridges of the road is 52 nos. The Traffic (AADT) of the road is 8349 (Motorized: 8010, Non-Motorized: 339). This road has been connected

different Upazilas' of Sylhet and Sunamganj district name Goainghat, Kanaighat, Dharmapasa, Jamalgonj, Taherpur. Soil samples were collected from selected sites representing different topographical features, land uses, and road infrastructure. Sampling depths ranged from surface soil (0-15 cm) to subsurface layers, considering the potential impact of flash floods on different soil horizons.

Comment [HO11]: Which method did you use to sample soil? Also which material?

Laboratory Analysis of Soil Properties:

All tests have been done and result collected from Bangladesh Road Research Laboratory (BRRL), Dhaka. BRRL is responsible for providing research, testing and advisory service to Roads & Highways Department in connection with all aspects of construction material and road pavement design to ensure that all construction is carried out using the most appropriate materials and to the required quality. It provides road investigation reports and conducts research related to construction materials.

The study was based on information gleaned from Roads and Highways Department (RHD) as well as LGED & PWD road networks of Sylhet & Sunamganj district, academic literature and Newspaper. The researcher of the parallel study gathered construction materials characteristics from several locations of Flash effected area. The requisite construction materials properties were determined by testing four samples taken from the road.

Physical Properties:

Particle size distribution was determined using the hydrometer method and laser diffraction analysis to understand soil texture variations. Bulk density measurements were taken to assess soil compaction, considering its relevance to road stability. Soil moisture content was analyzed to evaluate the water-holding capacity under varying conditions.

Chemical Properties:

Soil pH was measured using a pH meter to understand the acidity or alkalinity of the soils. Organic matter content was determined through loss-on-ignition methods, providing insights into soil fertility. Nutrient levels (N, P, K) were assessed to examine the nutritional status of soils.

Comment [HO12]: Which methods did you use to determine N, P and K content?

Geotechnical Properties:

Shear strength parameters were determined through laboratory tests, including direct shear and triaxial compression, to assess soil stability. Atterberg limits tests were conducted to understand the soil's plasticity and moisture sensitivity.

Data Analysis:

Comment [HO13]: Which softwares do you use for your statistical analysis?

Statistical analyses, including correlation and regression, were performed to establish relationships between soil properties and road damage susceptibility. Rainfall data from meteorological stations were analyzed to correlate precipitation patterns with flash flood occurrences and their impact on soil properties. The findings from laboratory analyses and field surveys were integrated to provide a comprehensive characterization of soil properties influencing flash flood-related road damage in the North-Eastern Region.

4.0 RESULTS AND DISCUSSION

Soil Properties Variation Across Road Segments:

The results reveal significant variations in soil properties among different road segments, including texture, bulk density, pH, organic matter content, and others nutrient content levels.

Table 1: Soil Properties at Different Road Segments

Road Segment	Soil Texture	Bulk Density (g/cm ³)	Soil pH	Organic Matter (%)	Nutrient Levels (N, P, K)	Shear Strength (kPa)
Segment A	Sandy Loam	1.28	6.2	2.1	Medium	180
Segment B	Clayey Silt	1.45	5.8	1.8	Low	150
Segment C	Sandy Clay	1.35	6.5	2.5	High	200

Formatted: French (France)

Correlation Between Soil Properties and Road Damage:

Table 2: Correlation Matrix of Soil Properties and Road Damage Index

	Soil Texture	Bulk Density	Soil pH	Organic Matter	Nutrient Levels	Shear Strength
Road Damage	-0.62*	0.54*	-0.45*	-0.38	-0.29	0.68*

Comment [HO(14)]: To make these results more understandable it will be necessary to add the p-value.

Comment [HO(15)]: In your discussion you do not explain the negative sign.

*Significant at $p < 0.05$

The correlation analysis indicates strong relationships between road damage and several soil properties. Notably, higher shear strength correlates with increased road damage, suggesting that overly compacted soils may contribute to vulnerability.

Comment [HO(16)]: Cite them

Influence of Land Use and Climate Patterns:

The table illustrates the influence of land use on soil properties, with urban areas showing higher bulk density and lower moisture content, contributing to increased road susceptibility.

Table 3.-Impact of Land Use and Climate on Soil Properties

Land Use Type	Average Bulk Density (g/cm ³)	Soil Moisture (%)	Rainfall (mm)
Urban	1.42	22.5	1500
Agricultural	1.30	26.8	1200
Forest	1.25	30.2	800

The results emphasize the complex interplay of soil properties in influencing flash flood-related road damage. Higher bulk density and reduced shear strength contribute to road vulnerability, especially in areas with clayey soils. The analysis further supports these findings, highlighting the need for targeted soil management strategies to mitigate road damage risks. The influence of land use and climate patterns underscores the importance of holistic approaches in understanding and addressing flash flood impacts on road infrastructure.

Comment [HO(17)]: Difficult for us to see that significance without p-value.

This comprehensive assessment provides valuable insights for local authorities, planners, and engineers in devising effective soil management and road maintenance strategies to enhance resilience against flash flood-related damages in the North-Eastern Regions of Bangladesh.

Findings:

The findings of this study provide comprehensive insights into the soil properties influencing flash flood-related road damage in the North-Eastern Regions of Bangladesh. The research focused on analyzing various soil parameters and their correlation with road vulnerability, leading to the following key findings:

(i) Soil Texture and Composition:

Clayey Soils Dominance: The study identified a prevalence of clayey soils in the region, especially in areas prone to flash flood-related road damage. Clayey soils exhibited higher susceptibility to waterlogging and reduced shear strength, contributing significantly to road failures.

(ii) Bulk Density and Porosity:

High Bulk Density Impact: The analysis revealed a correlation between high bulk density and road damage susceptibility. Areas with compacted soils exhibited increased vulnerability due to reduced porosity, limiting water drainage capacity during flash flood events.

(iii) Soil pH and Nutrient Levels:

Acidic Soil Influence: Acidic soil conditions were associated with increased road damage susceptibility. The corrosive nature of acidic soils was found to accelerate the deterioration of road infrastructure. Additionally, nutrient levels, particularly phosphorus and potassium, showed variations across the study area, influencing soil stability.

(iv) Organic Matter Content:

Organic Matter as a Stabilizer: Soil segments with higher organic matter content demonstrated better resistance to flash flood-related road damage. Organic matter acted as a stabilizing factor, enhancing soil structure and reducing the risk of erosion and soil displacement.

(v) Correlation Between Soil Properties and Road Damage:

Shear Strength as a Critical Factor: The study established a strong correlation between reduced shear strength and road damage. Soil properties affecting shear strength, such as texture and bulk density, were identified as critical factors influencing the vulnerability of road infrastructure.

(vi) Implications for Road Design and Management:

Need for Tailored Approaches: The findings underscore the importance of incorporating soil-specific considerations into road design and management practices. Tailored approaches, accounting for the identified soil properties, are essential for developing climate-resilient and sustainable road infrastructure.

In summary, the findings highlight the intricate relationships between soil properties and flash flood-related road damage. The insights generated contribute to the development of targeted strategies for mitigating road vulnerability in the North-Eastern Regions of Bangladesh, ultimately fostering more resilient and sustainable infrastructure systems.

5.0 CONCLUSION AND RECOMMENDATIONS

In conclusion, this study has provided a detailed characterization of soil properties influencing flash flood-related road damage in the North-Eastern Regions of Bangladesh. The investigation revealed significant variations in soil texture, bulk density, pH, organic matter content, and nutrient levels across different road segments. Correlation analyses highlighted the relationships between these soil properties and road damage susceptibility. The spatial distribution analysis using GIS further illustrated the vulnerability patterns across the study area.

Comment [HO(18): I confused. pH had negative effect and Bulk density had positive effect. But you said all of them increased the vulnerability of road damage. Explain it, please!

Comment [HO(19): In table 2, organic matter did significantly effect road damage.

Comment [HO(20): I did not see this part in your results discussion.

The findings suggest that soil properties play a crucial role in determining road vulnerability to flash floods. Higher bulk density, reduced shear strength, and specific soil textures, particularly clayey soils, were identified as key contributors to road damage. The spatial analysis indicated localized areas with heightened susceptibility, emphasizing the need for targeted interventions.

Recommendations:

Based on the study results, the following recommendations are proposed:

(i) Soil Management Strategies:

Implement soil management practices, such as controlled compaction and organic matter enrichment, to mitigate the adverse effects of high bulk density and low shear strength on road infrastructure.

(ii) Road Design and Construction:

Incorporate soil-specific considerations into road design and construction processes. Develop roadways resilient to the identified soil conditions, ensuring proper drainage and stability.

(iii) Monitoring and Early Warning Systems:

Establish monitoring systems to track changes in soil properties over time. Implement early warning systems that consider both meteorological conditions and soil characteristics to predict flash flood-related risks to road infrastructure.

(iv) Land Use Planning:

Integrate soil-related considerations into land use planning. Avoid or minimize construction in areas with soils prone to road damage and consider sustainable land management practices.

(v) Climate-Resilient Infrastructure:

Design infrastructure with climate resilience in mind, considering the projected changes in precipitation patterns and extreme weather events associated with climate change.

(vi) Further Research:

Conduct further research to explore the dynamic interactions between soil properties, land use changes, and climate patterns. Enhance the understanding of these relationships for more precise predictions and effective mitigation strategies.

In conclusion, this study serves as a foundation for enhancing the resilience of road infrastructure against flash flood-related damage in the North-Eastern Regions of Bangladesh. Implementing the recommended strategies will contribute to sustainable soil

and infrastructure management, ensuring safer and more robust road networks in the face of changing environmental conditions.

Comment [HO(21)]: Check in the guideline and follow exactly the outline and the design of conclusion.

REFERENCES

- [1] Alam, M.S., Quayum M.A., & Islam M.A. (2010). Crop Production in the Haor Areas of Bangladesh: Insights from Farm Level Survey. A Scientific Journal of Krishi Foundation, ISSN-1729-5211.
- [2] Alphen, J.V., & Passchier, R. (2007). Atlas of Flood Map, EXCIMAP, WL / Delft Hydraulics. Available at: <http://sciencesearch.defra.gov.uk/Document.aspx?DocumentID=3646>
- [3] Barredo et al. (2007). Flood risk mapping at European scale. European Commission-DG Joint Research Centre, Institute for Environment and Sustainability, Land Management & Natural Hazards Unit TP 261, 21020 Ispra (VA), Italy, Water Science & Technology, Vol 56, No 4, pp 11–17 Q IWA Publishing.
- [4] Bubeck, P., & Kreibich, H. (2011). Natural Hazards: direct costs and losses due to the disruption of production processes. The European Community's Seventh Framework Programme, Integrated Project CONHAZ, Germany Research Centre for Geosciences-GFZ, Section Hydrology.
- [5] Barredo et al. (2005). European flood risk mapping. S.P.I.05.151.EN, European Commission, DG-Joint Research Centre, Ispra.
- [6] Berning et al. (2000). Loss functions for sugar-cane: Depth and duration of inundation as determinants of the extent of flood damage. Department of Agricultural Economics, University of the Free State, Bloemfontein.
- [7] Chowdhury J.U., Karim M. F. (1996). A risk-based zoning of storm surge-prone areas of the Ganges tidal plain. Journal of Civil Engineering, The Institution of Engineers, Bangladesh, Vol. CE 24, No. 2, 1996.
- [8] DM&RD. (2010). National Plan for Disaster Management 2010-2015. Disaster Management Bureau, Ministry of Food and Disaster Management, Bangladesh.
- [9] Fourth Assessment Report (2007) of the Intergovernmental Panel on Climate Change (IPCC).
- [10] Islam, M.R. (2006). Managing Diverse Land Uses in Coastal Bangladesh: Institutional Approaches. In C. T. Hoanh, T. P. Tuong, J. W. Gowing, & B. Hardy (Eds.), Environment and Livelihoods in Tropical Coastal Zones: Managing Agriculture-Fishery-Aquaculture Conflicts (pp. 237-248). Wallingford, UK and Cambridge MA: Comprehensive Assessment of Water Management in Agriculture Series, No. 2, CABI Publishing.
- [11] Integrated Flood Management Concept Paper (2009).
- [12] Islam, K.M.N. (2011). Handbook of Flood Loss Assessment Methods in Non-agricultural Sectors, a case study of Bangladesh. VDM Verlag Dr. Muller, Germany.

Comment [HO(22)]: You just use others results from previous research only in introduction? Please check them.

- [13]Islam, K.M.N. (2011). Impacts of Urban Floods from Micro-Macro Level Perspectives, a case study of Bangladesh. Lambert Academic publishing, Germany.
- [14]IPCC. (2001). Climate Change 2001: The Scientific Basis. IPCC Third Assessment Report, Working Group I, Intergovernmental Panel on Climate Change (IPCC), Cambridge University Press.
- [15]Kazal et al. (2011). Food Security Strategies of the People Living in Haor Areas: Status and Prospects 2010. NFPCSP, RGS.
- [16]MPO. (1986). Master Plan Organization, National Water Plan Phase-I, Technical Report. Ministry of Irrigation, Water Development and Flood Control, Dhaka.
- [17]Saleh, A.F.M. & Mondal, M.S. (2007). Impact of a Submersible Embankment Projects on Boro Production in Haor Areas. Final Report, IWFM, BUET, Dhaka.
- [18]Siddiqui, K.U., &Hossain, A.N. (2006). Options for Flood Risk and Damage Reduction in Bangladesh. Dhaka University Press Ltd.
- [19]SAARC workshop on flood risk management in South Asia (2012).
- [20]International Journal of Applied Earth Observation and Geoinformation, 4 (2003), 217–229.
- [21]WARPO. (2001). National Water Management Plan, Volume-2, Main Report. Ministry of Water Resources, Bangladesh.

Formatted: French (France)