

# EDUCATIONAL VIDEO-ASSISTED CONSENT VERSUS TRADITIONAL INFORMED CONSENT FOR EXTRACORPOREAL SHOCKWAVE LITHOTRIPSY: A RANDOMISED CONTROLLED STUDY

---

## ABSTRACT

**Aims** :Traditionally, informed consent involves verbal and/or written material provided to the patient by a treating clinician. Multimedia interventions improve patients' knowledge and understanding during the informed consent process. This study aimed to compare pre-procedural anxiety levels and pain scores between educational video-assisted informed consent and traditional informed consent for extracorporeal shockwave lithotripsy (ESWL) at our centre.

**Study design** :This was a prospective, randomised, controlled study conducted at two centres.

**Place and duration of study** : The study was conducted in two Urology centre Department of Urology, Penang General Hospital and Department of Urology, Hospital Kuala Lumpur between 15<sup>th</sup> May 2022 till 15<sup>th</sup> October 2022 .

**Methodology** : The study group consisted of all adult patients undergoing ESWL in both centres. A video presentation explaining the ESWL procedure was developed in two languages, and group allocation was randomised using a computer-based random number generator. Anxiety levels were assessed using the Amsterdam Preoperative Anxiety and Information Scale (APAIS) questionnaire, visual analogue scale, and numerical rating scale used to collect data on pain scores.

### Results

A total of 54 respondents, with a predominance of male patients (57.4%) and the majority of patients having completed secondary school education (53.7%). In comparing the two groups, the video-assisted respondents exhibited significantly lower anxiety scores regarding the procedure than those in the traditional group ( $p < 0.05$ ). However, there were no statistically significant differences between the groups in terms of pain scores, both pre- and post-procedural.

### Conclusion

Video-assisted informed consent can reduce procedure-related patient anxiety levels before ESWL; however, there was no difference in pain scores between the two consent methods.  
patients.

*Keywords: Informed consent, video-assisted, ESWL, anxiety, pain score, randomized controlled trial*

## **1. INTRODUCTION**

Informed consent is essential for clinicians to provide comprehensive information regarding intended medical procedures. Informed consent describes the indications, risks, benefits, and alternatives of the procedure. It is important that patients have sufficient information about the procedure to allow them to make appropriate decisions regarding their treatment options.

Traditionally, the informed consent process involves verbal and/or written material provided to the patient by a treating clinician. However, this traditional approach can be time-consuming, increasing the likelihood of patient misunderstanding and inadequate comprehension of the procedure. Effectively communicating complex medical information to individuals who may not possess a deep understanding of the subject matter presents a significant challenge. Thus, there is a need to explore alternative methods to improve patient understanding of the informed consent process. Several methods have been explored to increase patient knowledge, including multimedia interventions, feedback modules, and extended discussions [1]. In a review by Nishimura et al. (2013), it was found that five of the sixteen total multimedia interventions (31%) resulted in a significant improvement in understanding of the informed consent process [1]. Chill, Dior, and Shveiky (2019) expressed a favourable clinical opinion regarding the use of multimedia during the informed consent process [2]. Tait and Voepel-Lewis (2015) also reported increasing effectiveness and acceptance of multimedia tools for research and healthcare consent [3]. Extracorporeal shockwave lithotripsy (ESWL) is a common procedure performed in our urology unit for the management of urolithiasis. Traditionally, oral and written informed consent is obtained prior to the procedure. However, elucidating how ESWL functions and its role in the management of urolithiasis can sometimes pose challenges. Patients are often unable to visualise how the procedure will be performed, potentially leading to incomplete information delivery to patients, which may further confuse them.

### **1.1 Anxiety**

A study conducted by Paton et al. (2018) demonstrated that the use of multimedia teaching tools significantly decreases anxiety levels [4]. Moerman et al. developed the Amsterdam Preoperative Anxiety and Information Scale (APAIS) in 1996, which comprises self-report questionnaires [5]. It includes six self-reported items that gauge anxiety and the need for information, rated on a five-point Likert scale. It was originally a Dutch version and has been validated in the Malay version by Mohd Fahmi et al. [6]. It is a simple and reliable instrument for assessing anxiety, widely utilised across the globe.

### **1.2 Pain score**

Cognitive impairment caused either by pain or opioids can interfere with the informed consent process [7]. In a study by Elke et al. (2018), the median pain score during ESWL was 5, despite the administration of analgesia [8]. Patients with

anxiety also experienced different levels of pain perception during the procedure. A study by Felice et al. (2020) suggested that effective anxiety management can reduce the need for analgesia during office procedures [9]. However, no study has compared the differences in pain scores between patients who underwent traditional informed consent and those who received educational video-assisted consent.

Currently, no studies have compared the use of educational video-assisted informed consent with traditional informed consent for ESWL. Our study aimed to investigate the impact of educational video-assisted informed consent on pre-procedural anxiety levels and pain scores. We hypothesised that educational video-assisted informed consent would reduce anxiety and result in lower pain scores

## **2. MATERIAL AND METHODS**

### **2.1 Aim**

This study aimed to assess pre-procedural anxiety levels between educational video-assisted informed consent and traditional informed consent for ESWL. Our secondary objective was to compare the pain scores between educational video-assisted informed consent and traditional informed consent for ESWL.

### **2.2 Study Type and Design**

This was a prospective, randomised, controlled study conducted in two tertiary healthcare centres. The study group included all adult patients undergoing ESWL between 15 May 2022 and 15 October 2022 in both centres.

A video presentation of the ESWL procedure was developed by the research team members and is available in both Malay and English languages. The video content included how ESWL was performed and what patients could expect during and after the procedure.

Group allocation was achieved through simple randomisation based on odd and even numbers generated using a computer-based random number generator.

#### **2.2.2 Anxiety level**

All participants provided informed consent and completed a validated questionnaire regarding their anxiety levels using the Amsterdam Preoperative Anxiety and Information Scale (APAIS). It includes six self-reported items assessing anxiety and the need for information, rated on a five-point Likert scale. Both the Amsterdam Preoperative Anxiety and Information Scale (APAIS) and validated Amsterdam Preoperative Anxiety and Information Scale (APAIS) in Malay were used to measure the anxiety levels.

#### **2.2.3 Pain score**

The visual analogue scale (VAS) and numeric rating scale (NRS) were used to assess the pain scores before and after the procedure. Participants were required to choose one of these pain measurement tools and self-report their pain on a data collection form.

## **2.3 Study Population**

All adult patients were scheduled for ESWL in the ESWL Suite at both centres during the study period.

### **2.4 Inclusion Criteria**

- a) Male and female individuals aged 18 and above undergoing ESWL
- b) Capable of reading and hearing either in the English language or Bahasa Malaysia
- c) No previous personal history of ESWL procedure

### **2.5 Exclusion Criteria**

- a) Participants with hearing impairments
- b) Participants with impaired vision
- c) Participants who refuse to provide consent to the study
- d) Participants below 18 years of age.

### **2.6 Withdrawal Criteria**

The participants had the option to withdraw from the study at any time. Subjects may be withdrawn if the investigator deems it detrimental or risky to continue their participation. Withdrawn participants will not be replaced.

### **2.7 Sample Size**

Based on the Miao et al. study, to achieve our objective of comparing the mean scores of the two groups, a sample size of 27 ( $n_1$ ) was required for the baseline/control group and a sample size of 27 ( $n_2$ ) for the other group in order to detect a mean difference of 2.00 with a power of 0.80 (80%) and a significance level ( $\alpha$ ) of 0.05 [6]. The mean difference was 2.00, which was considered the smallest important difference to detect. The standard deviation (SD) of the knowledge difference was estimated as 2.61 (Miao et al. study). This calculation was performed using ScalexMean version 1.0.2 (Naing, 2016).

## **2.8 Study Visits and Procedures**

During the study period, all patients who underwent ESWL pre-screening were randomised into two groups. All patients were seen on the same day of the procedure in the ESWL suite. After obtaining their consent to participate in the study, they were recruited into the study by a trained research team. Following an explanation of the study, the patients were randomised into two groups. A trained research team member explained the ESWL procedure in the traditional consent group. Patients in the intervention group watched an educational video after providing verbal and written consent. The video was available in either Malay or English, according to the patient's preference. After obtaining consent, both groups were given ample time to ask questions. Patients were provided with a form to self-

report their anxiety levels after signing their written consent. Patients underwent ESWL as per the standard hospital protocol with a standard dose of pain medication according to the hospital protocol. Pain scores post-procedural were recorded using the same data collection form.

### 3. RESULTS AND DISCUSSION

Data analysis was performed using IBM SPSS Statistics for Windows version 22. Descriptive statistics were employed for selected variables. The results were presented as frequencies and percentages for categorical data. Chi-square, Fisher's exact, and paired t-tests were used for statistical analysis. A value of  $P < 0.05$  is considered significant.

A total of 54 respondents were included in the study. Demographic data indicated that 31 participants (57.4%) were male, with a mean age of 51.2 years in the traditional consent and 52.2 years in the video-assisted consent group (Table 1). The majority of participants had an educational background in secondary school (53.7%) or a diploma (22.2%). Based on each anxiety question score, procedure-based questions were significant in the video-assisted consent group compared to the traditional consent group (Table 2). However, there was no significant difference in the anaesthesia-related questions between the two groups. Additionally, there was no statistical difference between the pre-and post-procedural pain scores in both groups. (Table 3).

Table 1: Descriptive data of the respondents.

Demographic data		Traditional		Video	
		Count	%	Count	%
Gender	Female	8	29.6%	15	55.6%
	Male	19	70.4%	12	44.4%
Education level	Primary	5	18.5%	0	0.0%
	Secondary	12	44.4%	17	63.0%
	Diploma	7	25.9%	5	18.5%
	Degree	3	11.1%	4	14.8%
	Master	0	0.0%	1	3.7%
Location of the stone	Distal ureteric stone	6	22.2%	1	3.7%
	Mid ureteric stone	3	11.1%	9	33.3%
	Proximal ureteric stone	3	11.1%	7	25.9%

	Renal stone	15	55.6%	10	37.0%
--	-------------	----	-------	----	-------

Table 2: Statistical analyses of anxiety level scores between two study groups.

APAIS		Traditional		Video		P value
Question 1	Not at all	4	14.8%	8	30.8%	0.093
	Somewhat anxious	7	25.9%	12	46.2%	
	Moderately anxious	10	37.0%	5	19.2%	
	Highly anxious	5	18.5%	1	3.8%	
	Extremely anxious	1	3.7%	0	0.0%	
Question 2	Not at all	3	11.5%	8	29.6%	0.055
	Somewhat anxious	7	26.9%	13	48.1%	
	Moderately anxious	12	46.2%	4	14.8%	
	Highly anxious	3	11.5%	2	7.4%	
	Extremely anxious	1	3.8%	0	0.0%	
Question 3	Not at all	1	3.7%	9	33.3%	0.026
	Somewhat anxious	8	29.6%	10	37.0%	
	Moderately anxious	11	40.7%	6	22.2%	
	Highly anxious	6	22.2%	2	7.4%	
	Extremely anxious	1	3.7%	0	0.0%	
Question 4	Not at all	4	14.8%	11	40.7%	0.012
	Somewhat anxious	3	11.1%	9	33.3%	
	Moderately anxious	13	48.1%	5	18.5%	
	Highly anxious	6	22.2%	2	7.4%	
	Extremely anxious	1	3.7%	0	0.0%	
Question 5	Not at all	2	7.4%	12	46.2%	0.001

	Somewhat anxious	5	18.5%	9	34.6%	
	Moderately anxious	11	40.7%	5	19.2%	
	Highly anxious	8	29.6%	0	0.0%	
	Extremely anxious	1	3.7%	0	0.0%	
Q6Possible	Not at all	1	3.7%	12	44.4%	0.000
	Somewhat anxious	4	14.8%	9	33.3%	
	Moderately anxious	9	33.3%	5	18.5%	
	Highly anxious	10	37.0%	1	3.7%	
	Extremely anxious	3	11.1%	0	0.0%	

Refer to appendix.  $P < 0.05$  is significant.

**Table 3: Mean pre- and post-procedural pain score between both study groups.**

Mean pain score	Traditional	Video	P value
Pre-procedural pain score	2.3	1.2	0.026
Post-procedural pain score	3.7	3.2	0.458

Informed consent is an important aspect of urological procedures, ensuring that patients possess a thorough understanding of the procedures, associated risks, benefits, and available alternatives before consenting to undergo urological procedures. Traditional consent relies on both verbal and written information. However, the integration of assisted technologies, such as videos, has been shown to reduce anxiety levels during various medical procedures. A study conducted by Gouda et al. (2022) among women undergoing colposcopy showed that video-assisted teaching reduced anxiety levels and alleviated pain experiences among these patients [10]. Similarly, research conducted by Shome et al. (2021) using video consent for blepharoplasty showed a decrease in preoperative anxiety levels [11]. Nevertheless, only a few studies have focused on video-assisted consent for urological procedures.

In our study, we used the Amsterdam Preoperative Anxiety and Information Scale, which is subdivided into two major components. Items 1 and 2 assessed anaesthesia-related anxiety, and items 4 and 5 evaluated surgery-related anxiety. The need for information was assessed using items 3 and 6. In our study, the video-assisted group exhibited lower anxiety scores than the traditional consent group in terms of surgery-related and need-for-information components. However, there was

no significant difference in anaesthesia-related anxiety between the two groups. This could be because the patient perceived the procedure as potentially painful, thus anticipating the need for anaesthesia or analgesia during the procedure, leading to a lower anaesthesia-related anxiety level.

Notably, our study did not reveal a significant difference in the pre-and post-procedural pain scores between the groups. We postulate that most of our patients were on analgesia prior to the procedure, as pain is one of the indications for active stone treatment at our centre. Various analgesic options were available for pre-procedural use at both centres. Additionally, it is conceivable that probably fewer shocks or energy were delivered during the procedure, and no data were collected if ESWL was completed with the intended delivered shockwave.

A limitation of the study is the absence of an assessment of participants' comprehension of the video content. In the era of internet and web browser access, patients may have conducted their own research on their medical condition and management options before the ESWL procedure. In the future, this study can be extended to explore the utilisation of internet resources, such as Google and social media, to evaluate patient comprehension, anxiety levels, and pain scores in ESWL or other urological-related procedures.

#### **4. CONCLUSION**

Video-assisted informed consent can reduce procedure-related anxiety in patients undergoing ESWL for stone management. Furthermore, integrating video-assisted consent into the process of prior urological procedures is a promising avenue for enhancing patient understanding and reducing anxiety.

#### **CONSENT (WHEREEVER APPLICABLE)**

All authors declare that written informed consent was obtained from the patient to participate in this study. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal.

#### **ETHICAL APPROVAL (WHEREEVER APPLICABLE)**

This study is registered with Malaysian National Medical Research Registry, Research ID NMRR-21-1677-60782. Medical research and Ethics committee approval has been obtained.

#### **REFERENCES**

- [1] Nishimura A, Carey J, Erwin PJ, Tilburt JC, Murad MH, McCormick JB (2013) Improving understanding in the research informed consent process: a systematic review of 54 interventions tested in randomized control trials. *BMC Med Ethics* 14:28. <https://doi.org/10.1186/1472-6939-14-28>.

[2] Chill HH, Dior U, Shveiky D (2019) Use of multimedia during informed consent: novelty or necessity. *Int Urogynecol J* 30:1635–7. <https://doi.org/10.1007/s00192-019-04046-0>. [Epub 2019 July 13]. PMID: [31302718](https://pubmed.ncbi.nlm.nih.gov/31302718/).

[3] Tait AR, Voepel-Lewis T (2015) Digital multimedia: a new approach for informed consent? *JAMA* 313:463–4. <https://doi.org/10.1001/jama.2014.17122>, PMID: [25647199](https://pubmed.ncbi.nlm.nih.gov/25647199/).

[4] Paton EA, Davis SK, Gaylord N, Cao X, Gosain A (2018) Impact of a multimedia teaching tool on parental anxiety and knowledge during the informed consent process. *PediatrSurg Int* 34:1345–52. <https://doi.org/10.1007/s00383-018-4352-z>.

[5] Moerman N, van Dam FS, Muller MJ, Oosting H (1996) The Amsterdam Preoperative Anxiety and Information Scale (APAIS). *AnesthAnalg* 82:445–51. <https://doi.org/10.1097/00000539-199603000-00002>.

[6] Mohd Fahmi Z, Lai LL, Loh PS (2015) Validation of the Malay version of the Amsterdam Preoperative Anxiety and Information Scale (APAIS). *Med J Malaysia* 70:243–8. PMID: [26358022](https://pubmed.ncbi.nlm.nih.gov/26358022/).

[7] Aja B. The effects of pain on informed consent; 2017.

[8] Bovelander E, Weltings S, Rad M, van Kampen P, Pelger RCM, Roshani H (2019) The influence of pain on the outcome of extracorporeal shockwave lithotripsy. *Curr Urol* 12:81–7. <https://doi.org/10.1159/000489424>.

[9] Sorrentino F, Petito A, Angioni S, D'Antonio F, Severo M, Solazzo MC et al (2021) Impact of anxiety levels on the perception of pain in patients undergoing office hysteroscopy. *Arch GynecolObstet* 303:999–1007. <https://doi.org/10.1007/s00404-020-05885-9>.

[10] Gouda ADK, Ismail LM, Hamdan-Mansour AM, Aboushady RMN (2022) Effect of video assisted-teaching on level of knowledge, anxiety and pain among women undergoing colposcopy. *Int J Health Sci* 9376–92. <https://doi.org/10.53730/ijhs.v6nS5.11975>.

[11] Shome D, Doshi K, Vadera S, Kumar V, Vyavahare SS, Kapoor R (2021) Personalized video consent in blepharoplasty: a new paradigm in the preoperative consent giving process. *J Cosmet Dermatol* 20:2211–23. <https://doi.org/10.1111/jocd.14172>.

## APPENDIX


### 1. Anxiety level / Tahapkegelisahan

Please circle the most appropriate feeling for each questions/ Sila bulatperasaanandauntuksetiapsoalan.

- 1 – Not at all / Tiada rasa gelisah
- 2 - Somewhat / Sedikit rasa gelisah
- 3 – Moderate / Sederhana rasa gelisah
- 4 – Moderately high / Sederhanatinggi rasa gelisah
- 5 – Extremely / Sangat gelisah

<b>Questions / Soalan</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>
I am worried about the anesthetic <i>Saya bimbangtentangpembiusansaya</i>	1	2	3	4	5
The anesthetic is on my mind continually <i>Saya bimbangtentangpembiusansaya</i>	1	2	3	4	5
I would like to know as much as possible about the anesthetic <i>Saya ingintahusebanyak yang mungkintentangpembiusansaya</i>	1	2	3	4	5
I am worried about the procedure Saya bimbangtentangprosedurpembedahansaya <i>Saya bimbangtentangprosedurpembedahansaya</i>	1	2	3	4	5
The procedure is on my mind continually <i>Prosedurpembedahansentiasaberada di fikiransaya</i>	1	2	3	4	5
I would like to know as much as possible about the procedure <i>Saya ingintahusebanyak yang mungkintentangpembedahansaya</i>	1	2	3	4	5

## PAIN SCORE



UNDER PEER