

Original Research Article

“Impact of priming with UV radiation on seed germination and seedling growth of Chakhao rice cultivar”

ABSTRACT

~~Development~~ The development of eco-friendly technology with innovative ideas is always essential to enhance plant growth and improve yield in a sustainable manner. Seed priming methods such as conventional and physical priming are ~~well~~ well-known pre-sowing techniques that are considered to be effective in improving the growth and yield of a crop. In view of this, an experiment was conducted to study the impact of physical priming with non-ionizing radiation like UV-C on seed germination and early seedling growth of Chakhao cultivars of rice (*Oryza sativa*). Three Chakhao rice cultivars namely Black (Amubi) - V₁, Red (Angangbi) - V₂ and White (Angoubi) - V₃ were used, and seeds were initially exposed to UV-C (8.32 K/J) radiation for 3 hours. Then seeds were soaked in water at 28±2^oC for 2 days and the ~~emerged~~ emerging seedlings were again treated with UV-C (8.32 K/J) for different durations i.e. 30 mins (T₃), 45 mins (T₄) and 60 mins (T₅). Seeds without any UV-C treatment (T₁) and only seed treatment for three hours (T₂) were taken as control. The data on germination rate, speed of germination, root length, shoot length, and seedling vigour index were recorded as per standard procedures. The results showed that all three rice cultivars recorded maximum germination percentage, speed of germination, shoot length, root length, and seedling vigor index under seed priming with UV- C for 60 mins followed by 45 mins. The results also indicated that control seeds recorded minimum germination and seedling growth.

Formatted: Font: Italic

Formatted: Not Superscript/ Subscript

Comment [j1]: Please add the values and state the conclusion of this study

Keywords: Chakhao Rice, UV-C, SVI, Germination rate, Seedling treatment and Seed priming

1. INTRODUCTION

Rice (*Oryzasativa*) is a major cereal crop that is consumed by more than half of the world's population as a primary staple food. It contributes to 21% of the world's per capita calorie intake, with over 90% of rice being produced and consumed in Asia. Rice can be categorized into pigmented and non-pigmented varieties, with "specialty rice" having unique aromas, flavours, and colors. In Manipur, India, a unique type of rice called Chakhao is consumed for its unique health benefits and flavour as well as high anthocyanin content. Chakhao rice has the potential to be used as a functional food and food colorant. ~~However, But~~ Chakhao rice is less popular among farmers due to its inherent undesirable traits such as low yield, high photo-insensitivity, longer vegetative phase, and tall stature which causes its lodging. Another constraint of Chakhao rice cultivation is its long harvest time ~~and~~ which ~~must need~~ needs to be compensated either by introducing early maturity traits s or by increasing productivity. Further, with the introduction of many ~~high-high~~-yielding rice varieties, farmers' preferences have also been changed, and as a result, many Chakhao landraces are getting degraded [13]. Thus, more research needs to be done ~~in-on~~ Chakhao rice to strengthen sustainability by preserving the local Chakhao rice of Manipur as well as improving desirable traits in the best rice varieties.

Low-level UV seed priming ~~has been found to be is~~ an environmentally safe technique for increasing crop production and yield as well as for inducing resistance to a variety of stresses in agricultural plants[3,5]. Ultra Violet radiation priming, a novel non-ionizing physical priming method proven to be beneficial on seed germination, seedling development, plant growth, and yield contribution [4]. UV priming has been shown to have a variety of physiological effects, including changes in growth, enhancement of photosynthetic pigments, taste or quality of the fruit, increased photo-protection, upregulation of the plant defence systems to strengthen crop production, and increased production of secondary metabolites [2,3,6,17].

Formatted: Indent: First line: 0.24"

Comment [j2]: Please provide examples of UV priming utilization on others species

Comment [j3]: Please state the objectives and potential benefits of this research

2. MATERIAL AND METHODOLOGY

The experiment was conducted in the RKVY Laboratory of Uttar BangaKrishiViswavidyalaya, Cooch Behar, West Bengal (26°19'86" N, 89°23'53" E) with an altitude of (43 m), India. The seeds were obtained ~~in-from~~ the local farmers with germination between 70% and 85%. The seeds were kept in a lineage bag in an ambient chamber, at temperature and relative humidity (27 ± 2 °C and $45 \pm 2\%$ respectively) until the beginning of the experiment.

To perform the experiment, a completely randomised design in a 3 × 5 factorial scheme with three replications was used. The treatments consisted of three

Chakhaor Rice cultivars i.e. Black rice, Red Rice, and White Rice, submitted to one UV-C radiation dose of 8.32 KJm⁻², corresponding to 0 hours i.e. control and 3 hours seedling priming, 3 hours seedling priming with 30 minutes seedling treatments, 3 hours seedling priming with 45 minutes seedling treatments and 3 hours seedling priming with 60 minutes seedling treatments. For each treatment, the seeds were exposed to UV-C radiation in a single batch.

The treatment with UV-C radiation was performed using a cylindrical irradiation chamber (46 cm in diameter, 90 cm in length), which has two lamps capable of emitting waves with a length of 254 nm, which corresponds to the range of UV-C radiation.

The seeds were placed on a petri disc for each radiation at a distance of 20 cm from the source of radiation and were treated accordingly to the purpose of the experiment, separately.

The seeds were arranged during the periods of time, so that they are/were as scattered as possible, in order to avoid that some seeds are on the other and not be struck by UV-C light.

In order to transform the exposure periods of the seeds into UV-C radiation at doses of UV-C (kJ m⁻²), we used the Lux Meter for measuring the light intensity, and with the conversion factors we converted it into kJ m⁻². As the seeds were arranged in the centre of the chamber, under the influence of the two lamps, they received a dose equivalent to 2.370 W m⁻² (0.0237 kJs⁻¹m⁻²).

The radiation intensity was kept constant at 2.37 W/m² for UV-C. The applied doses were obtained by varying the exposure time to the set distance, using the equation of López-Rubira et al. (2007):

$$D = \frac{(F \times T)}{1000}$$

Based on the exposure times and the dose emitted by the lamps, the equivalent doses were obtained at each time, by the product between these factors, resulting in the following doses: 8.32KJ/m².

After exposure of the seeds to different doses of UV-C radiation, they were subjected to physiological quality and later to sanitary quality tests.

a. Determination of Germination Percentage

The exposed seeds i.e. 3 hours of radiation treatment (approximately 10-12% moisture content) were kept in a desiccator. The seed germination study was done by using sterilized corning Petri dishes of 9 cm diameter. Each petri dish was layered with blotting paper and distilled water is/was used to moistening the blotting paper (Labouriau & Valadares 1976). The seeds were placed after 3 hours of imbibitions @ 50 seeds per Petri dish for germination. All Petri dishes were kept in the incubator

Comment [j4]: in the discussion you use different treatments (T₁ to T₅) and varieties (V₁, V₂, V₃), please indicate the T and V treatments accordingly

Comment [j5]: Please use a consistent citation format

maintained at 27± 2°C temperature. Seeds with radical length equal to or greater than 2 mm were considered as germinated. Germination percentage was calculated on the total no germinated seeds per treatment after three days of incubation. The experiment was done in three replicates and repeated for six months.

2.2 Speed of germination

The speed of germination was estimated according to the following formula given by Haugland and Brandsaeter (1996).

$$\bar{T} = \frac{\sum_{i=1}^k N_i}{\sum_{i=1}^k T_i}$$

Where T_i is the time from the start of the experiment to the i^{th} observation, N_i is the number of seeds germinated in the i^{th} time and k is the last time of observation.

2.3 Determination of seedling growth

To determine seedling growth, root length, and shoot length of different parts of seedlings were measured. Early seedling growth parameters were evaluated on 10-day-old seedlings. On the day of final measurements, seedlings were removed from the petri dish carefully and the length of shoot and root were measured separately.

2.3.1 Seedling vigor index

Based on the mean seedling length and mean seedling dry weight, the seedling vigor index (SVI) was calculated as per Abdul Baki and Anderson (1973) Using the following formulae;

SVI = Mean seedling length (cm) x Germination (%)

The vigor index (VI) of the seedlings was estimated as per the formula given by Abdul- Baki, and Anderson (1973):

SVI = (RL + SL) x GP

Where, RL - Root Length (cm) SL - Shoot Length (cm)

GP - Germination Percentage

2.3.2 Seedling length (cm)

For determining seedling length, five normal seedlings from each treatment were randomly selected for measuring the seedling length on the 12th day of germination. From the collar region to the tip of the root, the seedling length was measured. The average seedling length of five seedlings was computed and expressed in centimeters.

2.3.3 Root length (cm)

Five normal seedlings in each treatment were randomly selected from the germination test for measuring the root length on the 12th day of germination. From the collar region to the tip of the root, the root length was measured. The average root length of five seedlings was computed and expressed in centimeters.

2.3.4 Shoot length (cm)

The earlier five seedlings used for the root length measurement were also used for measuring shoot length. The shoot length was measured from the collar region to the point of

Comment [j6]: Please use a consistent citation format

Comment [j7]: Please use a consistent citation format

attachment of cotyledons. The average of five seedlings was computed and expressed in centimeters.

2.4 Statistical analysis

The data were submitted to ANOVA and the means of the qualitative treatments, when significant, were compared by the Tukey test ($p \leq 0.05$). The quantitative treatments were submitted to regression analysis ($p \leq 0.05$). The values of the parameters of the models were estimated as a function of the exposing duration of UV-C radiation. The degree of fit of the models to the experimental data was based on the magnitude and significance of [the](#) coefficient of determination, [the](#) significance of regression coefficients, [the](#) magnitude of coefficient of variation, and on verification of the fit of the statistical model to the experimental data.

3. RESULTS AND DISCUSSION

The present investigation was carried out on three different [C](#)ehakhaor Rice cultivars collected from different parts of [the](#) Manipur. The three cultivars includes Black rice (chakhaoamubi), Red rice (chakhaoangnangbi), and White rice (Chakhaoangoubi). The provided data appears to represent the mean values for various parameters across different treatments (T_1 to T_5) and varieties (V_1 , V_2 , V_3), along with analysis of variance at 0.05 level.

As depicted in [Figure 1](#), the Mean Germination Percentage varied between [63.655](#) and 73.630. Among all treatments, T_5 exhibited the highest Mean Germination Percentage for all varieties, with values ranging from 71.07 to 73.63 and 69.71. On the other hand, T_2 displayed the lowest Mean Germination Percentage (63.655) for V_1 . For V_2 and V_3 , the lowest germination percentage was observed at T_1 , with values of 66.85 and 67.46, respectively. Notably, significant differences were observed in all three factors, namely treatment, variety, and their interaction (Table 1). ~~And~~ according to Guajardo-Flores *et al.*[7], reactive oxygen species (ROS), which are beneficial during seed germination, are present at higher concentrations in plant tissues subjected to UV-C exposure. In line with this, Rupiasih and Vidyasagar[14] found that exposing wheat seedlings to UV-C for up to 180 minutes can enhance seed germination. Moreover, Neelamegam and Sutha[12] discovered that after 10 days of germination, groundnut seeds (*Arachis hypogaea* L.) exposed to UV-C showed a decline in germination rates, followed by an increase at 5 and 60 minutes, respectively.

As shown in [fig-Fig 1](#). The mean Speed of Germination (SOG) spans from [8.73](#) to [9.53](#). Specifically, the mean value of V_1 demonstrates [the](#) highest at T_2 ([9.53](#)) and attains its minimum at T_1 (8.93). In the case of V_2 , T_5 manifests the highest mean SOG (9.40),

Comment [j8]: Did you use statistical software? if yes, please state clearly including the manufacturer

Comment [j9]: Please consider to present the result and discussion for each parameter you observed in a sub-title

Comment [j10]: Please refer to the Journal format Figure 1 or Fig 1. ? please be consistent

Comment [j11]: 63.655% ?

Comment [j12]: in days or weeks?


Comment [j13]: please add the unit of each value

while T₂ records the lowest (9.06). Concerning V₃, T₃ and T₅ display the highest mean SOG (9.20), whereas T₁ exhibits the lowest (8.80). Significant values were observed for both the treatment and the interaction between treatment and variety; however, no significance was noted for variety alone (table 1).

Table: 1: Anova table of Impact of priming with UV radiation on germination rate, speed of germination, shoot length, root length seedling length, and sSeed vigor index.


Source of Variation	DF	Germination %	Speed of germination	Shoot Length	Root Length	Seedling length	SVI
		Mean Squares					
V	2	23.66**	0.032	0.763	0.543	0.211	14,071.57
Treatment	4	33.683**	0.167*	11.067**	9.722**	40.88**	413492.18**
VXT	8	8.054*	0.165**	2.307*	1.908*	5.04**	29330.056*
Error	30	3.338	0.044	0.996	0.841	1.569	11,442.24
Total	44						

**,* Significant at 1% and 5% levels of probability, respectively




Comment [j14]: The bars are difficult to see please add the vertical axis titles and the units

Fig 1: Impact of priming with UV radiation on Speed of germination and Germination Rate



Comment [j15]: please add the vertical axis titles and units

Fig 2: Impact of priming with UV radiation on shoot length, root length, and seedling length.



Comment [j16]: please add the vertical axis titles and units

Fig 3: Impact of priming with UV radiation on seedling vigor index.

The findings of Neelamegan and Shuta [12] corroborate the present study's observations, as both report that priming or treatment of seeds with UV radiation

Formatted: Indent: First line: 0.29"

results in the breakdown of radicles or molecules within the seed embryo, initiating metabolic processes at the earliest possible time. The supplied radiation energy triggers the breakdown of metabolic processes, which, when combined with water, leads to an increase in free radicals in the seeds, altering their ability to germinate. Kondrateva *et al.* [9] also observed the effects of UV radiation on seed germination, reporting an increase in the average length of sprouts and roots in seeds of naked oats by 11-21%.

According to Figure 2, the mean seedling length for V₁ and V₃ varieties ranged from 11.37 to 18.87 and 11.37 to 13.40, respectively whereas for V₂ it ranged from 14.57 to 17.60. T₅ showed the highest mean seedling length for both varieties V₁ and V₃ (18.87 and 19.67), similarly while T₁ had the lowest (11.37 and 13.40). For the V₂, T₄ had the longest seedling length (17.60), and T₂ had the shortest (14.57). In terms of shoot length, the mean values ranged from 6.333 to 10.700. T₅ has the maximum shoot length (9.83 and 10.70) for variety V₁ and V₃ respectively but for V₂, T₄ has the maximum (8.43) similarly T₁ had the lowest (6.33) for V₁, T₁ and T₂ both has the lowest (8.17) for V₂. Lastly, for V₃, T₂ has the lowest (6.87).

For root length, the mean values ranged from 5.033 to 9.033, and T₅ had the highest mean root length (9.03 and 8.97) for variety V₁ and V₃ respectively, while T₁ had the lowest (5.033) for V₁, T₂ (6.40) for V₂ and T₁ (6.53) for V₃. Significant values were observed in the treatment and its interaction with variety, but no significance was found for variety in any of the cases (Table 1) for seedling length, root length, and shoot length.

According to Neelamegam and Shuta [12], UV-C treatment displayed favorable effects on the seedling growth rate, and it was observed that UV-C irradiation for up to 60 minutes enhanced the growth parameters and biomass production of groundnut plants. Similarly, Scott *et al.* [15] demonstrated that UV-C treatments on tomato seeds resulted in a reduction in disease incidence and severity without adversely affecting germination or seedling growth. Additionally, Siddiqui *et al.* [16] reported that UV-C exposure led to an increase in shoot weight, shoot length, root length, root weight, leaf area, and the number of nodules in groundnut plants.

Figure 3 displays the mean values for the Seedling Vigor Index (SVI) ranging from 944.70 to 1729.13. Among the treatments, T₅ exhibited the highest mean SVI values for V₁, V₂, and V₃ (1688.50, 1570.23, and 1729.13), while T₁ displayed the lowest mean SVI for V₁, V₂, and V₃ (944.70, 1257.60, and 1138.33), respectively. The treatment and its interaction with variety were found to be significant, whereas no significant difference was observed for the variety alone. Previous research by

Comment [j17]: Please add the unit for each value so that easier for the reader to understand

Formatted: Indent: First line: 0.29"

Rupiasih and Vidyasagar[14]. has reported that exposure to UV-C radiation can yield positive results in terms of the germination and early seedling growth of groundnut plants.

Each treatment (T₁ to T₅) shows variations in mean values across different parameters. T₅ consistently shows higher mean values for speed of germination, shoot length, root length, SVI, and germination percentage. T₁ consistently shows lower mean values for these parameters.

4. Conclusion

The results suggest that Chakhao rice cultivars, namely black, red, and white, exhibited positive effects on seedling growth and development when exposed to UV-C radiation. According to the data, it is recommended that priming with UV radiation at 8.32 K/J for 3 hours before soaking and an additional 60 minutes of treatment after seedling emergence can significantly improve seedling growth and development, comparable to the control and other treatments. Each treatment (T1-T5) displayed variations in mean values across different parameters. T5 consistently exhibited higher mean values for germination speed, shoot length, root length, SVI, and germination percentage, while T1 showed lower mean values for these parameters.

Comment [j18]: significantly improved yet comparable ?

Comment [j19]: what is the T5 ?

Reference

1. Abdul-Baki AA, Anderson JD. Vigor determination in soybean seed by multiple criteria 1. *Crop sci.* 1973; 13(6): 630-3.
2. Aboul Fotouh MM, Moawad FG, El-Naggar HA, Tag El-din MA, Sharaf Eldeen HA. Influence of seed treatment with UV-C on saline stress tolerance in green beans (*Phaseolus vulgaris* L.). *J BiolChem Environ Sci.* 2014; 9(2): 391–414.
3. Badridze G, Kacharava N, Chkhubianishvili E, Rapava L, Kikvidze M, Chanishvili Sh, Shakarishvili N, Mazanishvili L, Chigladze L. Effect of UV radiation and artificial acid rain on ~~pre~~-productivity of wheat. *Russ J Ecol.* 2016; 47: 158–166.
4. Bera K, Dutta P, Sadhukhan S. Seed priming with non-ionizing physical agents: Plant responses and underlying physiological mechanisms. *Plant Cell Reports*, 2022; 41(1), 53-73.
5. Delibaltova V, Ivanova R. Impact of the pre-sowing irradiation of seeds by helium-neon laser on the dynamics of development of some cotton varieties. *J Environ Protection Ecol.* 2006; 7(4): 909–917
6. Dwivedi R, Singh VP, Kumar J, Prasad SM. Differential physiological and biochemical responses of two *Vigna* species under enhanced UV-B radiation. *J Radiat Res Appl Sci.* 2015; 8(2): 173–181.

7. Guajardo-Flores D, Serna-Guerrero D, Serna-Saldívar SO, JacoboVelázquez DA. Effect of germination and UV-C radiation on the accumulation of flavonoids and saponins in black bean seed coats. *Cereal Chem.*2014; 91:276–279
8. Haugland, E, Brandsaeter LO. Experiments on bioassay sensitivity in the study of allelopathy. *Journal of Chemical Ecology.* 1996;22:1845–1859.
9. Kondrateva NP, Kasatkina NI, Kuryleva AG, Baturina KA, Ilyasov IR, Korepanov RI. Effect of treatment of seeds of grain crops by ultraviolet radiation before sowing. In *IOP Conference Series: Earth and Environmental Science.* 2020; 433(1): 012039.
10. Labouriau LG, Valadares MB On the physiology of seed of *Calotropisprocera*. *Anais da Academia Brasileira de Ciência.*1976; 42(2): 235-64.
11. López RV, Artés Hernández FDA, Artés CF. Evaluación de la calidad de granadastratadas con UV-C y almacenadas en atmósferacontrolada; 2007.
12. Neelamegam R, Sutha T. UV-C irradiation effect on seed germination, seedling growth and productivity of groundnut (*Arachis hypogaea* L.). *Int J CurrMicrobiol App Sci.* 2015; 4:430–443
13. Roy S, Banerjee A, Pattanayak A., Roy SS, Rathi RS, Misra AK, Ngachan SV, Bansal KC. Chakhao (delicious) rice landraces (*Oryza sativa* L.) of North-east India: collection, conservation and characterization of genetic diversity. *Plant Genetic Res: Characterization and Utilization.* 2014; 12(3): 264 –272.
14. Rupiasih NN, Vidyasagar PB. Effect of UV-C radiation and hypergravity on germination, growth, and content of chlorophyll of wheat seedlings. In: *AIP conference proceedings* 1719:030035. AIP Publishing; 2016.
15. Scott G, Scott G, Almasrahi A, Mansoorkhani FM, Rugar M, DickinsonM, Shama G. Hormetic UV-C seed treatments for the control of tomato diseases. *Plant Pathology.* 2019; 68:700–707.
16. Siddiqui A, Dawar S, Zaki MJ, Hamid N. Role of ultraviolet (UV-c) radiation in the control of root infecting fungi on groundnut and mung bean; 2011.
17. Wargent JJ, Jordan BR. From ozone depletion to agriculture: understanding the role of UV radiation in sustainable crop production. *New Phytol.* 2013; 197(4): 1058–1076.

Formatted: Font: French (France)

Formatted: French (France)