

Effect of Drought Stress at Growth and Development of Pea(*Pisum sativum* L.)

Abstract

Peas belongs to a family Leguminosae which is cultivated as an edible seed all over the world and having the high nutritional importance. They are an effective source of fiber, vitamins, minerals, and plant-based protein. Legumes are essential for crop rotation because they fix nitrogen in the soil, enhancing soil fertility and lowering the demand for artificial fertilizers. Around 9,000 years ago, pea cultivation is said to have started in the Near East and the Mediterranean region. By altering numerous physiological and biochemical processes, including photosynthesis, water uptake, and nutrient assimilation, drought stress can drastically lower the productivity of pea. Lack of water can cause pea plants to grow slowly, have fewer leaves, and produce less photosynthesis. Additionally, it might lead to a decline in pea quality and production. Pea plants have a number of defense mechanisms to deal with drought stress, such as altered root systems, osmotic adjustment, the synthesis of antioxidants, and stomatal modulation. The activation of genes that control the synthesis of osmo-protectants, including proline and sugars, as well as the expression of stress-responsive proteins, like LEA proteins and chaperones, is part of the molecular defense system of pea against drought. [please mentioned the zest of your paper in brief](#)

Formatted: Indent: First line: 0.5"

Comment [W1]: remove the general statement

Comment [W2]:

Keywords: ~~Leguminosae~~, Drought, Photosynthesis, Osmotic Adjustment, ~~Chaperons~~, Antioxidant.

Introduction

The growth and development of plants can be influenced by a variety of environmental conditions, including soil quality, temperature, light, water, and nutrients which are climate variables that can have significant effects on plant physiology and define the kinds of plants that can flourish in a specific area. Especially, water deficiency and flooding cause the huge reduction in the productivity of crop (Habib-ur-Rahman et al., 2022). Natural disasters including drought may seriously hamper the agricultural growth and productivity, which in turn can have an impact on food security. Different types of crops may respond differently to drought depending on its intensity and length, but in general, drought stress can result in decreased photosynthesis, water and nutrient intake, and plant development. Plant development and survival may be hampered by drought stress or waterlogging caused by insufficient or excessive precipitation, respectively (Begna, 2020). A long period of particularly little rainfall defines a drought as a

Formatted: Indent: First line: 0.5"

natural calamity, which causes a lack of water supplies. Crop failure, livestock deaths, and a lack of water are all possible consequences for agriculture, wildlife, and human populations. Also, droughts can worsen other environmental issues like wildfires, soil erosion, and desertification. Droughts are predicted to become more frequent and severe in many parts of the world due to climate change, creating a serious worldwide challenge that calls for proactive management and adaptation solutions (Kopecká et al., 2023). The introduction part need to well structured, it is submitted in immature state. Lacking of key research highlights in the concerned title of the paper.

Comment [W3]: The introduction part need to well structured, it is submitted in immature state. Lacking of key research highlights in the concerned title of the paper

Comment [W4]:

Importance of Pea

Pea is a leguminous crop that is frequently consumed and has a number of health and environmental advantages. Pea cultivation has been practiced for a very long time; pea domestication artifacts have been discovered at ancient Egyptian, Greek, and Roman archaeological sites (Kumari and Deka, 2021). During the Middle Ages, peas were widely grown in Europe, and European settlers transported them to the Americas. Peas are an effective source of nutrients such as vitamins, nutritional fibers and minerals. They are a good source of vitamins B, C, and K. Iron, magnesium, and potassium are among the crucial minerals that are present in peas. They are also an excellent source of plant-based protein (Melchior et al., 2020). Peas have a number of health advantages. The high fiber content aids in controlling digestion and might aid in weight management. Flavonoids and carotenoids, two types of antioxidants found in peas, help lower the chance of developing chronic diseases like heart disease and some types of cancer. Peas are also anti-allergic, which benefits general health (Ma et al., 2018).

Peas are adaptable and can be used in a variety of delicious meals and can be used as dried, frozen, or fresh. Peas are often served as a side dish, in soups, stews, salads, and stir-fries. To prepare spreads or dips, they can also be crushed or pureed (Lu et al., 2020). Peas have a long tradition of production and consumption across numerous cultures. They are a common ingredient in many cuisines and are frequently used in traditional foods and celebrations (Mudryj et al., 2014; Martens et al., 2017). Because they can fix nitrogen in the soil, peas are said to be environmentally benign. Fixing nitrogen increases soil fertility and lessens the demand for

~~artificial-chemical~~fertilizers. Peas are a sustainable option because they require less water to grow than other crops(Garneau et al., 2018).

Comment [W5]: where is material and methods

Where is result and discussion part of the paper

Response of Plant under Osmotic Stress

Drought has adverse effect at the growth and development of peas. Water must be available continuously for peas to grow and develop properly. The development and productivity of pea crops can be significantly impacted when there is a water shortage as a result of a dry soil during drought conditions can prevent plant roots from absorbing as much nutrients(Bagheri et al., 2023). This may result in a decline in the plant's intrinsic health and vigor, stunted development, and decreased yields. Additionally, pea plants may become more vulnerable to pests and diseases as a result of drought stress. The amount of humidity has an impact on how quickly plants lose water through transpiration. Low humidity can cause an increase in water loss, whereas high humidity can reduce it. Some plants are adapted to specific amounts of humidity and may have difficulty surviving in areas with high humidity levels(Etienne et al., 2018).

Pea germination and production can be significantly influenced by drought stress. Pea seeds need enough moisture to germinate, and protracted droughts might stop or postpone this process. In addition, drought stress may result in poorer seedling development, lower plant biomass, and reduced yields(OKÇU et al., 2005). Through a variety of physiological pathways, drought stress can dramatically impair pea germination. One of the main reasons is the decrease in water availability, which can result in an increase in the pressure of water, which can harm the cell membrane and cause dehydration(Martin et al., 2006). As a result, the seed may not absorb water at the same pace, which is crucial for germination. Inhibiting the enzymes that degrade the seed's stored food stores is another way. Amylase, a key enzyme in turning stored starch into sugars that give energy for germination, can become less active under drought stress. This may cause partial or delayed germination(Li et al., 2013).

Moreover, the control of hormones in the seed, especially abscisic acid (ABA), might be impacted by drought stress. A hormone called ABA is essential for controlling seed germination

and dormancy. Increased ABA levels under drought stress conditions cause an increase in seed dormancy and a decrease in germination rate (Ali et al., 2020). The physiological and biochemical mechanisms of pea plants are susceptible to drought stress. For instance, drought stress can result in higher amounts of reactive oxygen species (ROS) and lower levels of antioxidants, which can harm plant cells through oxidative processes and adversely affect growth and yield (Carvalho, 2008; You and Chan, 2015).

The amount of chlorophyll in pea leaves can be significantly impacted by drought. The green color of plant is due to the presence of Chlorophyll and is essential to photosynthesis, the process through which plants turn sunlight into energy. Plants endure water stress during a drought, which may result in less water being available in the soil. As a result, plants reduce the intake of carbon dioxide (CO₂) required for photosynthesis by closing their stomata (tiny pores on the surface of the leaf). Due to this restriction, there is less CO₂ accessible for the photosynthetic process, which results in less glucose and other organic molecules being produced (Sarani et al., 2014; Hu et al., 2023).

Plants may see a drop in chlorophyll concentration along with decreased photosynthesis. The molecules of chlorophyll are vulnerable to environmental stresses, such as drought. The breakdown of chlorophyll molecules or the reduced production of new chlorophyll can both contribute to the decrease in chlorophyll content (Krasteva et al., 2013).

Defensive mechanism under Drought stress

Drought stress can have a substantial effect on the process of photosynthesis. Reduced water availability for plants during drought circumstances impairs their capacity to absorb moisture through the roots and transfer it to the leaves. In order to prevent excessive water loss, the stomata on the surface of the leaf close, which reduces carbon dioxide (CO₂) uptake and restricts photosynthesis (Yang et al., 2021). Pea plants can close their stomata to limit water loss, produce osmo-protectants to maintain cellular turgor, and activate stress-responsive genes to control physiological processes as some of their defenses against drought. Pea plants have evolved a number of chemical defenses against drought stress. The activation of genes that are responsive to stress and control the synthesis of proteins involved in stress tolerance is one such mechanism.

These proteins support the preservation of photosynthetic machinery, cellular homeostasis, and defense against oxidative damage(Lahuta et al., 2022).

Aside from that, drought stress causes the synthesis of signaling molecules like abscisic acid (ABA), which is essential for controlling how plants react to water constraint. Stomatal closure, water conservation, and decreased transpiration rates are all benefits of ABA. Furthermore, it regulates gene expression to encourage the production of defense-related proteins and enzymes that increase drought resistance in plants(Sah et al., 2016). Additionally, pea plants activate their antioxidant defense mechanisms to combat the rise in reactive oxygen species (ROS) brought on by drought stress. Chloroplasts, which are involved in photosynthesis, are among the biological components that ROS can harm. Superoxide dismutase (SOD) and catalase (CAT), two antioxidant enzymes, aid in scavenging ROS and shielding the photosynthetic machinery (Farooq et al., 2021; Pandey et al., 2023).

Generally, multiple molecular mechanisms, including gene regulation, ABA signaling, and activation of antioxidant defense systems, are involved in the response of pea plants to drought stress (Bharath et al., 2021). All of these systems work together to maintain plant survival during drought circumstances and to lessen the harmful impacts of water scarcity on photosynthesis. One of the strategies that helps the plant survive in a water-scarce environment is the activation of stress-responsive genes. These genes are in charge of making the proteins that shield the plant from harm brought on by water stress.

The DREB1 gene is one of the most significant stress-responsive genes in pea plants. This gene creates a protein that aids in controlling how the plant reacts to drought stress. The DREB1 protein activates additional genes that aid the plant in water conservation and cell protection when it is subjected to drought conditions(Nayak et al., 2009). another mechanism is LEA gene, a crucial stress-responsive gene in pea plants. This gene results in the production of a protein that aids in preventing dehydration-related cell damage in plants. By stabilizing other proteins and preventing their unfolding and destruction, the LEA protein serves as a molecular chaperone(Porcel et al., 2005).

[Where is conclusion part](#)

Wher is conflict of interest is mentioned

Reference

1. Ali, S., K. Hayat, A. Iqbal and L. Xie. 2020. Implications of Abscisic Acid in the Drought Stress Tolerance of Plants. *Agronomy*. 10: 1-28.
2. Bagheri, M., C. S. Santos, D. Rubiales and M. W. Vasconcelos. 2023. Challenges in pea breeding for tolerance to drought: Status and prospects. *Ann Appl Bio.*,183: 105-195.
3. Begna, T. 2020. Effects of Drought Stress on Crop Production and Productivity. *International Journal of Research Studies in Agricultural Sciences*. 6: 34-43.
4. Bharath, P., S. Gahir and A. S. Raghavendra. 2021. Abscisic acid-induced stomatal closure: An important component of plant defense against abiotic and biotic stress. *Front. Plant Sci.*, 12: 13-24.
5. Carvalho, M. H. C. 2008. Drought stress and reactive oxygen species. *Plant Signaling & Behavior*, 3: 156-165.
6. Etienne, P., S. Diquelou, M. Prudent, C. Salon, A. Maillard and A. Ourry. 2018. Macro and Micronutrient Storage in Plants and Their Remobilization When Facing Scarcity: The Case of Drought. *Agriculture*. 8: 1-17.
7. Farooq, M., R. Ahmad, M. Shahzad, Y. Sajjad, A. Hassan, M. M. Shah, S. Naz and S. A. Khan. 2021. Differential variations in total flavonoid content and antioxidant enzymes activities in pea under different salt and drought stresses. *ScientiaHorticulturae*, 287: 1-8.
8. Garneau, M. G., Q. Tan and M. Tegeder. 2018. Function of pea amino acid permease AAP6 in nodule nitrogen metabolism and export, and plant nutrition. *J. Exp. Bot.*, 69: 5205-5219.
9. Habib-ur-Rahman, M., A. Ahmad, A. Raza, M. U. Hasnain, H. F. Alharby, Y. M. Alzahrani, A. A. Bamagoos, K. R. Hakeem, S. Ahmad, W. Nasim, S. Ali, F. Mansour and A. EL Sabagh. 2022. Impact of climate change on agricultural production; Issues, challenges, and opportunities in Asia. *Front. Plant Sci.*, 13: 1-22.
10. Hu, F., Y. Zhang and J. Guo. 2023. Effects of drought stress on photosynthetic physiological characteristics, leaf microstructure, and related gene expression of yellow horn. *Plant Signal Behav.*, 18: 1-10.
11. Kopecká, R., M. Kameniarová, M. Černý, B. Brzobohatý and J. Novák. 2023. Abiotic Stress in Crop Production. *Int. J. Mol. Sci.*, 24: 1-47.
12. Krasteva, V., V. Alexandrov, M. Chepishcheva, S. Dambov, D. Stefanov, I. Yordanov and V. Goltsev. 2013. Drought induced damages of photosynthesis in bean and plantain

plants analyzed in vivo by chlorophyll a fluorescence. *Bulg. J. Agric. Sci., Supplement.*, 19: 39-44.

13. Kumari, T. and S. C. Deka. 2021. Potential health benefits of garden pea seeds and pods: A review. *25:1-13*.
14. Lahuta, L. B., J. Szablińska-Piernik and M. Horbowicz. 2022. Changes in metabolic profiles of pea (*Pisum sativum* L.) as a result of repeated short-term soil drought and subsequent re-watering. *Int. J. Mol. Sci.*, 23: 1-10.
15. Li, H., X. Li, D. Zhang, H. Liu and K. Guan. 2013. Effects of drought stress on the seed germination and early seedling growth of the endemic desert plant *Eremosparton songoricum* (Fabaceae). *EXCLI J.* 12: 89-101.
16. Lu, Z. X., J. F. He, Y. C. Zhang and D. J. Bing. 2020. Composition, physicochemical properties of pea protein and its application in functional foods. *Critical Reviews in Food Science and Nutrition*, 60: 2593-2605.
17. Ma, Z., J. I. Boye and X. Hu. 2018. Nutritional quality and techno-functional changes in raw, germinated and fermented yellow field pea (*Pisum sativum* L.) upon pasteurization. *LWT-Food Science and Technology*, 92: 147-154.
18. Martens, L. G., M. M. Nilsen and F. Provan. 2017. Pea hull fibre: Novel and sustainable fibre with important health and functional properties. *EC Nutrition*, 10: 139-148.
19. Martin, R.J., D. R. Wilson, M. U. Riddle and R. N. Gillespie. 2006. Response of pea seed yield to water deficit. *Agronomy*. 36: 36-43.
20. Melchior, S., S. Calligaris, G. Bisson, and L. Manzocco. 2020. Understanding the impact of moderate-intensity pulsed electric fields (MIPEF) on structural and functional characteristics of pea, rice and gluten concentrates. *Food and Bioprocess Technology*, 13: 2145-2155.
21. Mudryj, A. N., N. Yu and H. M. Aukema. 2014. Nutritional and health benefits of pulses. *Applied Physiology, Nutrition, and Metabolism*. 39: 1197-1204.
22. Nayak, S. N., J. Balaji, H. D. Upadhyaya, C. T. Hash, P. K. Kishor, D. Chattopadhyay, L. M. Rodriguez, M. W. Blair, M. Baum, K. McNally, D. This, D. A. Hoisington and R. K. Varshney. 2009. Isolation and sequence analysis of DREB2A homologues in three cereal and two legume species. *Plant Science*, 177: 460-467.
23. OKÇU, G., M. D. KAYA and M. ATAK. 2005. Effects of Salt and Drought Stresses on Germination and Seedling Growth of Pea (*Pisum sativum* L.). *Turk. J. Agric. For.*, 29: 237-242.
24. Pandey, J., E. Devadasu, D. Saini, K. Dhokne, S. Marriboina, A. S. Raghavendra and R. Subramanyam. 2023. Reversible changes in structure and function of photosynthetic apparatus of pea (*Pisum sativum*) leaves under drought stress. *The Plant Journal*, 113: 60-74.
25. Porcel, R., R. Azcón and J. M. Ruiz-Lozano. 2005. Evaluation of the role of genes encoding for dehydrin proteins (LEA D-11) during drought stress in arbuscular

- mycorrhizal *Glycine max* and *Lactuca sativa* plants. *Journal of Experimental Botany*, 56: 1933-1942.
26. Sah, S. K., K. R. Reddy and J. Li. 2016. Abscisic acid and abiotic stress tolerance in crop plants. *Front. Plant Sci.*, 7: 1-26.
27. Sarani, M., M. Namrudi, S. M. Hashemi and M. M. Raoofi. 2014. The effect of drought stress on chlorophyll content, root growth, glucosinolate and proline in crop plants. *Intl J Farm & Alli Sci.*, 3: 994-997.
28. Yang, X., M. Lu, Y. Wang, Y. Wang, Z. Liu and S. Chen. 2021. Response Mechanism of Plants to Drought Stress. *Horticulturae*, 7: 1-36.
29. You, J. and Z. Chan. 2015. Regulation During Abiotic Stress Responses in Crop Plants. *Front. Plant Sci.*, 6: 1-15.

UNDER PEER REVIEW