

## Distribution and antibiotic resistance profile of Enterobacteriaceae isolates from well-drinking water in the rural communities of Ezza South local government area of Ebonyi State

### Abstract

Enterobacteriaceae as an indicator of water sanitary quality and their frequency of occurrence as antibiotic resistant microbial pathogen in water remain a threat to human and environment. The aim of this research work was to determine the distribution and antibiotic resistance profile of Enterobacteriaceae isolates from different water sources in Ezza South Local Government Area of Ebonyi State, Nigeria. A total of 200 water sample were collected in sterile bottles and subjected to bacteriological analysis using Standard Microbiological protocol for isolation and identification. Antimicrobial resistance studies of Enterobacteriaceae was determined using the Kirby-Bauer disk diffusion method and the results were analyzed and were compared with the Clinical Laboratory Standard Institute (CLSI) zone diameter breakpoints. The result of isolation revealed a high Colony enumeration of bacterial isolates from well water revealed high bacterial counts of  $1.0 \times 10^4$ - $8.1 \times 10^4$  cfu/ml from Nsukara. *E. coli* distribution from borehole water revealed overall occurrence rate of 23(28.8 %) consisting high occurrence rate of 5(50.0 %) in Onueke Urban followed by 3 (30.0 %) in both Amuzu, Echara, UmunwaguIdembia and Amagu/Amaezekwe while the least occurrence of 2(20.0 %) was recorded against Ezzama, Ikwuato/Idembia and Amana respectively while the distribution of bacteria in well water revealed high proportion of *E. coli* 42(42.0 %) followed by *Salmonella* species 12(12.0 %) and *Shigella* species 7(7.0 %). The isolates exhibited high percentage of resistance to ceftriaxone 100 %, Sulthamethoxazole/trimethoprim 100 % tetracycline 66.7 %, 66.7 % but susceptible to Ciprofloxacin 100 %, and Imipenem 100 %. Our findings indicate the presence of antibiotic resistant bacteria in well water and amongst the communities, awareness should be given to the populace on the implication of antibiotic residues in the environment as well as the importance of maintaining a clean and hygienic environment around the wells to ensure the safety of water and also to prevent the spread of resistant determinant in the environment and human. Government should make provision for a portable water source for that will be accessible and well sustained in the communities.

**Keywords:** Well water, *E. coli*, *Salmonella* species, *Shigella* species, Antibiotic resistant, *Enterobacteriaceae*

### Introduction

In most rural locations, well water is the most frequent source of drinking water for residential usage and human consumption. It has grown more challenging to meet all of the water requirements in the rural settlements in the Ezza South local government region due to the mostly nonexistent public water supply, which is also inaccessible when it does exist. This has prompted many homes to resort to unsafe and non-potable water sources, resulting in many digging of wells.

Most Nigerian rural areas do not have access to improved water supplies [1, 2, 3, 4]. Their primary sources of free water are usually rivers, perennial streams, ponds, and unprotected wells, all of which are reservoir of virulence and drug resistant bacteria.

One of the main environmental problems caused by the inappropriate and negligent disposal of sewage, industrial, and chemical waste is groundwater pollution.

Based on multiple research, it has been determined that groundwater is easily contaminated by rainstorm overflows, runoff from farming areas and areas with septic systems and latrines that are improperly situated which makes the water clinically unsafe for human consumption [5, 6].

**Comment [P1]:** All SCIENTIFIC NAMES MUST BE WRITTEN WITH ITALIC

**Comment [P2]:** Not necessary

**Comment [P3]:** 1st too long just remove ones and put it in in the conclusions

**Comment [P4]:** Reduce these key words to 5 oder less

The majority of human diseases are caused by unsanitary drinking water supplies, which cause infections such as dysentery, diarrhea, cholera, and typhoid. It has been stated that over 20% of the world's population has scarcity of safe drinking water, and with over a million people die every year from illnesses related with drinking water due to inadequate sanitation and the convergence of antibiotic-resistant bacteria [7, 8, 9]. Most thriving bacteria family in water is the Enterobacteriaceae [10, 11]. Antibiotic resistance develops naturally, but misuse of antibiotics in humans and veterinary medicine accelerates the process. As a result, water acts not only as a vehicle for the spread of antibiotic-resistant organisms among humans and animals, but also as a reservoir of introducing Enterobacteriaceae harboring resistance genes into natural bacterial ecosystems. The rate of increasing cases of antimicrobial resistance have become a worldwide problem in different ecosystem [12].

Although the present guidelines for water quality are designed to decrease the number of cases of faecal contamination, they do not take into account the possibility that water could serve as a reservoir for antimicrobial resistance determinant that confer resistance against antimicrobials that are clinically important. But determining whether antimicrobial-resistant Enterobacteriaceae are present in well-sourced drinking water would reveal details about the spread and durability of these bacteria, which can live and even become mobilized within the human population and pose a health risk.

#### **Materials and Methods**

##### **Study Area**

The study was carried out at Ezza South local government area, located at latitude 6.1322°N and longitude 8.0216°E in Ebonyi state on south-eastern Nigeria. Its headquarters is Onueke, which also serve as a central unification town for the Ezza nation as well as headquarters of Ebonyi Central senatorial zone. It was created on October 1, 1996, amongst other local government areas in the then new Ebonyi state by the military government of late General Sani Abacha. Ezza South prior to its creation was part of old Ezza Local government area. The people are predominantly of Igbo stock. They speak Ezza dialect and the central Igbo language. Their major occupations are farming and trading as well as emerging civil servant class. It has an area of 324 km<sup>2</sup> and a population of 133,625 at the 2006 census.

##### **Sample Collection**

Following multiple initial visits to villages around the study area, nine communities—Onueke Urban, Amuzu, Ezzama, Echara, Umunwaguldembia, Ikwuato Idembia, Amagu Amaezekwe, Amana, Nsukara, and Amudo/Okoffia—were subsequently identified and selected for sampling.

Wells water was drawn using a fetcher found in the sampling points. A total of 200 water samples, each containing approximately 500 ml of water, were taken from ten wells in each of the selected communities. Water samples were stored in a flask at -4°C, and aseptically transported to the laboratory within 2 hours for bacteriological analysis.

##### **Isolation, Identification and Colony Enumeration of Bacterial Strains**

Ten folds serial dilutions were carried out following standard microbiological procedures, by using 1 ml of each water samples in 9 ml of sterile water [13]. After the dilution, 0.5 ml from dilution factor five (10<sup>5</sup>) was transferred to each sterile petri dish before pouring a sterilized plate count agar (Hi media, Mumbai, India). The plates were incubated at room temperature for 24 hours. After 24 hrs of incubation, colonies were counted using Colony counter (Reichert, Inc. Quebec ®) and a loopful of each colony were aseptically streaked on solidified eosin methylene blue agar plate, *Salmonella Shigella* agar (Hi media, Mumbai, India). The plates were incubated aerobically for 18-24 hrs at 37°C. Bacterial colonies with greenish-metallic sheen on eosin methylene blue agar plate, black centered colonies on *Salmonella Shigella* agar, and smooth and opaque or colorless on *Salmonella Shigella* agar were infer as the presence *Escherichia coli*, *Salmonella* species and *Shigella* species respectively. An API 20E kit (BioMérieux, Marcy l'Etoile, France) was used to identify and differentiate the Gram-negative bacteria of the family Enterobacteriaceae following the manufacturer's instructions.

##### **Antimicrobial Susceptibility Testing**

The antimicrobial susceptibility testing was performed using the Kirby-Bauer disc diffusion method on Mueller Hinton agar based on the Clinical Laboratory Standard Institute (CLSI) standards [14]. The suspension of standard inoculum from the pure culture was adjusted to achieve turbidity equivalent to 0.5 McFarland standard solutions, the suspension was emulsified using sterile cotton swabs onto Mueller Hinton agar plate. The antibiotics to be tested are: Meropenem (10 µg), Gentamicin (30 µg), Imipenem (10 µg), Ciprofloxacin (30 µg), Sulthamethoxazole/trimethoprim (12.5 µg), Ceftriaxone (30 µg), Amikacin (30 µg), Tetracycline (30 µg), Azithromycin (30 µg), Cefepime (30 µg) was placed on the inoculated agar and incubated at 37 °C for 16 to 18 h. The inhibition zone of each antimicrobial agent were interpreted according using CLSI standards [14, 15].

Comment [P5]: No need

**Result**

**Colony enumeration of bacteria isolated from different well water sources in Ezza south LGA**

Colony enumeration of bacteria isolate from well water revealed a high Bacteria Count of  $1.0 \times 10^4$ - $8.1 \times 10^4$  from Nsukara followed by Onueke Urban  $1.0 \times 10^3$ - $1.6 \times 10^3$ , Amuzu  $1.0 \times 10^3$ - $1.0 \times 10^4$ , Ezzama  $0.9 \times 10^3$ - $2.0 \times 10^3$ , Echara  $1.0 \times 10^3$ - $1.0 \times 10^4$ , UmunwaguIdembia  $1.3 \times 10^3$ - $1.4 \times 10^3$ , IkwuatoIdembia  $1.3 \times 10^3$ - $1.0 \times 10^4$ , Amagu/Amaezekwe  $1.3 \times 10^3$ - $1.0 \times 10^4$ , Amana  $1.3 \times 10^3$ - $2.0 \times 10^2$  and Amudo/Okoffia  $1.2 \times 10^3$ - $1.6 \times 10^3$  as shown in Table 1.

**Table 1:** Colony enumeration of bacteria isolated from different well water sources in Ezza south L.G.A

S/No.	Onueke Urban		Amuzu		Ezzama		Echara		UmunwaguIdembia		IkwuatoIdembia	
	Colonies	Cfu/ml	Colonies	Cfu/ml	Colonies	Cfu/ml	Colonies	Cfu/ml	Colonies	Cfu/ml	Colonies	Cfu/ml
1	59	$1.2 \times 10^4$	58	$1.2 \times 10^3$	45	$1.0 \times 10^3$	65	$1.3 \times 10^3$	66	$1.3 \times 10^3$	67	$1.3 \times 10^3$
2	48	$1.0 \times 10^4$	54	$1.0 \times 10^3$	81	$1.6 \times 10^3$	61	$1.2 \times 10^3$	71	$1.4 \times 10^3$	81	$1.6 \times 10^3$
3	47	$1.0 \times 10^4$	70	$1.4 \times 10^3$	58	$1.2 \times 10^3$	51	$1.0 \times 10^3$	75	$1.5 \times 10^3$	77	$1.5 \times 10^3$
4	81	$1.6 \times 10^3$	72	$1.4 \times 10^3$	39	$0.8 \times 10^3$	91	$1.8 \times 10^3$	80	$1.6 \times 10^3$	66	$1.3 \times 10^3$
5	55	$1.1 \times 10^3$	48	$1.0 \times 10^4$	101	$2.0 \times 10^3$	72	$1.4 \times 10^3$	61	$1.2 \times 10^3$	81	$1.6 \times 10^3$
6	56	$1.1 \times 10^3$	45	$1.0 \times 10^4$	60	$1.2 \times 10^3$	65	$1.3 \times 10^3$	58	$1.2 \times 10^3$	70	$1.4 \times 10^3$
7	70	$1.0 \times 10^3$	47	$1.0 \times 10^4$	42	$0.8 \times 10^3$	61	$1.2 \times 10^3$	70	$1.4 \times 10^3$	67	$1.3 \times 10^3$
8	40	$1.0 \times 10^3$	40	$1.0 \times 10^3$	46	$0.9 \times 10^3$	70	$1.4 \times 10^3$	81	$1.6 \times 10^3$	58	$1.2 \times 10^3$
9	50	$1.0 \times 10^3$	79	$1.6 \times 10^3$	50	$1.0 \times 10^2$	48	$1.0 \times 10^4$	67	$1.3 \times 10^3$	60	$1.2 \times 10^3$
10	62	$1.2 \times 10^3$	55	$1.1 \times 10^3$	47	$0.9 \times 10^3$	49	$1.0 \times 10^4$	72	$1.4 \times 10^3$	49	$1.0 \times 10^4$

**Table 1 contd:** Colony enumeration of bacteria isolated from different well water sources in Ezza south L.G.A

S/No.	AmaguAmaezekwe		Amana		Nsukara		Amudo/Okoffia	
	Colonies	Cfu/ml	Colonies	Cfu/ml	Colonies	Cfu/ml	Colonies	Cfu/ml
1	67	1.3x10 <sup>3</sup>	71	1.4x10 <sup>3</sup>	39	1.0x10 <sup>4</sup>	80	1.6x10 <sup>3</sup>
2	80	1.6x10 <sup>3</sup>	80	1.6x10 <sup>3</sup>	81	1.6x10 <sup>3</sup>	72	1.4x10 <sup>3</sup>
3	77	1.5x10 <sup>3</sup>	100	2.0x10 <sup>2</sup>	66	1.3x10 <sup>3</sup>	60	1.2x10 <sup>3</sup>
4	61	1.2x10 <sup>3</sup>	102	2.0x10 <sup>2</sup>	70	1.4x10 <sup>3</sup>	58	1.2x10 <sup>3</sup>
5	58	1.2x10 <sup>3</sup>	66	1.3x10 <sup>3</sup>	81	8.1 x10 <sup>4</sup>	60	1.2 x10 <sup>3</sup>
6	70	1.4x10 <sup>3</sup>	58	1.2x10 <sup>3</sup>	59	1.2x10 <sup>3</sup>	78	1.4x10 <sup>3</sup>
7	48	1.0x10 <sup>4</sup>	96	1.9x10 <sup>3</sup>	48	1.0x10 <sup>4</sup>	80	1.6x10 <sup>3</sup>
8	49	1.0x10 <sup>4</sup>	72	1.4x10 <sup>3</sup>	50	1.0x10 <sup>3</sup>	78	1.4x10 <sup>3</sup>
9	60	1.2x10 <sup>3</sup>	81	1.6x10 <sup>3</sup>	82	1.6x10 <sup>3</sup>	69	1.4x10 <sup>4</sup>
10	71	1.4x10 <sup>3</sup>	68	1.3x10 <sup>3</sup>	73	1.5x10 <sup>3</sup>	58	1.2x10 <sup>3</sup>
<b>Key:</b>		<b>Cfu-Colony</b>		<b>Forming</b>		<b>Unit,</b>		<b>ml-Milligram</b>

### Distribution of bacteria isolated from different well water sources in Ezza South LGA

The distribution of bacteria in well water revealed high proportion of *E. coli* 42(42.0 %) followed by *Salmonella* species 12(12.0 %) and *Shigella* species 7(7.0 %). *E. coli* was highly predominant in samples from Onueke Urban 60.0 % over AmaguAmaezekwe 50.0 %, Nsukara 40.0 % while *Salmonella* species comprising of 50.0 %, 30.0 % and 20.0 % from samples in Amuzu, Amana and Ezzama respectively. *Shigella* species accounted 30.0 %, 20.0 % and 20.0 % from Onueke Urban, Amuzu and Ezzama respectively as shown in Table 2.

**Table 2:** Distribution of bacteria isolated from different well water sources in Ezza South LGA

Location	No. sampled	<i>E. coli</i> (%)	<i>Salmonella</i> species (%)	<i>Shigella</i> species (%)
Onueke Urban	10	6(60.0)	2(20.0)	3(30.0)
Amuzu	10	4(40.0)	5(50.0)	2(20.0)
Ezzama	10	3(30.0)	2(20.0)	2(20.0)
Echara	10	4(40.0)	0(0.0)	0(0.0)
UmunwaguIdembia	10	5(50.0)	0(0.0)	0(0.0)
IkwuatoIdembia	10	4(40.0)	0(0.0)	0(0.0)
AmaguAmaezekwe	10	5(50.0)	0(0.0)	0(0.0)
Amana	10	3(30.0)	3(30.0)	0(0.0)
Nsukara	10	4(40.0)	0(0.0)	0(0.0)
Amudo/Okoffia	10	4(40.0)	0(0.0)	0(0.0)
<b>Total</b>	<b>100</b>	<b>42(42.0)</b>	<b>12(12.0)</b>	<b>7(7.0)</b>

### Antibiotic Susceptibility profile of *Shigella* species isolated from different well Water sources in Onueke Urban Ward, Ezzama Ward, Amuzu Ward in Ezza South LGA

In Onueke Urban ward, *Shigella* species from well water samples were 100 % resistant to tetracycline Trimethoprim-Sulfamethoxazole, Azithromycin, Cefepime and ceftriaxone but were sensitive to meropenem 100 %, Ciprofloxacin 100 %, Imipenem 100 %, Gentamicin 100 %. Isolate from Ezzama Wardwell water sample were resistant meropenem 100%, Amikacin 100%, Tetracycline 100 %, Trimethoprim-Sulfamethoxazole 100 %, Azithromycin 100 %, Cefotaxime 100 %but were susceptible to ciprofloxacin 100 %, Imipenem 100 % and Gentamicin 100%. Majority of the *Shigella* species from well water in AmuzuEzzaweresusceptible to meropenem 50.0 %, Ciprofloxacin 100 %, Imipenem 100 %, Gentamicin 100 % but were resistant to Tetracycline 100 %, Trimethoprim-Sulfamethoxazole 100 %, Cefepime 100 %, Ceftriaxone 100 % and Azithromycin 100 %as presented in Table 3

Comment [P7]: Use the name of the bacteria

Comment [P8]: Check the word

**Table 3:** Antibiotic Susceptibility profile of *Shigella* species isolated from different well Water sources in Onueke, Ezzama, Amuzu communities in Ezza South, Ebonyi State

Ezza South Communities of Ebonyi State						
Antibiotic (µg)	Onueke Urban Ward		Ezzama Ward		AmuzuEzza	
	Well Water (n=3)		Well Water (n=2)		Well Water(n=2)	
	S (%)	R (%)	S (%)	R (%)	S (%)	R (%)
MEM (10)	3(100)	0(0.0)	0(0.0)	2(100)	1(50)	1(50)

CN (30)	3(100)	0(0.0)	2(100)	0(0.0)	2(100)	0(0.0)
IPM (10)	3(100)	0(0.0)	2(100)	0(0.0)	2(100)	0(0.0)
CIP (30)	3(100)	0(0.0)	2(100)	0(0.0)	2(100)	0(0.0)
SXT (12.5)	0(0.0)	3(100)	0(0.0)	2(100)	0(0.0)	2(100)
CRO (30)	0(0.0)	3(100)	0(0.0)	2(100)	0(0.0)	2(100)
AK (30)	0(0.0)	3(100)	0(0.0)	2(100)	0(0.0)	2(100)
TE (30)	0(0.0)	3(100)	0(0.0)	2(100)	0(0.0)	2(100)
AT (30)	0(0.0)	3(100)	0(0.0)	2(100)	0(0.0)	2(100)
FEP (30)	0(0.0)	3(100)	0(0.0)	2(100)	0(0.0)	2(100)

**Key:** Mem: Meropenem (10), CN = Gentamicin 30, IPM = Imipenem 10, CIP = Ciprofloxacin 30, SXT = Sulthamethoxazole/trimethoprim 12.5, CRO = Ceftriaxone 30, Ak = Amikacin 30, TE = Tetracycline 30, ATM = Azithromycin 30, FEP = Cefepime 30 and % = percentage, S= Susceptible, R=Resistance, n=number of isolate

#### Antibiotic Susceptibility profile of *Salmonella* species isolated from different well Water sources in Onueke Urban Ward, Amuzu Ward, Ezzama Ward and in Ezza South LGA

Antibiotic susceptibility profile of *Salmonella* species shows that in Onueke Urban Ward, *Salmonella* species from well water samples were 100 % resistant to tetracycline Trimethoprim-Sulfamethoxazole, Azithromycin, Ceftriaxone, Cefepime, and meropenem but were sensitive to Ciprofloxacin 100 % Imipenem 100 %, Gentamicin 100 % while in Well Water from Amuzu Ward, majority of the isolates were susceptible to Imipenem 100 %, Gentamicin 100 %, ciprofloxacin 100 %, but were resistant to meropenem 40.0 %, Amikacin 60.0 %, Tetracycline 100 %, Trimethoprim-Sulfamethoxazole 100 %, Azithromycin 100 %, Ceftriaxone 100 % and Cefepime 100 %. In Ezzama community, isolate from well water sample demonstrated resistant to Amikacin 66.7 %, Tetracycline 100 %, Trimethoprim-Sulfamethoxazole 100 %, Ceftriaxone 100 %, Azithromycin 100 % and Cefepime 100 % but were susceptible to ciprofloxacin 100 %, Imipenem 100 % and Gentamicin 66.7 % while in well water from Amana Ward the isolate were susceptible to meropenem 66.7 %, Gentamicin 100 %, Imipenem 100 % and ciprofloxacin 100 % respectively but were resistant tetracycline, Cefepime, Ceftriaxone, Trimethoprim-Sulfamethoxazole, Azithromycin recording 100 % respectively as presented in Table 4

**Table 4:** Antibiotic Susceptibility profile of *Salmonella* species isolated from different well Water sources in Onueke Urban Ward, Amuzu Ward, Ezzama Ward and in Ezza South LGA

Antibiotic (µg)	Onueke Urban Ward		Amuzu Ward		Ezzama Ward		Amana Ward	
	Well Water (n=2)		Well Water (n=5)		Well Water (n=2)		Well Water (n=3)	
	S (%)	R (%)	S (%)	R (%)	S (%)	R (%)	S (%)	R (%)
MEM (10)	0(0.0)	2(100)	3(60)	2(40)	0(0.0)	(100)	2(66.7)	1(33.3)
CN (30)	2(100)	0(0.0)	5(100)	0(0.0)	2(66.7)	(33.3)	3(100)	0(0.0)
IPM (10)	2(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	3(100)	0(0.0)
CIP (30)	2(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	3(100)	0(0.0)
SXT (12.5)	0(0.0)	2(100)	0(0.0)	5(100)	0(0.0)	(100)	0(0.0)	3(100)
CRO (30)	0(0.0)	2(100)	0(0.0)	5(100)	0(0.0)	(100)	0(0.0)	3(100)
AK (30)	0(0.0)	2(100)	2(40)	3(60)	1(33.3)	2(66.7)	1(33.3)	2(66.7)
TE (30)	0(0.0)	2(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	3(100)

AT (30)	0(0.0)	2(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	3(100)
FEP (30)	0(0.0)	2(100)	0(0.0)	5(100)	0(0.0)	(100)	0(0.0)	3(100)

**Key:** Mem: Meropenem, CN = Gentamicin, IPM = Imipenem, CIP = Ciprofloxacin, SXT = Sulthamethoxazole/trimethoprim, CRO = Ceftriaxone, Ak = Amikacin, TE = Tetracycline, ATM = Azithromycin, FEP = Cefepime and % = percentage, S=Susceptible, R=Resistance n=number of isolate

**Antibiotic susceptibility profile of *E. coli* isolates from different well water sources in Onueke, Ezzama, Amuzu, Echara and Amudo/Okoffia communities in Ezza South, Ebonyi State**

*Escherichia coli* isolate from well water Onueke were susceptible to Amikacin 50.0 %, Ciprofloxacin 100 %, Imipenem 100 % and Gentamicin 100 % respectively but were resistant to meropenem 66.7 %, Azithromycin 100%, tetracycline 100 %. In Amuzu communities, *E. coli* from well water were resistant to Azithromycin 75.0 %, Trimethoprim-Sulfamethoxazole 100 %, and Ceftriaxone 100% but were sensitive to Amikacin 50.0 %, meropenem 75.0 %, Imipenem 100 %, Gentamicin 100 % and Ciprofloxacin 100 %

*E. coli* isolate from well water samples in Ezzama ward were susceptible to Meropenem 66.7 %, Amikacin 66.7 %, Ciprofloxacin 100 %, Imipenem 100 %, Gentamicin 100 % but were 100 % resistant to Tetracycline 100 %, Trimethoprim-Sulfamethoxazole, Azithromycin, Ceftriaxone 100 % and Cefepime 100 %. Isolate from well water samples in Echara were susceptible to meropenem 75.0 %, Imipenem 100 %, Gentamicin 100 % and Ciprofloxacin 100 % but were resistant to Trimethoprim-Sulfamethoxazole 100 %, Cefepime 100 %, tetracycline 100 % and Ceftriaxone 100 % while in well water from Amudo/Okoffia, the isolate were susceptible to Imipenem 100 %, Gentamicin 100 %, ciprofloxacin 100 %, but were resistant to Amikacin 100 %, Tetracycline 100 %, Azithromycin 100 %, Trimethoprim-Sulfamethoxazole 100 %, meropenem 100 %, Ceftriaxone 100 % and Cefepime 100 % as presented in Table 5

**Table 5:** Antibiotic Susceptibility profile of *E. coli* isolates from different Borehole and well Water sources in Onueke, Ezzama, Amuzu, Echara and Amudo/Okoffia communities in Ezza South, Ebonyi State

Antibiotic (µg)	Onueke/Ezzama		Amuzu/Echara		Amudo/Okoffia					
	Well water (n=6)	Well Water (n=3)	Well water (n=4)	Well Water (n=4)	Well Water (n=4)	Well Water (n=4)				
	S (%)	R (%)	S (%)	R (%)	S (%)	R (%)				
MEM (10)	2(33.3)	4(66.7)	2(66.7)	1(33.3)	3(75)	1(25)	3(75)	1(25)	0(0.0)	4(100)
CN (30)	6(100)	0(0.0)	3(100)	0(0.0)	4(100)	0(0.0)	4(100)	0(0.0)	4(100)	0(0.0)
IPM (10)	6(100)	0(0.0)	3(100)	0(0.0)	4(100)	0(0.0)	4(100)	0(0.0)	4(100)	0(0.0)
CIP (30)	5(83.3)	1(16.7)	3(100)	0(0.0)	4(100)	0(0.0)	4(100)	0(0.0)	4(100)	0(0.0)
SXT (12.5)	0(0.0)	6(100)	0(0.0)	3(100)	0(0.0)	4(100)	0(0.0)	4(100)	0(0.0)	4(100)
CRO (30)	0(0.0)	6(100)	0(0.0)	3(100)	0(0.0)	4(100)	0(0.0)	4(100)	0(0.0)	4(100)
AK (30)	3(50)	3(50)	2(66.7)	1(33.3)	2(50)	2(50)	0(0.0)	4(100)	0(0.0)	4(100)
TE (30)	0(0.0)	6(100)	0(0.0)	3(100)	0(0.0)	4(100)	0(0.0)	4(100)	0(0.0)	4(100)
AT (30)	0(0.0)	6(100)	0(0.0)	3(100)	1(25)	3(75)	0(0.0)	4(100)	0(0.0)	4(100)
FEP (30)	0(0.0)	6(100)	0(0.0)	3(100)	0(0.0)	4(0.0)	0(0.0)	4(100)	0(0.0)	4(100)

**Key:** Mem: Meropenem, CN = Gentamicin, IPM = Imipenem, CIP = Ciprofloxacin, SXT = Sulthamethoxazole/trimethoprim, CRO = Ceftriaxone, Ak = Amikacin, TE = Tetracycline, ATM = Azithromycin, FEP = Cefepime and % = percentage, S=Susceptible, R=Resistance, n=number of isolate

**Antibiotic Susceptibility profile of *E. coli* isolated from different Borehole and well Water sources in Umunwagu/Idenbua and Ikwuato/Idenbua in Ezza South LGA**

Isolate from Umunwagu/Idenbua well water sample were resistant meropenem 60.0%, Amikacin 60.0 %, Tetracycline 100 %, Trimethoprim-Sulfamethoxazole 100 %, Cefotaxime 100 % but were susceptible to ciprofloxacin 100 %, Imipenem 100 % and Gentamicin 100%, while the isolate from Well Water were sensitive to meropenem 75.0 %, Imipenem 100 %, ciprofloxacin 100 %, Gentamicin 100 % but were resistant tetracycline 100 %, Trimethoprim-Sulfamethoxazole, Azithromycin,

Ceftriaxone, Cefepime recording 100 % while in Well Water from Amagu/amezekwe., majority of the isolates were susceptible to Imipenem 100 %, Gentamicin 100 %, ciprofloxacin 100 %, but were resistant to Tetracycline 100 %, Trimethoprim-Sulfamethoxazole 100 %, Azithromycin 100 %, Ceftriaxone 100 % and Cefepime 100 %. and in well water from Amana Ward the isolate were susceptible to meropenem 66.7 %, ciprofloxacin 100 %, Imipenem 100 %, Gentamicin 100 % respectively but were resistant tetracycline, Cefepime, Ceftriaxone, Trimethoprim-Sulfamethoxazole, Azithromycin recording 100 % and Antibiotic susceptibility profile of *Escherichia coli* shows that in Nsukara, *E. coli* from well water samples were 100 % resistant to Ceftriaxone, Cefepime, tetracycline Trimethoprim-Sulfamethoxazole, Azithromycin but were sensitive to meropenem 66.7 %, Ciprofloxacin 100%, Imipenem 100 %, Gentamicin 100 % respectively as presented in Table 6

**Table 6:** Antibiotic Susceptibility profile of *E. coli* isolates from different Borehole and well Water sources in UmunwaguIdenbiaIkwuato, Amagu/amezekwe, Amana and Nsukara communities in Ezza South, Ebonyi State

Antibiotic (µg)	UmunwaguIdenbiaIkwuato		Amagu/amezekwe		Amana		Nsukara			
	Well water (n=5)	Well Water (n=4)	Well water (n=5)	Well Water (n=3)	Well Water (n=3)	Well Water (n=3)	Well Water (n=6)	Well Water (n=6)		
	S (%)	R (%)	S (%)	R (%)	S (%)	R (%)	S (%)	R (%)	S (%)	R (%)
MEM (10)	2(40)	3(60)	3(75)	1(25)	3(60)	2(40)	2(66.7)	1(33.3)	4(66.7)	2(33.3)
CN (30)	5(100)	0(0.0)	4(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	6(100)	0(0.0)
IPM (10)	5(100)	0(0.0)	4(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	6(100)	0(0.0)
CIP (30)	4(80)	2(40)	4(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	6(100)	0(0.0)
SXT (12.5)	0(0.0)	5(100)	0(0.0)	4(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	6(100)
CRO (30)	0(0.0)	5(100)	0(0.0)	4(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	6(100)
AK (30)	2(40)	3(60)	0(0.0)	4(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	6(100)
TE (30)	0(0.0)	5(100)	0(0.0)	4(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	6(100)
AT (30)	0(0.0)	5(100)	0(0.0)	4(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	6(100)
FEP (30)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	5(100)	0(0.0)	3(100)	0(0.0)	6(100)

## Discussion

Many different species of bacteria were isolated from well-sourced drinking water for our study. These include *Salmonella* species, which can cause typhoid fever and acute diarrheal infections, *Shigella* species, which can cause dysentery, and *E. coli*, which can cause gastroenteritis, urinary tract disease, neonatal meningitis, hemorrhagic colitis, and Crohn's disease.

Kalu *et al.* [16] stated that research by the WHO indicates that about 80% of diseases globally are linked to the consumption and use of contaminated water.

The presence of coliform bacteria such as *E. coli*, in these well water samples make them unsafe for drinking for human consumption. Members of the coliform group were found in drinking water in rural Peshawar, India [17], in well water in Shagamu and Iwo Nigeria [18, 19] and also in stream water [20]. Although no differentiation were made between pathogenic and non-pathogenic *E. coli* when this species is isolated from well water, therefore water that contains *E. coli* is unsafe for consumption due to strong association between *E. coli* and fecal contamination.

The high percentage of *Salmonella* species and *E. coli* in this study implies that the well in may be contaminated with feces due to human and animal activity. This finding supports previous report by Ekelozie *et al.* [21] and Tangwa *et al.* [22], who found *Salmonella* spp. and *E. coli* as the most prevalent bacteria in hand dug wells and boreholes in Ngaoundere municipality of Adamawa region in Cameroon, as well as water sources in two local government areas of Anambra State, Nigeria.

According to Gugu *et al.* [20], the most often isolated bacterium was *Salmonella* spp. [18(22.8%)], followed by *E. coli* [16(20.3%)]. In a related study, Geta and Kibret [23] found that *E. coli*, *Streptococcus* spp., *Klebsiella* spp., *Pseudomonas* spp., *Citrobacter* spp., and *Salmonella* spp. were the most commonly found bacteria from waste sources in Hotspot Environments in Bahir Dar City, Northwestern Ethiopia.

The presence of intestinal bacteria species such as *Salmonella*, *Proteus*, *Shigella* and *E. coli* indicates that the water was likely contaminated with feces. Enterobacteria species, especially *E. coli*, are commonly found organisms in various water sources such as rivers, streams, rainwater, well water, groundwater and even tap water [24].

Comment [P9]: Please mix/assign this discussion with ist related results

The failure of the government in providing safe drinking water led to people sourcing potable water by themselves by digging wells for household use. The total bacterial counts and total coliform counts in all water samples analyzed in this study indicated a high microbial load in the water. Colony enumeration of bacteria isolate from well water revealed a high Bacteria Count of  $1.0 \times 10^4$ - $8.1 \times 10^4$  from Nsukara followed by Onueke Urban  $1.6 \times 10^3$  which comparatively exceeds the WHO's standard limits. The standard limit of  $1.0 \times 10^3$  CFU/ml for heterotrophic bacteria and coliforms was set by WHO and USEPA [25, 26]. This could be due to the wells continually receiving dirty water from surface runoff and seepage from contaminated groundwater. Some of the wells are located in densely populated areas and receive doses of feces from the septic tank, water from an abattoir, sewage water, and pit latrines; also, some of the wells have been exposed. These results are similar to previous reports of high coliform counts in well and borehole waters analyzed. Also, the unhygienic nature of storing and placing of the fetcher may facilitate the spread of numerous bacteria into the well. This result aligns with findings from earlier research on the microbial assessment of drinking-water sources in Boko, Plateau State, Abakaliki Metropolis, Ebonyi State and Ekosodin Community, Edo state, where a high prevalence of heterotrophic bacteria and coliforms were found, many of which exceeded the acceptable water quality limits [20, 27, 28].

Resistance to third- and fourth generation cephalosporins was also frequent. Similar pattern of resistance has been documented in Environmental Water Sources from Southern Chile and water supplies used in poultry production in Ashanti region of Ghana [29, 30]. Atobatele, and Owosen [19] reported Multi-antibiotic-resistant bacteria were present in all (30/30) of the well water samples, and a high number of the identified bacteria (80%) were resistant to all antibiotics in the cephalosporins group. Cephalosporins resistant may attributed to the presence of ESBL-coding genes (*bla*TEM and *bla*CTX-M) in some strains and have been previously found in water sources [29, 31, 32, 33] indicating potential horizontal gene transfer between environmental and pathogenic bacteria [34, 35]. In addition, due to increased anthropogenic activities and incessant intake of antibiotics, many enterobacteria have acquired antibiotic-resistant genes (ARGs) and become pathogenic ARB.

*E. coli* demonstrated MDR to Azithromycin, sulphamethoxazole/trimethoprim and tetracycline. The findings of this study confirm the pattern of antibiotic resistance in *E. coli* isolates obtained from water similar to the study conducted by Sayah *et al.* [36] revealed that *E. coli* isolates obtained from surface water samples exhibited resistance to drugs such tetracycline and sulphamethoxazole/trimethoprim. The results align with a study conducted by Wose-Kinge *et al.* [37] that examined the antibiotic resistance profiles of *E. coli* isolates obtained from several water sources in the Mmabatho location of South Africa. The isolates were shown to possess resistance against antibiotics like tetracycline. The frequency of multi-drug resistant *E. coli* is consistent with a report by Ogunleye *et al.* [38], who revealed that all isolates of *E. coli* from poultry in Abeokuta, Southwestern Nigeria, were multi-drug resistant. According to Bueno *et al.* [39], these results emphasize the potential for these common water-borne bacteria to spread, as well as the existence of ARGs such as  $\beta$ -lactamase genes (*bla*SHV, *bla*CTX-M, *bla*KPC, and *bla*TEM) and other markers of antibiotic resistance that are frequently used in both human and animal medicine (quinolones, tetracyclines, and sulfonamides, among others). Such critically important pathogens, demand the urgent development of novel therapeutic strategies.

Between this to drug class aminoglycoside and carbapenems; most bacteria were resistant to amikacin but susceptible to gentamicin. Similar pattern was found along with some resistance to carbapenems (meropenem) over strain susceptible to imipenem. Although no alleles associated with mobile resistance were detected. Therefore, amikacin and meropenem resistance would be attributed to the presence of some intrinsic or acquired resistance mechanisms encoded at the chromosome level in these isolates.

This is a red flag in drug prescription due to the observed pattern of resistance against most of the tested antibiotic, could be reduced as majority of the isolate were susceptible with the range of 80-100 % to Gentamicin, ciprofloxacin and imipenem and will likely makes treatment of bacterial infections less difficult and reduces the severity of life threatening infection.

### Conclusion

Considering the presence of these ARB in the well drinking water, enterobacteria in the water samples presented a resistance phenotype to at least one antibiotic, the need for constant monitoring of the changes in their abundance and diversity at the different well drinking water source. Amongst the communities, awareness should be given to the populace on the implication of antibiotic residues in the environment as well as the importance of maintaining a clean and hygienic environment around the wells to ensure the safety of water. It is also advisable that every individual should embark on in-house water treatment in order to avoid water-borne diseases. A recommended

Comment [P10]: What is this? Acronym are needed

distance of 50–100 feet from potential sources of groundwater contamination like soakaways, pit latrines, etc., by health authorities should be maintained. Finally, it should be noted that, although the samples obtained are not representative of all well drinking water in Ebonyi State, Nigeria, it would be interesting to increase the sampling of well drinking water and the identification of GNB not only in the state but also in the regions of the country, in order to correlate the presence of ARG-encoding species with different environmental factors and also relate them to genomic transfer events

Comment [P11]: ???????

Comment [P12]: No ending point???

## REFERENCES

1. Ishaku, J. M., Ahmed, A. S. & Abubakar, M. A. 2011 Assessment of groundwater quality using chemical indices and GIS mapping in Jada. *Journal of Earth Sciences and Geotechnical Engineering* 1 (1), 35–60.
2. Abubakar, I. R. 2018 Strategies for coping with inadequate domestic water supply in Abuja, Nigeria. *Water International* 43 (1), 1–21.
3. Balogun, I. I., Sojobi, A. O. & Galkaye, E. 2017 Public water supply in Lagos State, Nigeria: review of importance and challenges, status and concerns and pragmatic solutions. *Cogent Engineering* 4, 1329776
4. Akoteyon, I. S. 2019 Factors affecting household access to water supply in residential areas in parts of Lagos metropolis, Nigeria. *Bulletin of Geography. Socio-Economic Series* 43 (43), 7–24.
5. Sparks, D. L. (2005). Toxic metals in the environment: the role of surfaces. *Elements* 1, 193–197.
6. Mile, I. I., Jande, J. A. & Dagba, B. I. 2012 Bacteriological contamination of well water in Markurdi Town, Benue State, Nigeria. *Pakistan Journal of Biological Sciences* 15 (21), 1048–1051.
7. Karnwal, A., Dohroo, A. & Mannan, M. A. 2017 Microbial analysis of potable water and its management through useful plant extracts. *International Journal of Sciences and Research* 73 (3), 44–49.
8. Haseena, M., Malik, M.F., Javed, A., Arshad, S, and Hanif, J. (2017). Water pollution and human health. *Journal of Environmental Risk Assessment Remedy* 1(3):16-9.
9. Singh, A.K., Das, S., Kumar, S., Gajamer, V.R., Najar, I.N., Lepcha, Y.D., Tiwari, H.K. (2020). Distribution of Antibiotic Resistant Enterobacteriaceae Pathogens in Potable Spring Water of Eastern Indian Himalayas: Emphasis on Virulence Gene and Antibiotic Resistance Genes in *Escherichia coli*. *Journal of Microbiology*; 11:5-8.
10. Djenadi, K. (2017). Antibiotic Resistance Bacteria from Rivers Water in Algeria. *Journal of Bacteriology* 8:3-19.
11. Alawi, M., Smyth, C., Drissner, D., Zimmerer, A., Leupold, D., Müller, D., Do, T. T., Velasco-Torrijos, T and Walsh, F (2024). Private and well drinking water are reservoirs for antimicrobial resistant bacteria. *Journal of antimicrobials and resistance*, 2:7-23
12. Teklu, D. S., Negeri, A. A., Legese, M. H., Bedada, T. L., Woldemariam, H. K and Tullu, K. D(2019). Extended-spectrum beta lactamase production and multi-drug resistance among Enterobacteriaceae isolated in Addis Ababa, Ethiopia. *Antimicrobial Resistance and Infection Control*, 8:39-89.
13. Joseph IS, Okolo IO, Udenweze EC, Nwankwo CE, Peter IU, Ogbonna IP, Iroha IR, Comparison of Antibiotic-Resistant Pattern of Extended Spectrum Beta-Lactamase and Carbapenem-Resistant *Escherichia Coli* Isolates from Clinical and Non-Clinical Sources, *Journal of Drug Delivery and Therapeutics*. 2023; 13(7):107-118
14. Clinical and Laboratory Standards Institute (CLSI) Performance Standards for Antimicrobial Susceptibility Testing, 28th ed.; Supplement M100; Clinical and Laboratory Standards Institute: Wayne, PA, USA, 2019.
15. Oke B., Iroha I. R., Moses I. B., Egbu I. H., Elom E., Uzoh C. V., Nwode V.F., Okpada J. O., Agbom J. N., Peter I U and Ibeka G. U (2020). Killing Rate Kinetics of Commercially Available

Brands of Ciprofloxacin and Cefotaxime on Clinical Bacterial Isolates Subjected to *in vitro* Antibiotic Treatments. *International Journal of Pharmaceutical Science Review Research*, **64**(2):87-97.

16. Kalu, CM., Mudau, KL., Masindi, V., Ijoma, GN and Tekere, M (2024). Occurrences and implications of pathogenic and antibiotic-resistant bacteria in different stages of drinking water treatment plants and distribution systems. *Heliyon* 10 (2024) e26380
17. Amin, R., Ali, S. S., Anwar, Z. & Khattak, J. Z. K. 2012 Microbial analysis of drinking water and water distribution system in new urban Peshawar. *Current Research Journal of Biological Sciences* 4 (6), 731–737.
18. Idowu, O. A., Martins, O., Oluwasanya, O., Awomeso, J. A. & Olaoye, O. A. 2015 Groundwater recharge using baseflow recession analysis In southwestern Nigeria. *Ife Journal of Science* 17
19. Atobatele, B. O and Owosen A (2023). Distribution of multiple antibiotic-resistant Gram-negative bacteria in potable water from hand-dug wells in Iwo, Nigeria. *H2Open Journal* Vol 6 No 1, 40 doi: 10.2166/
20. Gugu M. F., Vomagai K. N., Mwanja P. Y., Ombugadu A., Adejoh V. A. and Attah S. A. (2023). Antibiotic Resistance Profile of Bacteria Isolated from Stream Water in Bokkos Local Government Area of Plateau State, Nigeria. *UMYU Scientifica*, 2(3), 009 – 014
21. Ekelozie, I. S., Ekejindu, I. M., Ochiabuto, O., Obi, M., Onwuasony, U. and Obeagu, E. I. (2018). Evaluation of Salmonella Species in Water Sources in Two Local Government Areas of Anambra State. *Cohesive J Microbiol infect Dis.*, 1(1). CJMI.000501.
22. Tangwa, B. V., Keubou, H., Nfor, E. N. and Ngakou, A. (2019). Antimicrobial Resistance Profile of Bacteria Isolated from Boreholes and Hand Dug Wells Water in Ngaoundere Municipality of Adamawa Region in Cameroon. *Advances in Microbiology*, 9: 629-645
23. Geta, K, and Kibret, M. (2022). Antibiotic Resistance Profiles of Bacteria Isolated from Hotspot Environments in Bahir Dar City, Northwestern Ethiopia. *Journal of Multidiscipline and Healthcare*, 15:1403-1414.
24. Environment Protection Agency (EPA). (2002). US Environment Protection Agency: Safe Drinking Water Act Amendment, 2002. P. EPA 816 – F – 03 – 016.
25. WHO 2002 Aeromonas. In: Guidelines for Drinking-Water Quality, 2nd edn (World Health Organization, ed.) Addendum Microbiological agents in drinking water. World Health Organization, Geneva.
26. World Health Organisation (2018). Fact Sheet: Drinking Water. Geneva: World Health Organization
27. Onwa, N. C., Uzomaka, I. C., Maduako, A. L., Elom, E. E., Ikeanumba, M. O. and Nwode, V. F. (2019). Antibiotic Susceptibility of Bacterial Species Isolated from Underground Waters in Abakaliki Metropolis of Ebonyi State, Nigeria. *International Journal of Pharmaceutical Science Invention(IJPSI)*, 8(2):55-65.
28. Ekhosuehi, A., Akharam, M. O. and Obayuwana, P. (2018). Bacteriological Quality and Antibigram of Isolates from Potable Water Sources in Ekosod in Community, Benin City, Nigeria. *Journal of Applied Science and Environmental Management*, 22(1) 129-133.

29. Jofré Bartholin, M.; Barrera Vega, B.; Berrocal Silva, L. Antibiotic-Resistant Bacteria in Environmental Water Sources from Southern Chile: A Potential Threat to Human Health. *Microbiol. Res.* **2023**, *14*, 1764–1773.
30. Osei, F. B., Boamah, V. E., Boakye, Y. D., Agyare, C. and Abaidoo, R. C. (2021). Antibiotic Resistance of Bacteria Isolated from Water Supplies Used in Poultry Production in Ashanti Region of Ghana. *Open Microbiology Journal*, 15:7-15.
31. Despotovic, M.; De Nies, L.; Busi, S.B.; Wilmes, P. Reservoirs of antimicrobial resistance in the context of One Health. *Curr. Opin. Microbiol.* **2023**, *73*, 102291.
32. Caltagirone, M.; Nucleo, E.; Spalla, M.; Zara, F.; Novazzi, F.; Marchetti, V.M.; Piazza, A.; Bitar, I.; De Cicco, M.; Paolucci, S.; et al. Occurrence of extended spectrum  $\beta$ -lactamases, KPC-type, and MCR-1.2-producing Enterobacteriaceae from wells, river water, and wastewater treatment plants in Oltrepò Pavese area, northern Italy. *Front. Microbiol.* **2017**, *8*, 2232.
33. Said, B.L.; Jouini, A.; Klibi, N.; Dziri, R.; Alonso, C.A.; Boudabous, A.; Ben Slama, K.; Torres, C. Detection of extended-spectrum beta-lactamase (ESBL)-producing Enterobacteriaceae in vegetables, soil and water of the farm environment in Tunisia. *Int. J. Food Microbiol.* **2015**, *203*, 86–92
34. Orji CO, Ogba RC, Emeruwa AP, Peter IU, Uzoeto HO, Agumah BN. Prevalence of Staphylococcus aureus nasal carriage that have developed resistance to second and third generation cephalosporins, Journal of Advances in Microbiology Research. 2024;5(1):130-135.
35. Ilang, D. C., Peter, I. U., Iroha, I. R (2023). Characterization of VIM, VEB and CTX-M Beta-lactamase Gene in *Escherichia coli* and *Pseudomonasaeruginosa* Isolated from Urine Samples of Patients Visiting a Tertiary Hospital in Abakaliki, *International Journal of Medical Sciences and Pharma Research*,9(4):7-11
36. Sayah RS, Kaneene JB, Johnson Y, Miller R. Patterns of antimicrobial resistance observed in *Escherichia coli* isolates obtained from domestic- and wild-animal fecal samples, human septage, and surface water Appl Environ Microbiol 2005; 71(3): 1394-404.
37. Wose-Kinge CN, Ateba CN, Kawadza DT. Antibiotic resistance profiles of *Escherichia coli* isolated from different water sources in the Mmabatho locality, North-West Province, South Africa S Afr J Sci 2010; 106(1-2): 44-9.
38. Ogunleye AO, Oyekunle MA, Sonibare AO. Multidrug resistant *Escherichia coli* isolates of poultry origin in Abeokuta, South Western Nigeria Vet Arh 2008; 78(6): 501-9.
39. Bueno, I.; Verdugo, C.; Jimenez-Lopez, O.; Alvarez, P.P.; Gonzalez-Rocha, G.; Lima, C.A.; Travis, D.A.; Wass, B.; Zhang, Q.; Ishii, S.; et al. Role of wastewater treatment plants on environmental abundance of Antimicrobial Resistance Genes in Chilean rivers. *Int. J. Hyg. Environ. Health* **2020**, *223*, 56–64.