

# Crude protein, economics and water use efficiency of ryegrass under different irrigation regimes and nitrogen levels

## ABSTRACT

A field experiment was undertaken at the Instructional-cum-Research (ICR) Farm, Assam Agricultural University, Jorhat during 2017-2018 and 2018-2019 in order to study the effect of irrigation regimes and nitrogen levels on crude protein, economics and water use efficiency of ryegrass. The experiment was laid out in split-plot design with three replications. The treatments consisted of five levels of irrigation in main plot viz., I<sub>0</sub>: Rainfed, I<sub>1</sub>: Irrigation at critical growth stages, I<sub>2</sub>: Irrigation at IW:CPE ratio of 1.0, I<sub>3</sub>: Irrigation at IW:CPE ratio of 1.2 and I<sub>4</sub>: Irrigation at IW:CPE ratio of 1.4 along with four levels of N - N<sub>0</sub>: 0 kg N/ha, N<sub>1</sub>: 30 kg N/ha, N<sub>2</sub>: 60kg N/ha and N<sub>3</sub>: 90 kg N/ha in sub-plots. The soil of the experimental site was sandy loam in texture, acidic in reaction, medium in organic carbon, medium in available N, available P<sub>2</sub>O<sub>5</sub> and low in available K<sub>2</sub>O. The study revealed that application of irrigation at IW:CPE ratio of 1.4 recorded the highest crude protein yield, highest value in terms of evapotranspiration, total water use, crop water use efficiency, field water use efficiency, gross return, net return and B:C ratio. Application of 90 kg N/ha recorded the highest crude protein yield. The highest values in terms of evapotranspiration, total water use, crop water use efficiency, field water use efficiency, gross return, net return and B:C ratio were observed under 90 kg N/ha.

**Key words:** Irrigation; nitrogen; crop water use efficiency; field water use efficiency; economics.

## 1. INTRODUCTION

Since the times of immemorial, livestock plays an important role in agriculture and it contributed to the growth of national economy through providing milk, meat, wool, various milk products and farm yard manure. Moreover animal has been considered as the major source of energy for draught power in performing various agricultural operations among farmers in rural India. The success of livestock industry mainly depended upon the availability and supply of nutritious and adequate quantity of good quality green fodder to meet the nutritional requirement for the growth, maintenance and production of livestock. Annual ryegrass (*Lolium multiflorum*) is an important and promising cereal fodder which grow well under agro-climatic conditions of Assam. The grasses are leafy and produced high quality and digestible fodder. It responds considerably to applied fertilizer and soil water. Jat et al.[1] found that Five irrigations produced significantly higher crude protein, crude fat, mineral matter and total digestible nutrients of fodder oats but there was no significant differences in crude fibre, nitrogen free extract and total digestible nutrients at first cutting whereas minimum were noted with two irrigations in all the above parameters productions. Tahir et al.[2] found maximum crude fibre (37.35%), crude protein (9.13%) and total ash (14.4%) of oats with six irrigations at 21, 35, 42, 56, 70 and 84 DAS than other treatments. Scheduling irrigation at IW/CPE ratio 0.5 recorded higher WUE in summer (20.7 kg/ha-cm) and *kharif* (17.4 kg/ha-cm) over 0.75 and 1.0 IW/CPE ratio in sandy clay loam

**Comment [SN1]:** From where did you get this wheat varieties?

**Comment [SN2]:** Use each word one time in title of the manuscript

**Comment [SN3]:** Please specify the whole abstract  
Write experimental sites first then weather condition, nutrients requirement, experimental design, best performing treatments parameters and value of that parameters in high performing irrigation and nitrogen levels.

**Comment [SN4]:** add the full form instead of using abbreviation (IW, CPE and B:C)

**Comment [SN5]:** Need to improvement first enter worldwide data

**Comment [SN6]:** Rephrase it and add references where needed

soil [3]. Irrigation scheduling at IW/CPE ratio 0.6 recorded higher WUE (6.96 kg/ha-mm) of maize over ratios of 0.8 (6.37 kg/ha-mm) and 1.0 (6.33 kg/ha-mm) in sandy soil during rabi season [4]. Irrigations scheduled as per critical growth stages recorded higher WUE (5.86 kg/ha-mm) of maize over 40 mm (3.29 kg/ha-mm), 60 mm (3.86 kg/ha-mm) and 80 mm (4.19 kg/ha-mm) CPE [5]. Irrigation level at 1.1 IW:CPE recorded maximum gross return (Rs. 27100 ha<sup>-1</sup>), net return (Rs. 15014 ha<sup>-1</sup>) and B:C of 1.24 while, minimum gross return (Rs. 19950 ha<sup>-1</sup>), net return (Rs. 8599 ha<sup>-1</sup>) and B:C of 0.79 average of two years data recorded irrigation level at 0.7 IW:CPE [6]. The fertility levels up to 110 kg N ha<sup>-1</sup> produced significantly higher crude protein, crude fibre, crude fat, mineral matter, nitrogen free extract and total digestible nutrient production of fodder oat over 70 and 90 kg N ha<sup>-1</sup> whereas crude fibre production did not influence by nitrogen levels at first cutting [1]. Increase in nitrogen level from 0 to 160 kg ha<sup>-1</sup> decreased the dry matter content, digestibility and cell content whereas the reverse trend was noticed for crude protein content, acid-detergent fibre, neutral detergent fibre and hemicelluloses content of oat [7]. Increase in nitrogen levels increased crude protein content but reduced crude fibre content of maize forage [8]. With regard to the effect of nitrogen on the percentage of crude protein of teff grass, it was found that increasing the level of nitrogen increased the percentage of crude protein [9]. Application of nitrogen up to 120 kg N ha<sup>-1</sup> produced significantly higher crude protein yield of fodder oat (9.38 kg ha<sup>-1</sup>) [10]. Jehangir et al. [11] found that increase in fertility level upto 150 kg N ha<sup>-1</sup> increased crude protein content and crude fiber content of oats. Mahdi et al. [12] reported that increase in N level from 60 to 120 kg/ha significant increase in protein content of maize. Shehzad et al. [13] found that significant increase in crude protein (10.52%), crude fat (2.82%), crude fibre (31.77%) and ash content (10.54%) in maize with application of 180 kg N/ha over other nitrogen levels. Dadarwalet al. [14] reported that increasing dose of fertilizer application increase the net returns and B:C ratio. They found higher net return (Rs. 18,910) and B:C ratio (2.47) with application of 180:60:45 kg NPK/ha. Application of 80 kg N ha<sup>-1</sup> recorded maximum gross return (Rs. 56232.00 ha<sup>-1</sup>) and net return (Rs. 42853.74 ha<sup>-1</sup>) and B:C of 1:3.20 [15]. Binadhaniet al. [16] reported application of nitrogen upto 120 kg ha<sup>-1</sup> significantly give higher net return and B:C ratio.

## 2. MATERIALS AND METHODS

Experiment was conducted during 2017-2018 and 2018-2019 at the Instructional-cum-Research (ICR) Farm, Assam Agricultural University, Jorhat. The experiment was laid out in a split-plot design with three replications. The treatments consisted of five levels of irrigation in main plot viz., Rainfed, Irrigation at critical growth stages, Irrigation at IW: CPE ratio of 1.0, Irrigation at IW:CPE ratio of 1.2 and Irrigation at IW:CPE ratio of 1.4 along with four levels of N- 0 kg N/ha, 30 kg N/ha, 60kg N/ha and 90 kg N/ha in sub-plots. Ryegrass variety Makhan grass at the seed rate of 20 kg/ha were dry seeded in the research plots. The nutrients were applied in the form urea, single super phosphate (SSP) and muriate of potash (MOP) as per requirement in the treatment. Nitrogen was applied in three split doses i.e. 1/2 of N is applied in final ploughing, 1/4 at 1<sup>st</sup> cut and remaining 1/4 at 2<sup>nd</sup> cut as per the treatment. All the phosphatic and potassic fertilizers were applied at the rate of 188 kg/ha of SSP and 50 kg/ha of MOP, respectively one day ahead of sowing ryegrass. Each sub-plot was provided with a uniform depth of 6 cm irrigation for ryegrass crop according to different IW:CPE ratios.

### Evapotranspiration by crop

**Comment [SN7]:** Rephrase it and don't discuss whole experiment in research paper like thesis describe it in 1-2 lines

**Comment [SN8]:** Don't use the quantity of source but use the amount of nutrients

**Comment [SN9]:** Also mentioned method or instrument used for water calculation for WUE

**Comment [SN10]:** Check the font size Font size should be same throughout the manuscript

The evapotranspiration by crop was computed from the soil moisture data by using the following formula:

$$ET = \sum_{i=1}^n (E_0 \times K_c K_p) + \sum_{i=1}^n \frac{(M_{1i} - M_{2i})}{100} \times ASG_i \times D_i + ER + GWC$$

Where,

ET= Evapotranspiration (cm)

E0 = Pan evaporation value (cm) from USWB class A pan evaporimeter from the day of irrigation to the day when sampling in wet soil is possible

N = Time interval (days)

M1i = Per cent soil moisture of the ith layer on the date of sampling after irrigation

M2i = Per cent soil moisture of the ith layer on the date of sampling before irrigation

ASGi = Apparent Specific Gravity of ith soil layer

Di = Depth (cm) of the ith layer of the soil

ER= Effective Rainfall during the period under consideration

n= Number of soil layers

Kc= Crop coefficient

GWC= Ground water contribution

### Water use efficiency

Water use efficiency (WUE) of the crop for different treatments was calculated as follows:

$$\text{Crop WUE (kg/ha-cm)} = \frac{\text{Dry matter yield (kg/ ha)}}{\text{ET of crop (cm)}}$$

$$\text{Field WUE (kg/ha-cm)} = \frac{\text{Dry matter yield (kg/ ha)}}{\text{Total water used (cm)}}$$

## 3. RESULTS AND DISCUSSION

### Crude protein content

The effect of irrigation regimes on crude protein content of ryegrass was found to be non-significant at all three cuts, during both the years (Table 1). The highest crude protein content was found irrigation at IW:CPE ratio of 1.4 and lowest was recorded in rainfed treatment. The effect of nitrogen levels on crude protein content of ryegrass was found to be significant at all three cuts, during both the years (Table 1). The highest crude protein content was recorded with the application of 90 kg N/ha followed by 60 kg N/ha. The lowest crude protein content observed under 0 kg N/ha. Higher application of N resulted in an increased in the crude

**Comment [SN11]:** If data were significant then use lettering from maximum to minimum

protein content because there was better assimilation of N resulting on increased protein synthesis.

### Interaction effect

The interaction effect between irrigation regimes and nitrogen levels was found to be non-significant on crude protein content at different growth stages of ryegrass during both the years.

**Table 1. Effect of irrigation regimes (I) and nitrogen levels (N) on crude protein content of ryegrass**

Treatments	Crude protein (%)					
	1 <sup>st</sup> Year			2 <sup>nd</sup> Year		
	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut
<b>Irrigation regimes (I)</b>						
I <sub>0</sub>	7.61	8.24	8.13	7.60	8.27	8.14
I <sub>1</sub>	7.82	8.38	8.27	7.84	8.43	8.32
I <sub>2</sub>	8.12	8.87	8.78	8.10	8.89	8.86
I <sub>3</sub>	8.38	9.12	8.98	8.35	9.17	9.10
I <sub>4</sub>	8.54	9.23	9.16	8.62	9.25	9.17
S.Ed (±)	0.41	0.34	0.36	0.31	0.33	0.41
CD (P=0.05)	NS	NS	NS	NS	NS	NS
<b>Nitrogen levels (N)</b>						
N <sub>0</sub>	7.74	8.45	8.23	7.65	8.37	8.06
N <sub>1</sub>	7.96	8.56	8.47	7.96	8.59	8.55
N <sub>2</sub>	8.15	8.89	8.75	8.20	8.94	8.89
N <sub>3</sub>	8.52	9.17	9.20	8.59	9.30	9.37
S. Ed (±)	0.24	0.21	0.22	0.17	0.27	0.25
CD (P=0.05)	0.56	0.47	0.52	0.40	0.63	0.59
<b>Interaction (I×N)</b>						
S.Ed (±)	0.54	0.46	0.50	0.38	0.61	0.57
CD (P=0.05)	NS	NS	NS	NS	NS	NS

N.S: Non-significant

**Comment [SN12]:** As you mentioned in paragraph that nitrogen significantly effect crude protein contents so add lettering with values


### Crude protein yield

The effect of irrigation regimes on crude protein yield was found to be significant in ryegrass at 1<sup>st</sup> cut, 2<sup>nd</sup> cut and 3<sup>rd</sup> cut, during both years (Table 2). The application irrigation at IW:CPE ratio of 1.4 recorded the highest crude protein yield (7.69 q/ha and 7.31 q/ha, respectively in both the years) which was significantly higher than other irrigation regimes. The lowest crude protein yield was found under rainfed treatment. The crude protein yield is the function of crude protein content and dry matter yield. The higher dry matter yield was found with application of irrigation at IW:CPE ratio of 1.4. So, the highest crude protein yield were recorded under this treatment. Similar findings were reported by Agrawal et al. [17] and Jat et al. [1] in case of fodder oat. The crude protein yield as influenced by different nitrogen levels was found to be significant in ryegrass at 1<sup>st</sup> cut, 2<sup>nd</sup> cut and 3<sup>rd</sup> cut, during both the years are presented in Table 2. Application of 90 kg N/ha recorded the highest crude protein yield (6.54 q/ha and 6.64 q/ha, respectively in both the years) followed by 60 kg N/ha and lowest was recorded in 0 kg N/ha. Highest dry matter yield was found with application of 90 kg N/ha which ultimately resulted in higher crude protein yield. These results are in conformity with the findings of Gasim [8] and Sarkar et al. [18].


**Table 2. Effect of irrigation regimes (I) and nitrogen levels (N) on crude protein yield (q/ha) of ryegrass**

Treatments	Crude protein yield (q/ha)							
	1 <sup>st</sup> Year				2 <sup>nd</sup> Year			
	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	Total	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	Total
<b>Irrigation regimes (I)</b>								
I <sub>0</sub>	0.70	1.03	0.89	2.62	0.79	1.06	0.91	2.75
I <sub>1</sub>	1.13	1.43	1.25	3.96	1.17	1.46	1.32	3.95
I <sub>2</sub>	1.29	1.70	1.51	4.50	1.29	1.72	1.53	4.54
I <sub>3</sub>	1.58	2.07	1.88	5.52	1.68	2.12	1.91	5.72
I <sub>4</sub>	2.19	2.83	2.66	7.69	2.14	2.67	2.51	7.31
S.Ed (±)	0.17	0.094	0.18	0.62	0.095	0.16	0.15	0.35
CD (P=0.05)	0.39	0.22	0.42	1.44	0.22	0.37	0.35	0.81
<b>Nitrogen levels (N)</b>								
N <sub>0</sub>	0.68	1.09	0.95	2.83	0.79	1.05	0.92	2.76
N <sub>1</sub>	1.35	1.74	1.53	4.62	1.34	1.74	1.51	4.59
N <sub>2</sub>	1.56	2.02	1.86	5.44	1.58	2.00	1.84	5.43
N <sub>3</sub>	1.92	2.41	2.22	6.54	1.94	2.43	2.27	6.64

S. Ed ( $\pm$ )	0.099	0.091	0.094	0.26	0.076	0.087	0.091	0.24
CD (P=0.05)	0.23	0.21	0.22	0.60	0.18	0.20	0.21	0.55
<b>Interaction (I×N)</b>								
S.Ed ( $\pm$ )	0.22	0.20	0.21	0.58	0.17	0.19	0.20	0.53
CD (P=0.05)	0.45	0.41	0.43	1.18	0.35	0.39	0.42	1.08
CV (%)	19.61	13.67	15.69	14.58	14.80	13.13	15.30	13.36



**Fig. 1: Indicating the effect of irrigation regimes and nitrogen levels on crude protein yield (q/ha) of ryegrass (1<sup>st</sup> year)**



**Fig. 2: Indicating the effect of irrigation regimes and nitrogen levels on crude protein yield (q/ha) of ryegrass (2<sup>nd</sup> year)**

**Interaction effect**

The interaction effect between irrigation regimes and nitrogen levels was found to be significant in respect of crude protein yield of ryegrass at 1<sup>st</sup> cut, 2<sup>nd</sup> cut and 3<sup>rd</sup> cut during both the years are presented in Table 3, Table 4, Table 5, Table 6, Table 7, Table 8, Table 9 and Table 10. The higher crude protein yield was recorded with the application of irrigation at IW:CPE ratio of 1.4 in combination with 90 kg N/ha at 1<sup>st</sup> cut, 2<sup>nd</sup> cut and 3<sup>rd</sup> cut during both the years followed by irrigation at IW:CPE ratio of 1.2 in combination with 60 kg N/ha. Rainfed treatment in combination with 0 kg N/ha recorded the lowest crude protein yield in all three cuts during both the years due to lowest dry matter accumulation in the respective treatments.

Comment [SN13]: Rephrase this paragraph

**Table 3. Interaction effect of irrigation regimes (I) × nitrogen levels (N) on crude protein yield (q/ha) of ryegrass at 1<sup>st</sup> cut (1<sup>st</sup> year)**

Nitrogen levels (N)	Irrigation regimes (I)					Mean
	I <sub>0</sub>	I <sub>1</sub>	I <sub>2</sub>	I <sub>3</sub>	I <sub>4</sub>	
N <sub>0</sub>	0.39	0.54	0.61	0.64	1.21	0.68
N <sub>1</sub>	0.68	1.05	1.25	1.50	2.27	1.35
N <sub>2</sub>	0.93	1.22	1.37	1.83	2.44	1.56
N <sub>3</sub>	0.78	1.69	1.94	2.33	2.83	1.92
<b>Mean</b>	0.70	1.13	1.29	1.58	2.19	1.38
		<b>I</b>		<b>N</b>		<b>I×N</b>
S.Ed (±)		0.17		0.099		0.22
CD (P=0.05)		0.39		0.23		0.45

**Table 4. Interaction effect of irrigation regimes (I) × nitrogen levels (N) on crude protein yield (q/ha) of ryegrass at 2<sup>nd</sup> cut (1<sup>st</sup> year)**

Nitrogen levels (N)	Irrigation regimes (I)					Mean
	I <sub>0</sub>	I <sub>1</sub>	I <sub>2</sub>	I <sub>3</sub>	I <sub>4</sub>	
N <sub>0</sub>	0.66	0.83	1.03	1.06	1.86	1.09
N <sub>1</sub>	0.91	1.36	1.74	2.05	2.63	1.74
N <sub>2</sub>	1.21	1.59	1.74	2.36	3.21	2.02
N <sub>3</sub>	1.36	1.95	2.27	2.81	3.63	2.41
<b>Mean</b>	1.03	1.43	1.70	2.07	2.83	1.81
		<b>I</b>		<b>N</b>		<b>I×N</b>
S.Ed (±)		0.094		0.091		0.20
CD		0.22		0.21		0.41

(P=0.05)						
----------	--	--	--	--	--	--

**Table 5. Interaction effect of irrigation regimes (I) × nitrogen levels (N) on crude protein yield (q/ha) of ryegrass at 3<sup>rd</sup> cut (1<sup>st</sup> year)**

Nitrogen levels (N)	Irrigation regimes (I)					Mean
	I <sub>0</sub>	I <sub>1</sub>	I <sub>2</sub>	I <sub>3</sub>	I <sub>4</sub>	
N <sub>0</sub>	0.51	0.70	0.92	0.91	1.72	0.95
N <sub>1</sub>	0.78	1.25	1.39	1.82	2.40	1.53
N <sub>2</sub>	1.11	1.41	1.60	2.12	3.06	1.86
N <sub>3</sub>	1.16	1.66	2.12	2.66	3.48	2.22
<b>Mean</b>	0.89	1.25	1.51	1.88	2.66	1.64
		<b>I</b>		<b>N</b>		<b>I×N</b>
S.Ed (±)		0.18		0.094		0.21
CD (P=0.05)		0.42		0.22		0.43

**Table 6. Interaction effect of irrigation regimes (I) × nitrogen levels (N) on total crude protein yield (q/ha) of ryegrass (1<sup>st</sup> year)**

Nitrogen levels (N)	Irrigation regimes (I)					Mean
	I <sub>0</sub>	I <sub>1</sub>	I <sub>2</sub>	I <sub>3</sub>	I <sub>4</sub>	
N <sub>0</sub>	1.56	2.64	2.56	2.61	4.79	2.83
N <sub>1</sub>	2.37	3.65	4.38	5.38	7.30	4.62
N <sub>2</sub>	3.25	4.22	4.71	6.31	8.71	5.44
N <sub>3</sub>	3.30	5.30	6.33	7.80	9.95	6.54
<b>Mean</b>	2.62	3.96	4.50	5.52	7.69	4.86
		<b>I</b>		<b>N</b>		<b>I×N</b>
S.Ed (±)		0.62		0.26		0.58
CD (P=0.05)		1.44		0.60		1.18

**Table 7. Interaction effect of irrigation regimes (I) × nitrogen levels (N) on crude protein yield (q/ha) of ryegrass at 1<sup>st</sup> cut (2<sup>nd</sup> year)**

Nitrogen levels (N)	Irrigation regimes (I)					Mean
	I <sub>0</sub>	I <sub>1</sub>	I <sub>2</sub>	I <sub>3</sub>	I <sub>4</sub>	
N <sub>0</sub>	0.46	0.66	0.75	0.80	1.30	0.79
N <sub>1</sub>	0.69	1.07	1.29	1.58	2.10	1.34
N <sub>2</sub>	0.94	1.32	1.38	1.97	2.31	1.58
N <sub>3</sub>	1.06	1.64	1.74	2.38	2.85	1.94
<b>Mean</b>	0.79	1.17	1.29	1.68	2.14	1.41
		<b>I</b>		<b>N</b>		<b>I×N</b>
S.Ed (±)		0.095		0.076		0.17
CD (P=0.05)		0.22		0.18		0.35

**Table 8. Interaction effect of irrigation regimes (I) × nitrogen levels (N) on crude protein yield (q/ha) of ryegrass at 2<sup>nd</sup> cut (2<sup>nd</sup> year)**

Nitrogen levels (N)	Irrigation regimes (I)					Mean
	I <sub>0</sub>	I <sub>1</sub>	I <sub>2</sub>	I <sub>3</sub>	I <sub>4</sub>	
N <sub>0</sub>	0.68	0.85	0.99	1.08	1.63	1.05
N <sub>1</sub>	0.95	1.43	1.71	2.08	2.53	1.74
N <sub>2</sub>	1.23	1.60	1.82	2.39	2.99	2.00
N <sub>3</sub>	1.36	1.96	2.38	2.94	3.51	2.43
<b>Mean</b>	1.06	1.46	1.72	2.12	2.67	1.81
		<b>I</b>		<b>N</b>		<b>I×N</b>
S.Ed (±)		0.16		0.087		0.19
CD (P=0.05)		0.37		0.20		0.39

Comment [SN14]: Insert lettering if data of main factor or interaction was significant

**Table 9. Interaction effect of irrigation regimes (I) × nitrogen levels (N) on crude protein yield (q/ha) of ryegrass at 3<sup>rd</sup> cut (2<sup>nd</sup> year)**

Nitrogen levels (N)	Irrigation regimes (I)					Mean
	I <sub>0</sub>	I <sub>1</sub>	I <sub>2</sub>	I <sub>3</sub>	I <sub>4</sub>	
N <sub>0</sub>	0.51	0.73	0.95	0.95	1.47	0.92

<b>N<sub>1</sub></b>	0.78	1.30	1.39	1.83	2.23	1.51
<b>N<sub>2</sub></b>	1.12	1.45	1.64	2.17	2.84	1.84
<b>N<sub>3</sub></b>	1.21	1.81	2.14	2.70	3.49	2.27
<b>Mean</b>	0.91	1.32	1.53	1.91	2.51	1.64
		<b>I</b>		<b>N</b>		<b>I×N</b>
S.Ed (±)		0.15		0.091		0.20
CD (P=0.05)		0.35		0.21		0.42

**Table 10. Interaction effect of irrigation regimes (I) × nitrogen levels (N) on total crude protein yield (q/ha) of ryegrass (2<sup>nd</sup> year)**

Nitrogen levels (N)	Irrigation regimes (I)					Mean
	I <sub>0</sub>	I <sub>1</sub>	I <sub>2</sub>	I <sub>3</sub>	I <sub>4</sub>	
<b>N<sub>0</sub></b>	1.65	2.23	2.69	2.83	4.40	2.76
<b>N<sub>1</sub></b>	2.42	3.81	4.38	5.49	6.85	4.59
<b>N<sub>2</sub></b>	3.29	4.37	4.83	6.53	8.14	5.43
<b>N<sub>3</sub></b>	3.64	5.41	6.26	8.02	9.86	6.64
<b>Mean</b>	2.75	3.95	4.54	5.72	7.31	4.86
		<b>I</b>		<b>N</b>		<b>I×N</b>
S.Ed (±)		0.35		0.24		0.53
CD (P=0.05)		0.81		0.55		1.08

### Evapotranspiration, Water use and water use efficiency

Data on evapotranspiration, total water use, crop water use efficiency and field water use efficiency as influenced by different irrigation regimes are presented in Table 11. Increasing levels of irrigation regime increased the evapotranspiration and total water use. Application of irrigation at IW:CPE ratio of 1.4 recorded the highest value in terms of evapotranspiration (26.33 cm and 24.74 cm, respectively in both the years), total water use (26.65 cm and 25.16 cm, respectively in both the years), crop water use efficiency (324.49 kg/ha-cm and 324.12 kg/ha-cm, respectively in both the years) and field water use efficiency (322.69 kg/ha-cm and 318.84 kg/ha-cm, respectively in both the years) followed by irrigation at IW:CPE ratio of

**Comment [SN15]:** Mentioned the method of evapotranspiration measurement in materials and methods chapter

1.2 and lowest values were recorded in rainfed treatment. The evapotranspiration and total water use increased with increasing level of irrigation regimes due to more numbers of irrigation applied which facilitated higher availability water in the soil profile leading to more loss of water through evapotranspiration. The crop and field water use efficiency also increased under IW:CPE ratio of 1.4 because of higher yield. Data pertaining to the evapotranspiration, total water use, crop water use efficiency and field water use efficiency as influenced by different nitrogen levels are presented in Table 11. The highest values in terms of evapotranspiration (21.16 cm and 18.89 cm, respectively in both the years), total water use (21.16 cm and 19.05 cm, respectively in both the years), crop water use efficiency (343.19 kg/ha-cm and 376.87 kg/ha-cm, respectively in both the years) and field water use efficiency (344.33 kg/ha-cm and 372.88 kg/ha-cm, respectively in both the years) were observed under 90 kg N/ha. The higher evapotranspiration and total water use observed due to higher crop growth rate and yield of the crop which exhibited more extraction of soil moisture. Higher crop and field water use efficiency recorded because higher yield in the respective treatments in both the years.

**Comment [SN16]:** Improve write-up and also mentioned significant or non- significant effect on various parameters

**Table 11. Effect of irrigation regimes (I) and nitrogen levels (N) on water use and water use efficiency of ryegrass**

Treatments	1 <sup>st</sup> year				2 <sup>nd</sup> year			
	Evapotranspiratin (cm)	Total water use (cm)	Crop water use efficiency(kg/ha-cm)	Field water use efficiency (kg/ha-cm)	Evapotranspiratin (cm)	Total water use (cm)	Crop water use efficiency(kg/ha-cm)	Field water use efficiency (kg/ha-cm)
<b>Irrigation levels (I)</b>								
I <sub>0</sub>	14.12	14.12	234.69	234.69	12.32	12.32	277.18	277.18
I <sub>1</sub>	17.15	17.24	271.03	270.84	15.46	16.12	309.76	299.40
I <sub>2</sub>	21.13	22.31	245.90	234.54	18.61	18.72	278.96	277.79
I <sub>3</sub>	23.43	24.81	260.40	253.08	20.17	20.64	317.33	310.64
I <sub>4</sub>	26.33	26.65	324.49	322.69	24.74	25.16	324.12	318.84
<b>Nitrogen levels (N)</b>								
N <sub>0</sub>	18.96	20.54	173.68	160.37	18.12	18.42	189.65	186.02
N <sub>1</sub>	20.21	21.08	266.03	258.72	17.65	18.15	302.55	294.82
N <sub>2</sub>	21.40	21.33	286.31	289.24	18.37	18.76	336.79	333.35
N <sub>3</sub>	21.16	21.16	343.19	344.33	18.89	19.05	376.87	372.88

### Economics of cultivation of ryegrass

Data presented in Table 12 indicated that in spite of increase in cost of cultivation, the gross return, net return and B:C ratio increased with the increasing irrigation regime from IW:CPE ratio of 1.0 to IW:CPE ratio of 1.4. The highest gross return (Rs. 78073.00 and Rs. 78254.00, respectively in both the years), net return (Rs. 51252.00 and Rs. 52433.00, respectively in both the years) and B:C ratio (1.87 and 1.99 respectively, in both the years) was obtained in Irrigation at IW:CPE ratio of 1.4 and followed by Irrigation at IW:CPE ratio of 1.2 in terms of gross return, net return and B:C ratio. The lowest gross return, net return and B:C ratio was observed under rainfed treatment in both the years. The highest cost of cultivation in irrigation at IW:CPE ratio of 1.4 due to higher cost of irrigation but increase yield led to higher gross return, net return and B: C ratio in both the years. Data presented in Table 12 showed that the higher gross return, net return and B:C ratio was obtained in 90 kg N/ha treatment over the other nitrogen levels. The cost of cultivation of ryegrass was observed to be comparatively higher with application of 90 kg N/ha. Yet, the higher gross return (Rs. 72376.00 and Rs. 70383.00, respectively in both the years), net return (Rs. 45038.00 and Rs. 43645.00, respectively in both the years) and B:C ratio (1.63 and 1.62 respectively in both the years) was obtained in 90 kg N/ha treatment followed by 60 kg N/ha. The highest cost of cultivation recorded with application of 90 kg N/ha because higher amount of fertilizer used compared to other nitrogen levels but higher gross return, net return and B:C ratio were obtained due to higher yield.

**Table 12. Economics of ryegrass as affected by irrigation regimes (I) and nitrogen levels (N)**

Treatments	2017-18				2018-19			
	Cost (Rs.)	Gross return (Rs.)	Net return (Rs.)	B:C ratio	Cost (Rs.)	Gross return (Rs.)	Net return (Rs.)	B:C ratio
<b>Irrigation regimes (I)</b>								
I <sub>0</sub>	24321	36195	12275	0.48	24321	36267	12386	0.49
I <sub>1</sub>	25321	48022	23516	0.90	25071	49711	24640	0.96
I <sub>2</sub>	25821	53384	27146	1.02	25321	52224	26903	1.04

I <sub>3</sub>	25821	63244	37418	1.41	25321	63659	38338	1.48
I <sub>4</sub>	26821	78073	51252	1.87	25821	78254	52433	1.99
<b>Nitrogen levels (N)</b>								
N <sub>0</sub>	22420	32505	11058	0.48	22020	34824	13156	0.59
N <sub>1</sub>	26038	55586	29215	1.11	25638	56023	30385	1.17
N <sub>2</sub>	26688	62667	35975	1.33	26288	62862	36574	1.38
N <sub>3</sub>	27338	72376	45038	1.63	26738	70383	43645	1.62

#### 4. CONCLUSION

From the ongoing discussion it can be concluded that application of irrigation at IW:CPE ratio of 1.4 recorded the significantly higher crude protein yield as compared to other irrigation treatments. The highest value in terms of evapotranspiration, total water use, crop water use efficiency and field water use efficiency recorded under this treatment. The highest gross return, net return and B:C ratio obtained in irrigation at IW:CPE ratio of 1.4 over the rest of the treatments. The crude protein content of ryegrass was significantly influenced by different nitrogen levels. The highest values of crude protein content was observed under 90 kg N/ha. Application of 90 kg N/ha recorded significantly higher crude protein yield over rest of the treatments. The higher values in terms of evapotranspiration, total water use, crop water use efficiency and field water use efficiency observed with application of 90 kg N/ha. Economic analysis showed that the higher gross return, net return and B:C ratio were recorded in 90 kg N/ha. Among the different treatment combination of irrigation regimes and nitrogen levels, the highest crude protein yield was recorded with application of irrigation at IW:CPE ratio of 1.4 in combination with 90 kg N/ha.

**Comment [SN17]:** Repeated same as mentioned in previous line

**Comment [SN18]:** Describe in single line that these these parameters perform highest under 90 kg N/ha not separately each parameters

**Comment [SN19]:** Conclusion is lengthy conclude it with in 5-7 lines

#### REFERENCES

1. Jat H, Kaushik MK, Nepalia V, Singh D. Effect of irrigation schedule and nitrogen fertilization on growth, yield and quality of fodder oat (*Avena sativa* L.). Journal of Pharmacognosy and Phytochemistry. 2017;6(4): 2040-2042.
2. Tahir GM, Azraf-ul-Haq, Khaliq T, Rehman M, Hussain S. Effect of Different Irrigation Levels on Yield and Forage Quality of Oat (*Avena sativa*). Applied Science Reports. 2014; 3(1): 42-46.
3. Singh SD. Effect of irrigation regimes and nitrogen levels on growth, yield and quality of baby corn. Madras Agricultural Journal. 2001;88(7/9): 367-370.
4. Hussaini MA, Ogunlela VB, Ramalan AA, Falaki AM. Growth and development of maize (*Zea mays* L.) in response to different levels of nitrogen, phosphorus and irrigation. Crop Research. 2001;22(2): 141-149.

**Comment [SN20]:** Tally the references with citation and it should be according to the journal style

5. Jiotode DJ, Lambe DL, Dhawad CS. Growth parameters and water use studies of maize as influenced by irrigation levels and row spacing. *Crop Research*.2002;24(2): 292-295.
6. Patel MR. Effect of irrigation, nitrogen and biofertilizer on yield and quality of forage oat (*Avena sativa* L.). M.Sc. (Ag.) Thesis. Anand Agricultural University, Gujrat.2006.
7. Bhilare RL, Joshi YP. Productivity and quality of oat (*Avena sativa*) in relation to cutting management and nitrogen levels. *Indian Journal of Agronomy*.2007;52(3): 247-250.
8. Gasim S A. Effect of nitrogen, phosphorus and seed rate on growth, yield and quality of forage maize (*Zea mays*). M.Sc. Thesis, Faculty of Agriculture, University of Khartoum.2001.
9. Adam MY. Effect of seed rate and nitrogen on growth and yield of teff grass (*Eragrostis tef* Zucc.). Trotter. B. Sc. (Agric.), a dissertation submitted to the University of Khartoum in partial fulfilment for the requirements of the degree of Master of Science (Agri.).2000.
10. Dubey A, Rathi GS, Sahu R. Effect of nitrogen levels on green fodder yield of oat (*Avena sativa* L.) varieties. *Forage Research*.2013; 39(1): 39-41.
11. Jehangir IA, Khan HU, Khan MH, F. Ur-Rasool, Bhat RA, Mubarak T, Bhat MA, Rasool S. Effect of sowing dates, fertility levels and cutting managements on growth, yield and quality of oats (*Avena sativa* L.). *African Journal of Agricultural Research*.2013; 8(7): 648-651.
12. Mahdi S, Hasan B, Singh L. Influence of seed rate, nitrogen and zinc on fodder maize (*Zea mays*) in temperate conditions of western Himalayas. *Indian Journal of Agronomy*.2012;57: 85-88.
13. Shehzad MA, Maqsood M, Bhatti MA, Ahmad W, Shahid MR. Effect of nitrogen fertilization rate and harvest time on maize (*Zea mays* L) fodder yield and its quality attributes. *Asian Journal of Pharmaceutical and Biological Research*.2012;2(1): 19-26.
14. Dadarwal RS, Jain NK, Singh D. Integrated nutrient management in baby corn (*Zea mays*). *Indian Journal of Agricultural Sciences*.2009;79(12): 1023-1025.
15. Luikham E, Kamei S, Mariam, Anal PS. Yield, quality and economics of oat fodder (*Avena sativa* L.) as influenced by nitrogen and varieties. *Forage Research*.2012;38: 112-114.
16. Bindhani A, Barik KC, Garnayak LM, Mahapatra PK. Nitrogen management in baby corn (*Zea mays* L.). *Indian Journal of Agronomy*.2007;52: 135-138.
17. Agrawal RK, Varma SC, Singh KK. Influence of nitrogen and moisture regimes on growth and yield of oats cultivars. *Range Management and Agroforestry*.2000;21: 206-212.
18. Sarkar RK, Mahasin M. Productivity, quality and profitability of oat based intercropping system under graded fertility levels. *Indian Journal of Agricultural Sciences*.2007;77(12):862-865

UNDER PEER REVIEW

