

Evaluation of Genotoxicity by Comet Assay in Tissues of *Clarias gariepinus* Exposed to Cassava Effluent

ABSTRACT

Untreated wastewater discharge into waterbodies poses a number of risks, including the potential for aquatic organisms' DNA to be damaged. Using the comet assay, this study assessed the genotoxic impact of cassava wastewater on the gonads, liver, and gills of post-juvenile *Clarias gariepinus*. The post-juvenile *C. gariepinus* fish were acquired from a fish farm in Edo State, Nigeria, and were subjected to different concentrations of cassava wastewater (0.2%, 0.3%, 0.5%, and 0.7%) for a duration of 96 hours. In contrast to the typical actions seen in the control groups, the fish exposed to the effluent exhibited restless, erratic movements, and gasping for air. The first set of repetitions showed that catfish mortality increased with concentration; after 96 hours, exposure to the lowest concentration (0.2%) and the highest concentration (0.7%) of cassava wastewater caused 10% and 40% of the fish to die, respectively. The second set of repetitions showed that catfish mortality increased with increasing concentration; after 96 hours, exposure to the lowest concentration (0.2%) and the highest concentration (0.7%) of cassava wastewater resulted in 20% and 50% catfish mortality, respectively. Following the exposure time, the fish were brought to the lab where their gonads, liver, and gills were removed in order to use the comet assay for genotoxic assessment. The genotoxins present in the wastewater can be the reason for the DNA impairment found in the genotoxic assessment result at different effluent concentrations. The investigation also showed that DNA damage increased with increasing concentration, indicating a dose-dependent genotoxic effect of cassava wastewater on post-juveniles of *C. gariepinus*. This research demonstrates the critical importance of treating wastewater before to discharge.

Keywords: Genotoxicity, Comet Assay, *Clarias gariepinus*, Cassava Effluent

INTRODUCTION

The word "genotoxicity" in genetics describes the existence of chemicals that can damage a cell's DNA and RNA, thereby harming the integrity of the cell. Mutagens that cause genotoxicity, such as radiation and chemicals, can cause damage to DNA or chromosomes and consequent mutations. Damage to embryonic cells can result in inherited abnormalities that cause birth defects, while damage to somatic cells' genetic material may cause cancer in eukaryotic creatures (Mohamed et al., 2017).

Researchers examine DNA impairment in cells exposed to these dangerous substances so as to determine the genotoxicity of various substances. Single- or double-strand breaks, point mutations, loss of excision repair, cross-linking, and structural or numerical anomalies in chromosomes are examples of this type of DNA damage. It is acknowledged that genetic material compromise has a role in the development of malignant diseases marked by unchecked cell expansion. This result has led to the development of a variety of complex techniques, including in vitro and in vivo toxicological examinations, the Ames Test, and the Comet Assay, to evaluate a substance's potential to instigate damage to genetic material that could ultimately lead to the advancement of cancer (Nagarathna et al., 2013).

One major global concern is the possibility of genotoxic chemical contamination of water. Mutagens in drinking water can have a negative impact on people's health, possibly leading to the development of cancer. Water ecosystems can also be harmed by environmental

contamination (Efekemo et al., 2024). Therefore, a great deal of study has been done to assess the mutagenic potential of drinking water as well as surface and groundwater sources. This research has revealed a wide range of materials that have the prowess to harm DNA (Ferk et al., 2009).

Some refer to cassava (*Manihot esculenta*) as the "bread of the tropics," the "food of the poor," or even the "poverty fighter." It is a staple crop that is grown on a local scale by farmers in Nigeria and other tropical African countries. Because of its drought resistance and versatility in a range of soil types, including marginal ones, cassava is an important source of carbohydrates in tropical Africa. Numerous high-calorie foods, including garri (toasted granules), fufu (fermented cooked paste), starch, tapioca, and other confections, can be made from cassava tubers (Oghenejoboh et al., 2021; Tonukari et al., 2016). On the other hand, processing cassava generates a significant amount of liquid and solid waste. Leaves, stems, peels, effluent, and starch bagasse are some of these byproducts. Despite the obvious harm these methods inflict on the environment and public health, Nigerian farmers frequently dispose of these wastes carelessly in open spaces near major roads and streets, undeveloped land plots, unfinished buildings, and water bodies (Oghenejoboh et al., 2021; Orororo et al., 2014). About 60–70% of raw cassava tubers are made up of water, much of which is converted into wastewater during processing. Furthermore, a significant amount of water is used in the processing of cassava roots into food items like starch and garri, which is subsequently released into the environment. Remarkably, depending on the intended end product, processing one ton of cassava tubers can produce nothing less than 250kg of wastewater (Oghenejoboh et al., 2021).

Due to its enormous cyanide content and significant chemical oxygen demand (COD), cassava wastewater poses significant environmental risks and if left untreated, can contaminate soil, rivers, and groundwater. Cassava is naturally high in cyanogenic glycosides, which can hydrolyze into hydrogen cyanide during processing, especially during pressing and washing. This release of cyanide into the environment, in the form of hydrocyanic acid, poses a significant environmental hazard (Fawole et al., 2014). High concentrations of cassava effluent negatively affect the physicochemical parameters of water, which in turn affect aquatic life. Using cassava effluent-contaminated water for irrigation can stunt plant growth or even cause plant death, depending on the concentration of cyanide.

The effluents from sewage treatment plants, industrial wastewater, and agricultural runoff are the main sources of surface water contamination that contain different genotoxic chemicals. A small number of harmful substances can not only be analyzed for surface water monitoring to be successful, given the diversity of chemical contaminants present. Although very sensitive, genotoxicity studies that rely on the metabolic or enzymatic activities of test organisms are not very specific. Therefore, in order to reliably determine the effects of genotoxic chemicals on a wide variety of organisms, a combination of biotests involving eukaryotic as well as prokaryotic organisms (fish, mussels, *Daphnia*, bacteria and algae) becomes required (Lah et al., 2004).

One method for quantifying DNA strand breakage at the individual cell level is the comet assay. Known also as single-cell gel electrophoresis (SCGE), this method is frequently used to evaluate the deleterious effects of toxins on genetic material. Because of its wide variety of uses, simplicity, sensitivity, adaptability, and time economy, this method is widely recognized for evaluating DNA damage. It is extensively employed in environmental monitoring to evaluate the effects of different toxicants on aquatic life. Environmental research has made extensive use of this assay, especially for assessing the genotoxicity of pollutants and toxins in aquatic environments. It makes it possible for scientists to assess how environmental stressors impact aquatic animals' DNA integrity (De Lapuente et al., 2015). To summarize, the comet assay involves applying single-cell suspensions in agarose to microscope slides,

treating them with detergent and concentrated NaCl to break down cell membranes and extract histones, and then electrophoresizing the results. When DNA strands break and are observed under fluorescence microscopy, they travel toward the anode and resemble comets when dyed fluorescently (Cordelli, Bignami, and Pacchierotti, 2021).

The African catfish, or *Clarias gariepinus*, is a popular freshwater fish farmed in Nigeria because of its quick growth and tolerance to a variety of environments. For toxicological research, it has been widely utilized as a model (Mahboub et al., 2021; Basari et al., 2024). African catfish are an significant source of protein, and their health a good measure of the quality of their habitat. Because African catfish respond well to ecological contamination, many researchers have selected it as a bio-indicator for genotoxicity and oxidative stress (Osman et al., 2012; Arojojoye et al., 2018; Zaghoul et al., 2020).

Fish are important members of aquatic ecosystems since they can react to low levels of hazardous compounds in the water directly and indirectly through eating polluted aquatic creatures. Hence, according to Cavas and ErgeneGözükara (2005), fish can serve as markers for the early detection of contamination in the maritime environment. A primary factor contributing to water contamination is the careless disposal of effluents into bodies of water. Given the economic significance of fish to humans and the high toxicity of unprocessed cassava products and effluents (Egbune et al., 2023), more research evaluating the toxic effects of cassava waste water on fish is required, even though some studies have been carried out on the genotoxic effect of cassava wastewater on the African Catfish (Ariyomo et al., 2017; Olufayo et al., 2016; Olorunfemi et al., 2019; Olorunfemi et al., 2024). Additionally, there aren't many research using the comet assay to explain the genotoxic effects of cassava effluent on *Clarias gariepinus*. Therefore, this study evaluated the genotoxic effects of cassava effluent on selected tissues of *C. gariepinus* using the comet assay.

MATERIALS AND METHODS

Study Area

This study was carried out in Iguedo, Usen district of Ovia South-West Local Government Area, Edo State, Nigeria. Iguedo is situated at Ovia Local Government Area at Latitude 6.72306 East and Longitude 5.38609 North (Figure 1). The effluent was collected from a cassava processing industry at the point of discharge.




Fig 1: Diagram illustrating the study's geographic region

Test animal: collection and acclimatization

One hundred post-juvenile *Clarias gariepinus* fish were acquired from Vinvok farms in Orerokpe, Delta State, Nigeria. They were then transported to the Wet Laboratory, Department of Fisheries, Faculty of Agriculture, University of Benin, Benin City, Edo state, Nigeria, under well-ventilated circumstances. Before the test, the fish were given five days to get used to the lab environment. The fish were fed commercial fish meal twice a day during the adaption period, and the water in the tanks where they were housed was changed daily. The fish were exposed to the cassava effluent after the adaptation phase.

Treatment Fish Samples and Acute toxicity tests

After a series of tests, concentrations were chosen. Concentrations were selected following an array of experiments. For 96 hours, ten fish were exposed to every concentration. The fish were subjected to tests on five concentrations that were prepared by serial dilution: 0.2%, 0.3%, 0.5%, 0.7%, and control (distilled water). In order to prevent contamination of the test environment, the number of dead fish was counted at the commencement of the tests and then every 24 hours after that. The fish were then removed and disposed of appropriately (Sogbanmu et al., 2018). When fish showed no indications of life and their carcasses were either sinking to the bottom of the test solution or floating horizontally close to the surface, they were declared dead. To prevent repeating the test in the event that the first setup failed, a replicate per test concentration was employed, which also served to strengthen the statistical baseline.

COMET ASSAY PROTOCOL

Extraction of organs and Slide Preparation: The fish samples are sacrificed and their gills, liver and gonads were excised and placed in separate well labelled eppendorf tubes. Comet assay slides are sterilized and allowed to dry. The slides are then labelled using a diamond pen and were then embedded in 0.5% low melting point agarose gel and thereafter placed on the slides and covered with cover slips. The coverslip was gently removed after a while and 1% low melting point agarose was added. The slides were then dried and kept on a tray.

Lysing: The cells are lysed to release DNA. This was done using a solution containing sodium hydroxide, EDTA, Triton-X and DMSO for 24 hours. Sodium hydroxide was used for the osmotic pressure for the destruction of the cell membrane. EDTA chelates co-factors which are usually divalent metals that affect nuclear activities. DMSO helps in osmotic pressure. Triton-X acts as a detergent and breaks the phospholipid bilayer of animal cells.

Electrophoresis: The DNA samples were then subjected to electrophoresis, which caused the DNA to move through the agarose gel. It involved the use of an electrophoresis buffer that contains sodium hydroxide and EDTA. This allowed the DNA to separate into fragments. This was done by placing the slides into electrophoresis tank and passing current of 300 amperes at 25V through it for thirty minutes. The process was conducted in the dark to avoid light-induced damage to the DNA samples.

Neutralization and Fixation: The slides were neutralized with Tris to keep the pH at 7.4. Neutralization was done three times every five minutes to stop the electrophoresis process. The slides were then fixed with cold ethanol and allowed to dry. Application of cold ethanol dehydrated any gel left on the slide and prevented contamination of the slide.

Staining: The slides were then stained with Giemsa stain and rinsed with distilled water. This allowed the DNA to be seen under a microscope.

Viewing and Scoring: The DNA fragments were then imaged with the aid of a light microscope fitted with a camera. The visual representations were thereafter analysed to determine the amount of DNA damage in the sample. A comet comprises a head and a tail, with the tail indicating the extent of DNA damage.

Analysis: Using image analysis techniques on individual cells is the most versatile way to gather comet data, and there are a number of commercially available specialist software packages for this purpose. Individual comet photos were examined for a number of parameters using Graph Pad Prism A, such as DNA content (overall intensity), head and tail length, DNA percentage in tail, and tail moment (fraction of migrating DNA multiplied by a tail length measurement). The most often reported measurements of these are tail moment and/or tail length; the usage of percent DNA in the tail provides a clear indicator of the comet appearance and is also linearly related to the frequency of DNA breaks over a broad range of damage levels. Therefore, the comet tail moment parameter (TM) was employed in this investigation to ascertain the extent of DNA damage. It was computed by multiplying the comet tail length by the DNA proportion. The tail moment is the sum of the comet tail length, which represents the smallest observable size of moving DNA, and the tail DNA intensity, which represents the quantity of relaxed/broken bits (Patek et al., 2018).

RESULTS

Acute toxicity mortality rate of *Clarias gariepinus* to cassava effluent exposure

Table 1 displays the mortality rates of *C. gariepinus* when exposed to cassava wastewater. In the first set of replicates, exposure to the lowest concentration (0.2%) and the highest concentration (0.7%) of cassava wastewater resulted in 10% and 40% catfish mortality, respectively, after 96 hours, demonstrating an increase in mortality as the concentration increased. In the second set of replicates, exposure to the lowest concentration (0.2%) and the highest concentration (0.7%) of cassava wastewater led to 20% and 50% catfish mortality, respectively, after 96 hours, also indicating a rise in mortality with increasing concentration. Behavioural responses were observed to be dose dependent, and also showed direct proportion with increased concentrations. As concentrations of the test effluent increased, the fish samples displayed initial disturbed swimming movements. This was accompanied with avoidance response manifested via surfacing behaviour. This was followed by sudden quick movement, unusual lethargy, and bleaching of the skin. It was also observed that the fish samples settled at the bottom of the aquaria. They were motionless and had slow opercula movements as they gulped for air. Fishes exposed to 100% concentration of cassava wastewater died in three minutes. Fishes in control (0% concentration) showed normal behaviour.

Table 1: Mortality rate of *Clarias gariepinus* exposed to varied concentrations of cassava wastewater

Concentration (%)	Duplicate mortality hour 1				Duplicate mortality hour 2			
	24	48	72	96	24	48	72	96
Control	0	0	0	0	0	0	0	0
0.2	0	0	1	0	0	1	1	0
0.3	0	1	0	1	0	0	1	1
0.5	2	1	0	1	1	0	0	1
0.7	2	1	1	0	2	2	0	1

Genotoxicity in Tissues of *Clarias gariepinus* Exposed to Cassava Effluent by Comet Assay

The genotoxic effects of Cassava Effluent in tissues of *Clarias gariepinus* is displayed in Plates 1 (gills), 2 (gonads) and 3 (liver), while Fig. 2 shows the total genotoxic potential of Cassava effluent on *Clarias gariepinus*. The result showed that cassava wastewater caused DNA damage in the tissue of African Catfish. The genotoxic effect of the wastewater increased at higher concentration indicating that the effect was concentration dependent. The outcome indicated a higher degree of DNA damage in both the gonads and liver at lower concentration of 0.2% and 0.3% respectively. The gills showed higher DNA damage at higher concentration of 0.7% compared to other organs.




Plate 1: Micrographs of DNA damage in the gills of *Clarias gariepinus* dosed with cassava wastewater at concentrations 0 %, 0.2 %, 0.3 %, 0.5 % and 0.7 %




Plate 2: Micrographs of DNA damage in the gonads of *Clarias gariepinus* dosed with cassava wastewater at concentrations 0 %, 0.2 %, 0.3 %, 0.5 % and 0.7 %




Plate 3: Micrographs of DNA damage in the liver of *Clarias gariepinus* dosed with cassava wastewater at concentrations 0 %, 0.2 %, 0.3 %, 0.5 % and 0.7 %




Fig. 2: Genotoxic potential of Cassava on *Clarias gariepinus*

DISCUSSION

This study used the comet assay to assess the genotoxic effects of cassava effluent on *Clarias gariepinus* tissues. Fish are essential to aquatic habitats because they are multifunctional members of the food chain. They react indirectly by eating contaminated aquatic creatures and show direct susceptibility to even very small quantities of harmful compounds in the water (Cavas & ErgeneGözükara, 2005). Fish can therefore act as early warning signs of pollution of aquatic ecosystems. Fish are good indicators of wastewater contamination in aquatic environments because of their frequent metabolic oscillations in reaction to wastewater (Cattaneo et al., 2011). Wastewaters and effluents are a substantial contribution to water pollution.

Death was caused by exposure to both low and high concentrations of cassava effluent in a dose-dependent manner (Table 1). Behaviors like erratic swimming patterns, avoidance-related surfacing behaviors that resulted in skin bleaching, abrupt, rapid movements, uncommon lethargy, and fish that settled at the lowermost part of the aquarium in a motionless state with sluggish opercula movements and gasping for air were also dose-dependent. Additionally, they are comparable to other related investigations (Cattaneo et al., 2011). The outcome provided a clear picture of how the effluent affected the aquatic habitat and aquatic organisms. Since we all come into contact with water and depend on it for living, this effect extends beyond aquatic habitats to other ecosystems and organisms.

The comet assay is used to quantify DNA strand breaks, an important indicator of genotoxicity in fish (Mitchelmore & Chipman, 1998). As a result of exposure to a variety of pollutants present in aquatic settings, the comet test is a commonly used, simple, and highly

sensitive tool for evaluating DNA damage in several fish organs, including gills, liver, gonads, kidney, and blood (Dhawan et al., 2009). Under alkaline circumstances, the comet assay can detect multiple types of DNA damage, such as single-strand breaks, alkaline labile sites, and wastewater-induced DNA cross-links (Kumar et al., 2010). Since the comet assay does not require cells to be dividing mitotically, it has significant advantages over conventional cytogenetic techniques (Ali et al., 2009; Ali & Kumar, 2008; Nwani et al., 2010).

The study's findings demonstrate that African catfish tissue suffers DNA damage from cassava wastewater. Higher concentrations of the wastewater had a greater genotoxic effect, suggesting that the effect was concentration dependent. The results showed that the liver and gonads had more DNA damage at lower concentrations of 0.3% and 0.2%, respectively. Elevated levels of DNA damage in gill cells may be caused by the gills' ongoing exposure to water contaminants that can harm DNA. Fish gills are essential organs that help in breathing, osmoregulation, and excretion. Gills are the main point of entry for effluents, hence they are good candidates for environmental impact studies since they are frequently regarded as great indicators of water quality. Respiratory distress can result from any damage to the gills caused by xenobiotic substances or modifications in membrane permeability, which can have a substantial impact on oxygen absorption rates. (Olufayo et al., 2016).

It is well known that a variety of genotoxic chemicals generate electrophilic free-radical metabolites and reactive oxygen species, which interact with DNA and cause disruption (Orororo et al., 2022; Efekemo and Orororo, 2022; Orororo et al., 2018). Research conducted by Chandra and Khuda-Bukhsh (2004) has demonstrated that the metabolism of azadirachtin generates free radicals and electrophilic ions that interact with nucleophilic sites in DNA, causing breaks and associated damage. According to Moore, Yedjou, and Tchounwou (2010), oxidative stress occurs naturally in an organism and is a major factor in the induction of cytotoxicity and genotoxicity in a number of critical tissues. According to Olourunfemi et al. (2024), the oxidative stress caused by the effluent's constituents in the fish tissues may be the cause of the genotoxic effects of cassava effluent that were reported in the study.

fish showed indications of liver and gill damage in a similar investigation examining the haematological and histological effects of cassava effluent on adult female African catfish, *Clarias gariepinus*. According to Olourunfemi et al. (2011), histological analyses of the kidney, gills, and liver of Nile Tilapia fingerlings (*Oreochromis niloticus*) subjected to cassava effluent revealed damage. It is clear that cassava wastewater can be dangerous to people if it is not properly cleaned before being released into water bodies. It is also harmful to aquatic life.

The inhibition of enzymes involved in DNA replication or repair may also be a significant contributing factor to DNA damage (Guilherme et al., 2012). Normal cellular functions depend on the integrity of DNA, and changes to DNA can cause tissue development, cell division, loss of cell structure, and degradation, which can ultimately lead to the loss of cellular regulatory systems. According to studies (Guilherme et al., 2012; Nwani et al., 2010), wastewater and cellular DNA interact physicochemically to cause a variety of primary alterations in DNA, such as single-strand breaks, double-strand breaks, DNA-protein crosslinks, and damage to purine and pyrimidine bases. These findings are consistent with the results obtain from this investigation.

CONCLUSION

According to the study, the comet assay method may be used to determine the genotoxic effects of cassava effluent in the tissues of the African cat fish, *C. gariepinus*. It's possible that the hazardous chemical cyanide is what causes the genotoxic consequences of cassava effluent. Therefore, it's crucial to adequately treat effluent from the cassava processing

companies before releasing it into bodies of water. Regulatory organizations should also be established to keep an eye on the effluent discharge procedure and production process in order to protect the aquatic ecology.

REFERENCES

- Amazeze, N.H. (2020). Comparative assessment of the acute toxicity, haematological and genotoxic effects of ten commonly used pesticides on the African Catfish, *Clarias gariepinus* Burchell 1822. *Heliyon*, **6**(8): e04768
- Ali, D., & Kumar, S. (2008). Long-term genotoxic effect of monocrotophos in different tissues of freshwater fish *Channa punctatus* (Bloch) using alkaline single cell gel electrophoresis. *Science of the Total Environment*, **405**(1): 345–350.
- Ali, D., Nagpure, N. S., Kumar, S., Kumar, R., Kushwaha, B., & Lakra, W. S. (2009). Assessment of genotoxic and mutagenic effects of chlorpyrifos in freshwater fish *Channa punctatus* (Bloch) using micronucleus assay and alkaline singlecell gel electrophoresis. *Food and Chemistry Tox*, **47**(3): 650–656.
- Alimba, C.G, Adekoya, K.O. and Soyinka, O.O. (2019). Exposure to effluent from pharmaceutical industry induced cytogenotoxicity, hematological and histopathological alterations in *Clarias gariepinus* (Burchell, 1822). *PubMed*, **18**: 63–78.
- Ariyomo TO, Jegede T, Adeniran AF, Omobepade BP (2017). Toxicity of Cassava Effluents to Catfish, *Clarias gariepinus*, and the Effect on Some Target Organs. *Annals of Aquaculture and Research*, **4**(2): 1034.
- Ayanda, O.I., Tolulope, A. and Oniye, S.J. (2021). Mutagenicity and genotoxicity in juvenile African catfish, *Clarias gariepinus* exposed to formulations of glyphosate and paraquat. *Science Progress*, **104**(2): 1-5
- Barabadi, H., Najafi, M., Samadian, H., Azarnezhad, A., Vahidi, H., Mahjoub, M.A., Koohiyan, M. and Ahmadi, A. (2019). A systematic review of the genotoxicity and antigenotoxicity of biologically synthesized metallic nanomaterials: are green nanoparticles safe enough for clinical marketing? *Medicina*, **55**(8): 439
- Busari, A. A., Efejene, I. O., Olayemi, S. O., Orororo, O. C., and Egbune, E. O. (2024). Evaluation of antibiotic use and analysis of ciprofloxacin and gentamicin residue in fish samples from farms in Lagos, Nigeria. *Environmental Monitoring and Assessment*, **197**(2): 127, 2024.
- Bolognesi, C. and Cirillo, S. (2014). Genotoxicity biomarkers in aquatic bioindicators. *Current Zoology*, **60**(2): 273–284
- Buckley, L. J. (1980). Changes in ribonucleic acid, deoxyribonucleic acid and protein content during ontogenesis in winter flounder, *Pseudopleuronectes americanus* and effect of starvation. *Fishery Bulletin*, **77**: 703–708.
- Cattaneo, R., Clasen, B., Loro, V. L., de Menezes, C. C., Pretto, A., Baldisserotto, B., De Avila, L. A. (2011). Toxicological responses of *Cyprinus carpio* exposed to a commercial formulation containing glyphosate. *Bulletin of Environmental Contamination and Toxicology*, **87**: 597–602.
- Cavas, T., & Ergene-Gözükara, S. (2005). Induction of micronuclei and nuclear abnormalities in *Oreochromis niloticus* following exposure to petroleum refinery and chromium processing plant effluents. *Aquatic Toxicology*, **74**: 264–271.
- Cordelli, E., Bignami, M. and Pacchierotti, F. (2021). Comet assay: a versatile but complex tool in genotoxicity testing. *Toxicology Research*, **10**(1): 68-78.

- Chandra, P., & Khuda-Bukhsh, A. R. (2004). Genotoxic effects of cadmium chloride and azadirachtin treated singly and in combination in fish. *Ecotoxicology and Environmental Safety*, **58**: 194–201
- De Lapuente, J. *et al.* (2015). The Comet Assay and its applications in the field of ecotoxicology: a mature tool that continues to expand its perspectives. *Frontiers in Genetics*, **6**(180)
- De Souza, J.M. *et al.* (2021). Mechanical and durability properties of compressed stabilized earth brick produced with cassava wastewater. *Journal of Building Engineering*, **44**: 103290.
- Dhawan, A., Bajpayee, M., & Parmar, D. (2009). Comet assay: a reliable tool for the assessment of DNA damage in different models. *Cell Biology and Toxicology*, **25**: 5–32
- Efekemo, O. and Orororo O. C. (2022). Effect of mixture of leaf extract of *Ocimum gratissimum* and *Vernonia amygdalina* on cadmium induced hepatotoxicity in wistar rats. *Journal of Biochemistry International*, 9(4):73-81
- Efekemo O., Davies I. C., Orororo C. O. (2024), Water Quality Assessment and Heavy Metal Levels in Mudskipper (*Periophthalmus Papilio*), Sediments and Water of Mangrove Swamps, Rivers State, Nigeria. *African Journal of Environment and Natural Science Research* 7(1), 128-145.
- Egbune, O. U., Egbune, E. O., Orororo, O. C., Ezedom, T., Onojakpor, O., Sabo, A. M. and Amadi, K. (2023). Chronic cassava meal modulates body weight, histology and weight of reproductive organs in male albino rats. *Toxicology and Environmental Health Sciences* <https://doi.org/10.1007/s13530-023-00179-4>
- Fawole, Olatunde & Yekeen, Taofeek & Ayandele, Abiodun & O, Ajayi & O., Aina & Omosehin, Omolaso & A., Odebunmi. (2014). Cassava mill effluent toxicity assessment using two bioassays and bacterial attributes. *Pollution Research*, **32**: 243–250.
- Ferk, F., Mišák, M., Grummt, T., Majer, B., Fuerhacker, M., Buchmann, C., Vital, M., Uhl, M., Lenz, K., Grillitsch, B. and Parzefall, W. (2009). Genotoxic effects of wastewater from an oncological ward. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, **672**(2): 69-75.
- Guilherme, S., Santos, M. A., Barroso, C., Gaivao, I., & Pacheco, M. (2012). Differential genotoxicity of roundup formulation and its constituents in blood cells of fish (*Anguilla anguilla*): considerations on chemical interactions and DNA damaging mechanisms. *Ecotoxicology*, **21**: 1381–1390
- Ibor, O.R. *et al.* (2020) “Contaminant levels and endocrine disruptive effects in *Clarias gariepinus* exposed to simulated leachate from a solid waste dumpsite in Calabar, Nigeria,” *Aquatic Toxicology*, **219**: 105375
- Ideta-Otsuka, M. *et al.* (2017). Epigenetic toxicity of environmental chemicals upon exposure during development - Bisphenol A and valproic acid may have epigenetic effects. *Food and Chemical Toxicology*, **109**: 812–816.
- Izah, S.C., Bassey, S.E. and Ohimain, E.I. (2017). Changes in the Treatment of Some Physico-Chemical Properties of Cassava Mill Effluents Using *Saccharomyces cerevisiae*. *Toxics*, **5**(4): 28.
- Kumar, R., Nagpure, N. S., Kwshwaha, B., Srivastava, S. K., & Lakra, W. S. (2010). Investigation of the genotoxicity of malathion to freshwater teleost fish *Channa punctatus* (Bloch) using the micronucleus test and comet assay. *Archives of Environmental Contamination and Toxicology*, **58**: 123–130.

- Lah, B., Malovrh, S., Narat, M., Cepeljnik, T. and Marinsek-Logar, R. (2004). Detection and quantification of genotoxicity in wastewater-treated *Tetrahymena thermophila* using the comet assay. *Environmental Toxicology: An International Journal*, **19**(6): 545-553.
- Lee, R.F. and Steinert, S. (2003) "Use of the single cell gel electrophoresis/comet assay for detecting DNA damage in aquatic (marine and freshwater) animals," *Mutation Research-reviews in Mutation Research*, **544**(1): 43–64
- Liney, K.E., Hagger, J.A., Tyler, C.R., Depledge, M.H., Galloway, T.S. and Jobling, S. (2006). Health effects in fish of long-term exposure to effluents from wastewater treatment works. *Environmental health perspectives*, **114**(1): 81-89.
- Luke Simon, Douglas T. Carrell. (2018). Laboratory Evaluation of Sperm Chromatin Structure, Editor(s): Michael K. Skinner. *Encyclopaedia of Reproduction (Second Edition)*, 106-111
- Mahaye, N., Thwala, M., Cowan, D.A. and Musee, N. (2017). Genotoxicity of metal based engineered nanoparticles in aquatic organisms: A review. *Mutation Research/Reviews in Mutation Research*, **773**: 134-160.
- Mahboub, H.H. *et al.* (2022). Silica nanoparticles are novel aqueous additive mitigating heavy metals toxicity and improving the health of African catfish, *Clarias gariepinus*. *Aquatic Toxicology*, **249**: 106238.
- Mitchelmore, C. L., & Chipman, J. K. (1998). Detection of DNA strand breaks in brown trout (*Salmo trutta*) hepatocytes and blood cells using the single cell gel electrophoresis (comet) assay. *Aquatic Toxicology*, **41**: 161–182.
- Mohamed, S.A.K.S., Upreti, S., Rajendra, S.V. and Dang, R. (2017). Genotoxicity: Mechanisms, testing guidelines and methods. *Global Journal of Pharmacy and Pharmaceutical Sciences*, **1**(5): 133-138.
- Moore, P. D., Yedjou, C. G., & Tchounwou, P. B. (2010). Malathion-induced oxidative stress, cytotoxicity and genotoxicity in human liver carcinoma (HepG2) cells. *Environmental Toxicology*, **25**(3): 221–226.
- Moussa, M.A., Mohamed, H.R.H. and Abdel-Khalek, A.A. (2022). Metal Accumulation and DNA Damage in *Oreochromis niloticus* and *Clarias gariepinus* After Chronic Exposure to Discharges of the Batts Drain: Potential Risk to Human Health. *Bulletin of Environmental Contamination and Toxicology*, **108**(6): 1064–1073
- Nagarathna, P.K.M., Wesley, M.J., Reddy, P.S. and Reena, K. (2013). Review on genotoxicity, its molecular mechanisms and prevention. *Int. J. Pharm. Sci. Rev. Res*, **22**(1): 236-243.
- Naguib, M. *et al.* (2022). Genotoxic evaluation of silver nanoparticles in catfish *Clarias gariepinus* erythrocytes; DNA strand breakage using comet assay. *Scientific African*, **16**: 01260.
- Nwani, C. D., Lakra, W. S., Nagpure, N. S., Kumar, R., Kushwaha, B., & Sirvastava, S. K. (2010). Mutagenic and genotoxic effects of carbosulfan in freshwater fish *Channa punctatus* (Bloch) using micronucleus assay and alkaline single-cell gel electrophoresis. *Food and Chemical Toxicology*, **48**(1): 202–208.
- Oghenejoboh, K.M. (2015). Effects of Cassava wastewater on the quality of receiving water body intended for fish farming. *British Journal of Applied Science and Technology*, **6**(2): 164–171.
- Oghenejoboh, K.M. *et al.* (2021). Value added cassava waste management and environmental sustainability in Nigeria: A review. *Environmental Challenges*, **4**: 100127.
- Oladokun, E.I., Sogbanmu, T.O. and Anikwe, J.C. (2020). Sublethal concentrations of dichlorvos and paraquat induce genotoxic and histological effects in the *Clarias gariepinus*. *Environmental Analysis, Health and Toxicology*, **35**(3): 2020013.

- Olorunfemi D. I., Oteri, O., Orororo O.C., Tina-Bayoko B. E., Pere C. and Osioma, E. (2024). Genotoxic Effects of Cassava Effluent on the Expression of Selected Genes in the African Catfish, *Clarias gariepinus*. *Journal of Advances in Biology & Biotechnology*; 27(5): 352-364.
- Olufayo M. O., & David O.T. (2016). The effect of cassava mill effluent on histology of *Clarias gariepinus* juveniles. *Revue Scientifique Et Technique Forêt Et Environnement Du Bassin Du Congo*, 7: 10–17.
- Orororo O. C., Tonukari, N. J., Avwioroko, O. J. and Ezedom, T. (2014). Effect of Supplementation of Animal Feed with Dried Cassava (*Manihot esculenta*) Peels, and Stems of *Vernonia amygdalina* and *Pennisetum purpureum* on Some Biochemical Parameters in Pigs. *Nigerian Society for Experimental Biology (NISEB) Journal* 14(4):177-183
- Orororo, O. C., Asagba, S. O, Tonukari, N. J., Okandeji, O. J., and Mbanugo J. J. (2018). Cadmium-Induced Testicular Damage in Wistar Rats: Protective Effects of *Hibiscus sabdariffa* L. Anthocyanins. *International Journal of Biochemistry Research and Review*; 21(4): 1-8
- Orororo, O. C., Efekemo, O. and Udi, O. A. (2022). Changes in liver histology, hematological parameters and lipid profile of cadmium-exposed rats treated with combined leaf extract of *Vernonia amygdalina* and *Occimum gratissimum*. *Asian Journal of Medicine and Health*; 20(11):195-203
- Osman, A.G.M. (2014). Genotoxicity tests and their contributions in aquatic environmental research. *Journal of Environmental Protection*, 5(14): 1391–1399.
- Osman, A.G.M. *et al.* (2008). Monitoring of DNA breakage in embryonic stages of the African catfish *Clarias gariepinus* (Burchell, 1822) after exposure to lead nitrate using alkaline comet assay. *Environmental Toxicology*, 23(6): 679–687.
- Sogbanmu, T.O., Osibona, A.O., Oguntunde, A.O. and Otitoloju, A.A. (2018). Biomarkers of toxicity in *Clarias gariepinus* exposed to sub-lethal concentrations of polycyclic aromatic hydrocarbons. *Afr. J. Aquat. Sci*, 43: 281-92
- Tonukari, N. J. Egbune E. Oliseneku, Oghenetega J, Avwioroko, Osuvwe C. Orororo, Akpovwehwee A. Anigboro (2016). A Novel Pig Feed Formulation containing *Aspergillus niger* CSA35 pretreated cassava peels and its effect on growth and selected biochemical parameters of pigs. *African Journal of Biotechnology*, 15(9): 776-785.
- Turan, F. *et al.* (2020). Heavy metal bioaccumulation, oxidative stress and genotoxicity in African catfish *Clarias gariepinus* from Orontes river. *Ecotoxicology*, 29(9): 1522–1537
- Wu, K., Zhou, Q. and Ouyang, S. (2021). Direct and indirect genotoxicity of graphene family nanomaterials on DNA—A review. *Nanomaterials*, 11(11): 2889.
- Yang, C., Song, G. and Lim, W. (2020). A review of the toxicity in fish exposed to antibiotics. *Comparative Biochemistry and Physiology C-toxicology & Pharmacology*, 237: 108840.
- Zaghloul, K. *et al.* (2020). Genotoxic and Histopathological Effects of Water Pollution on *Clarias gariepinus* Fish at Fayoum Governorate, Egypt. *Natural Resources*, 11(11): 499–519