

**Length-weight relationship and condition factor of *Synodontis schall***  
**((Bloch & Schneider, 1801), from Roseires reservoir, Sudan**

**Abstract:**

This study was carried out to investigate the length-weight relationship and condition factor of the Mochokid catfish *S. schall*, a commercially important fish species in the Roseires reservoir on the Blue Nile, Sudan. A total of 643 specimens were collected using the multi-filament gillnets of mesh sizes (4.0, 8.0, 10.0, and 12.0 cm) and twine (No. 2 and 12) during the period from November 2015 to October 2016; Samples were collected from four sites namely: Awal Bab, Elregiba, Kirma, and Wad El Mahi.; The length and weight of sampled fish were measured and recorded on the sampling sites and used to determine a length-weight relationship. The growth coefficient ( $b$ ) ranged from 1.196 to 2.649, with moderate to high correlations ( $r$ ) between 0.437 and 0.798. Values of ( $b$ ) were less than 3 indicating a negative allometric growth pattern, whereas weight increased at a lower rate than length. The condition factor ( $K$ ) ranged from  $1.966 \pm 0.696$  to  $3.856 \pm 3.856$ , suggesting the fish were in good health condition throughout the sampling sites. The findings of this study provide valuable insights into the biological aspects of *S. schall*, which can form the basis of sustainable management and conservation policy for the fishery resource of this economically important fish species in the Roseires reservoir.

**Keyword:**

*Synodontis schall*, Roseires Dam reservoir, Length-weight relationship, Condition factor.

**Introduction:**

The Sudan inland fisheries are located primarily around the Nile and its tributaries, major man-made lakes (Nubia and Merowe) and dam reservoirs, like Jebel Aulia, Roseires, Sinnar, Khashm El-Girba, Upper Atbara and Sittit Dams. The Roseires Dam, established in 1966 on the Blue Nile River at Ad Damazin town, is one of the most important dams in Sudan which was originally built to store water for irrigation purposes and hydropower generation, [1;2]. The height of Roseires Dam was increased by 10.0 meters during the period 2006-2012, allowing greater water storage, and creating a 75 km long lake, forming an important fishing ground with a maximum sustainable production of 1,000 tons/year [2].

The expansion of Roseires Dam reservoir is expected to change the fish abundance and distribution, alter primary productivity of the water, promote individual growth rates of fishes, and increase overall fish production. Additionally, it is anticipated that the significant increase in the volume of water in Lake Roseires will lead to a corresponding increase in fish production from the reservoir [3], and future changes in the fisheries sector, which can achieve positive changes in the socioeconomic status and livelihood of people inhabiting the region [2].

Approximately 112 species of the genus *Synodontis* were recorded in the Nile River and Eastern Rift Valley Lakes (i.e. Egypt, Ethiopia, Sudan), and across West African countries (i.e. Ivory Coast, Ghana, Guinea, Guinea-Bissau, Liberia, Niger, Nigeria, Senegal, Sierra Leone, [4 and 5]).

Studies on length-weight relationships are of considerable importance for proper fishery management of the stocks and biomass estimation of any fish species [6; 7 and 8]. LWR can be used to predict the weight of a fish when its length is known (or vice versa), for assessing fishery yields [9, 10 and 11], and provide information on the fish growth pattern [12, 13, 14, 15, 16 and 17]. LWR has also been widely used in local and inter-regional morphological comparison of populations [18 and 10], as well as estimating weight at age, and assessing age structure and function of fish populations [19, 16, 20, 21 and 11]. Moreover, LWR is used to evaluate the index of well-being of the fish population and obtain information on the condition of fishes in order to determine whether somatic growth is isometric or allometric [22 and 23].

The condition factor (K) of a fish (also known as Fulton's condition factor) is an important biological parameter that indicates the suitability of a specific body of water for growth of fish, and as an index of species average size; It also reflects the physico-chemical and biological factors, and interactions of feeding conditions, parasitic infections and physiological factors [24]. In addition, it indicates the changes in food reserves in the water body, and an indicator of the general health condition, and an expression of the relative fatness of the fish [10]. Therefore, understanding the condition factor ('K') is important in fishery assessment and management of fish populations, as it gives us clear knowledge regarding maturity, spawning and life cycle of fish at different body lengths during their life span, availability of food in a water body, and environmental factors affecting growth of a fish [25, 22 and 20].

Analysis of LWRs and condition factors has evolved from log-transformed mean weights to non-linear fitting approaches that leverage modern statistical software and computing power,

allowing for a more accurate representation of the relationship between fish length and weight [26, 27, 28, 20 and 30].

The present study aims to provide the baseline information on the length-weight relationships and condition factor of the commercially important Mochokid catfish, *S. schall* in Roseires Dam reservoir on the Blue Nile, Sudan. The study's results will be useful in the sustainable management and exploitation of this species' fishery and the protection of its natural habitats.

## Materials and Methods:




Fig. 1: Roseires reservoir area before and after the heightening of the Dam.

### Study area:

Fish (*S. schall*) samples were collected from the following sites:

**Awal bab sampling site:** This is one of the biggest fishing sites on the western bank of the Blue Nile. It lies about 4 km southeast of Ad Damazin town.

**EL-Regayba sampling site:** is located on the western bank of the Blue Nile about 16 km southeast of Ad Damazin town.

**Kirma sampling site:** is located on the eastern bank of the Blue Nile, and lies about 43 Km east of Ad Damazin town.

**Wed-ELmahi sampling site:** located on the eastern bank of the Blue Nile, about 80 km south of Roseires city on the end of the Dam lake.

The study extended 12 months (November 2015 - October 2016). Sampling of fish took place monthly. During each sampling, samples of fish representing different sizes were randomly selected from the catch.

### Sample collection:

During the study period, about 643 specimens of *Synodontis schall* were collected from the four sampling sites, using four multi-filament gill nets of various mesh sizes and twine numbers as shown below:

Table 1. Specifications of gillnets used for collection of fish samples during the study period, October 2015 to September 2016.

Twine No.	Length (m)	Depth (m)	Mesh size (cm)
2	50	2	4
12	90	4	8
12	95	4	10
12	100	4.5	12

### Measurements of length and weight:

The total length (tip of snout to tip end of caudal fin) and standard length (tip of snout to base of caudal fin) of each fish were measured to the nearest 0.01 mm. using a standard measuring board. The body weight of sampled fish was taken to the nearest gm., using a digital weighing balance, version FRUIT 2000B.

### Length-weight relationship:

Excel package was used to plot the curve of the relationship between the standard length and weight of *S. schall*, and the linear equation was then obtained from logarithm transformation. Values of the constants 'a' and 'b' for the fish were obtained from the length-weight relationship according to [17], using the equation:

$$W = aL^b$$

Where:

W = total weight in grams

L = total length in cm

b = a constant of the relationship representing the slope of the equation.

a = a constant of the relationship representing the intercept on the "y" axis

### Condition Factor (CF) or (Fulton's factor) (FCF)

The well-being of *S. schall* was calculated according to [17], according to the following formula:

$$FCF = \frac{W}{L^b} \times 100$$

Where W: weight of fish; L: standard length of fish, and 'b': constant.

### Statistical Analysis

Statistical analysis was made by using a personal computer and the advanced computer Excel program.

### Results and Discussion:




Fig.2: Linear fit of the length-weight relationship of *S. schall* in Awal Bab site during the study period (2015/2016).




Fig.3: Linear fit of the length-weight relationship of *S. schall* in Elregiba site during the study period (2015/2016).




Fig.4: Linear fit of the length-weight relationship of *S. schall* in Kirma site during the study period (2015/2016).




Fig.5: Linear fit of the length-weight relationship of *S. schall* in Wad Elmahi site during the study period (2015/2016).

Table 2: Shows linear fit of length-weight relationship *S. schall* during the study period (2015/2016).

Site	<i>b</i>	<i>a</i>	<i>r</i>
Awal Bab	2.649	-1.344	0.798
Elregiba	1.196	-0.375	0.437
Kirma	2.156	-0.807	0.539
Wad El Mahi	2.426	-1.117	0.626

The genus *Synodontis* comprises several commercially important species, contributing up to 40% of the total landing weight in some regions of Africa [30]. The length-weight relationship in fishes is not constant throughout the year and is affected by factors such as the availability of food, rate of feeding, development of gonads, and spawning season [32 and 31].

Previous studies of *Synodontis schall* reported 'b' values (the exponent in the length-weight relationship) ranging from 2.4977 to 3.522, indicating both allometric and isometric growth patterns [31, 32, 33, 34, 35 and 36].

The results of the current study indicated a negative allometric growth pattern of *S. schall* in all sample sites. The values of the exponent ( $b$ ) were found to be 2.649, 1.196, 2.156, and 2.426, with high to moderate correlation ( $r = 0.798, 0.437, 0.539, \text{ and } 0.626$ ) in Awal Bab, Elregiba, Kirma, and Wad El-Mahi, respectively (Fig. 2-5 and Table 2). These results agree with previous findings in the Lower Ogun River (Nigeria), Okpara Stream (Benin), Khashm El Girba reservoir and Atbara River, Upper Atbara and Sittit dam complex, and the River Benue (Nigeria) [32,31; 43,1 and 35].

However, [36] reported that the length-weight relationships of Mochokidae species in the Niger River (Benin) showed both negative and positive allometric growth, with values of 'b' ranging from 2.236 to 3.380 and high correlations ( $r = 0.87 \text{ to } 0.99$ ). Regarding *S. schall*, they recorded a negative growth coefficient of  $b = 2.6327$ . [37] stated that 'b' value of *S. schall* in Lake Ayamé 2 (Côte d'Ivoire) ranged between 2.516 and 2.785 during the dry season, showing negative allometric growth, but increased to 3.0008 - 3.001 during the rainy season, indicating isometric growth. In contrast, [35] reported that the regression coefficient 'b' for the length-to-body weight relationship of *S. schall* in the White Nile showed isometric growth, with 'b' ranging from 2.985 and 3.033.

The results of the current study indicated a negative allometric growth pattern of *S. schall* in all sampling sites. The values of the exponent 'b' were found to range from 2.649, 1.196, 2.156, and 2.426, with high to moderate correlation ( $r = 0.798, 0.437, 0.539, \text{ and } 0.626$ ) in Awal Bab, Elregiba, Kirma, and Wad El-Mahi, respectively (Fig. 2-5, and Table 2).

These results agree with previous findings recorded for other tropical African rivers, such as, the Lower Ogun River (Nigeria), Okpara Stream (Benin), Khashm El Girba reservoir and Atbara River, Upper Atbara and Sittit dam complex (Sudan), and the River Benue (Nigeria) [31, 30,32, 1 and 33].

However, [36] reported that, the length-weight relationships of Mochokid species in the Niger River (Benin) showed both negative and positive allometric growth, with 'b' values ranging from 2.236 to 3.380 and high correlations ( $r = 0.87 \text{ to } 0.99$ ). With respect to *S. schall*, they recorded a negative

growth coefficient of “ $b = 2.6327$ . [37] stated that the value of ‘ $b$ ’ of *S. schall* in Lake Ayamé 2 (Côte d'Ivoire) ranged between 2.516 and 2.785 during the dry season, indicating negative allometric growth, and increased to values between 3.0008 and 3.001, during the rainy season, indicating isometric growth. In contrast, [35] reported that the regression coefficient ‘ $b$ ’ for the length-to-body weight relationship in *S. schall* from the White Nile showed isometric growth, with ‘ $b$ ’ ranging between 2.985 and 3.033.

Table 3: Shows condition factor (C.F) of *S. schall* in Awal Bab, Elregiba, Kirma and Wad El-Mahi during the studyperiod (2015/2016).

Month	Awal bab	Elregiba	Kirma	Wad El-Mahi
Dec-15	3.455 ± 0.048	1.535 ± 0.549	0 ± 0.000	2.302 ± 0.476
Jan-16	3.760 ± 0.000	1.890 ± 0.201	2.030 ± 0.707	3.082 ± 1.408
Feb-16	3.521 ± 1.287	2.491 ± 0.268	1.690 ± 0.268	1.962 ± 0.416
Mar-16	3.398 ± 0.331	2.349 ± 0.081	1.758 ± 0.000	2.072 ± 1.251
Apr-16	3.029 ± 0.679	2.414 ± 0.507	2.074 ± 0.000	1.970 ± 0.744
May-16	3.549 ± 0.000	2.586 ± 0.253	3.127 ± 0.121	1.912 ± 0.154
Jun-16	3.902 ± 0.694	2.700 ± 0.662	1.979 ± 0.000	1.914 ± 0.504
Jul-16	3.108 ± 0.108	2.666 ± 1.124	1.850 ± 0.470	1.828 ± 0.166
Aug-16	3.002 ± 0.975	2.245 ± 0.905	2.417 ± 0.845	1.969 ± 0.594
Sep-16	3.808 ± 0.800	2.573 ± 0.000	2.080 ± 0.415	1.733 ± 0.257
Oct-16	3.957 ± 0.242	2.627 ± 1.185	1.619 ± 0.246	1.871 ± 0.597
Nov-16	0.812 ± 0.983	2.129 ± 0.120	1.865 ± 0.399	2.228 ± 0.760
Average	3.856 ± 3.856	2.428 ± 0.738	1.982 ± 0.614	1.966 ± 0.696

The condition factor (K) is a useful index for assessing the health and general well-being of fish, with values typically ranging from 2.5 to 4, for mature freshwater fish [38, 39 and 29]. Previous values of condition factor reported for *Synodontis schall* ranged from 1.422 to 3.269 [32, 33, 35, 36 and 37].

The findings of the current study indicate that the maximum values of condition factor recorded for *S. schall* were 3.957, 2.700, 3.127, and 3.082, while minimum values were 0.812, 1.535, 0.000, and 1.733, for Awal Bab, Elregiba, Kirma, and Wad El-Mahi sites, respectively. It can be inferred from the foregoing data that the high values of condition factor of *S. schall* in Awal Bab and Elregiba sites on the western bank of the Blue Nile, and relatively low values of condition factor recorded in

Kirma and Wad El-Mahi sites, on the eastern bank of the Blue Nile, throughout the year (Table, 3) This may be due to abundance of food items, and favorable environmental conditions for growth and reproduction which resulted in good health conditions of *S. schall* on the western bank of the Blue Nile.

These findings are consistent with those of [32], who reported good health conditions (K ranging from 0.506 to 3.415) for six species of fish, including *S. schall* (K ranging from 1.422 to 1.844), collected from the Khashm El Girba and Atbara River. Similarly, [40] reported condition factors ranging from 0.443 to 2.191 for 21 species in the Nile within Egypt, including *S. schall* (K = 1.437). Additionally, [1] reported a condition factor of 2.902 for *S. schall* in the Upper Atbara and Sittit dam complex, while [33] recorded mean condition factors of *S. schall* of 2.874, 2.838, and 2.855 for female, male, and combined sexes, respectively. [36] recorded condition factors ranging from 0.409 to 7.276 for *S. frontosus* and *S. sorex*, and 2.732 for *S. schall*. [37] reported good physiological condition for *S. schall* in Lake Ayamé 2 (Côte d'Ivoire), during the rainy season, with 'K' values varying between 1.0375 and 1.357.

#### **Conclusion:**

The present study of *Synodontis schall* provides baseline information on the length-weight relationship of one of the economically important fishes in Sudan. It indicated negative allometric growth patterns of the fish in all sampling sites in Roseires Reservoir. On the other hand, condition factor showed high values on the western bank of the Reservoir and relatively low values on the eastern bank. This may be attributed to anthropogenic influences, physiochemical and biological changes in the local environment, over-exploitation of this species, and changes in state of the gonadal development of the fish. The results of this study may be useful in the management of the fisheries resources of *S. schall* in Roseires Reservoir.

#### **Disclaimer (Artificial intelligence):**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### **References:**

[1] Adam, H., A., H. and Hamad, A., E. (2021). Length-weight relationship and condition factor of *Labeo niloticus*, *Synodontis schall* and *Auchenoglanis occidentalis*, in Upper Atbara and Setit

dam complex, Gadarif state, Sudan. *Glob., J. Fish., Sci.*, Vol., 3(4): 37-34. <https://doi.org/10.31248/GJFS2021.027>.

[2] Karra, A. M. (1999). Biological studies on indicator fish species prior to Expected Heightening of Roseires Dam *M.Sc. Thesis University of Khartoum – Sudan*.

[3] Hamid, A. M. A. (2018). Aspects study of some biological of four commercially important fishes in Roseires Reservoir after heightening of the Dam. *M. Sc. Thesis, Sudan Academy of Sciences Animal Research Council*. Khartoum, Sudan.

[4] Froese, et. al., (2011). Rainer and Daniel Pauly, eds, (2011). “*Synodontis schall*” in *fishbase*. December 2011 version.

[5] De Weirdt, D., Huyghe, C.E.T. and Vreven, E.J.M.W. N.(2023). On the occurrence and distribution of *Synodontis nigrita* and *Synodontis violaceus* (Siluriformes: Mochokidae) in the Nilo-Sudan, and Upper and Lower Guinea ichthyofaunal provinces. *J. Fish Biol.*, 103(3), pp.741-746. <https://doi.org/10.1111/jfb.15414>.

[6] Hossain, M. Y., Rahman, M. M., Fulanda, B., Jewel, M. A. S., Ahamed, F., Ohtomi, J. (2012). Length-weight and length-length relationships of five threatened fish species from the Jamuna (Brahmaputra River tributary) River, northern Bangladesh. *J. Appl., Ichthyo.*, 28, 275-277. <https://doi.org/10.1111/j.1439-0426.2011.01900.x>.

[7] Tah, L., Bi Goore, G., Da Costa, K. S. (2012). Length-weight relationships for 36 freshwater fish species from two tropical reservoirs: Ayamé I and Buyo, Côte d’Ivoire. *Revista de biologia tropical*, 60, 1847-1856. Available from: [http://www.scielo.sa.cr/scielo.php?script=sci\\_arttext&pid=S0034-77442012000400036&lng=en](http://www.scielo.sa.cr/scielo.php?script=sci_arttext&pid=S0034-77442012000400036&lng=en).

[8] Lederoun, D., LalPyP K. R, Boni, A. R., Amouso, G., Vodougnon, H., Adjibogoun, H., LalPyP, P.A. (2018b). Length-weight and length-length relationships of some of the most abundant species in the fish catches of Lake Nokoué and Porto-Novo Lagoon (Benin, West Africa). *Lakes & Reservoirs*, 1-7. <https://doi.org/10.1111/lre.12243>.

[9] Froese, R., Thorson, J. T., Reyes, R. B. (2014). A Bayesian approach for estimating length-weight relationships in fishes. *J. Appl., Ichthyo.*, 30, 78-85. <https://doi.org/10.1111/jai.12299>.

[10] Froese, R. (2006). Cube law, condition factor and weight-length relationships: History, meta-analysis and recommendations. *J. Appl., Ichthyo.*, 22 (2006), pp. 241-253. <https://doi.org/10.1111/j.1439-0426.2006.00805.x>.

- [11] Olopade, O. A., Dienye, H. E. and Eyekpegba, A., (2018). Length frequency distribution, length-weight relationship and condition factor of cichlid fishes (Teleostei: Cichlidae) from the New Calabar River, Nigeria. *Iranian J. Ichthyol.*, 5(1), pp.74-80. doi: [10.22034/iji.v5i1.251](https://doi.org/10.22034/iji.v5i1.251).
- [12] Mehanna, S.F. and Farouk, A.E. (2021). Length-Weight Relationship of 60 Fish Species from the Eastern Mediterranean Sea, Egypt (GFCM-GSA 26). *Front., Mar., Sci.*, 8. Article 625422, 10.3389/fmars.2021.625422. <https://doi.org/10.3389/fmars.2021.625422>.
- [13] Jisr, N.; Younes G., Sukhn C., and El-Dakdouki M. H. (2018). Length-weight relationships and relative condition factor of fish inhabiting the marine area of the Eastern Mediterranean city, Tripoli-Lebanon. *Egyptian J. Aquat. Res.*, 44: 299-305. <https://doi.org/10.1016/j.ejar.2018.11.004>.
- [14] Kuriakose, S. (2017). Estimation of length weight relationship in fishes. Fishery Resources Assessment Division, ICAR-Central Marine Fisheries Research Institute. 215-220. url: <http://eprints.cmfri.org.in/id/eprint/12178>.
- [15] King, M., 2007. Fisheries biology, assessment and management. *Wiley Blackwell*, 400P.
- [16] Morato, T., Afonso, P., Lourinho, P., Barreiros, J. P., Sanstos, R. S. and Nash, R. D. M., (2001). Length-weight relationships for 21 coastal fish species of the Azores, North-eastern Atlantic. *Fish., Res.*, **50**: 297-302. [https://doi.org/10.1016/S0165-7836\(00\)00215-0](https://doi.org/10.1016/S0165-7836(00)00215-0).
- [17] Abdalla, M. Y. M. (2018). Biological aspects of *Oreochromis niloticus* and *Labeo niloticus* from Khashm El-Girba reservoir and Atbara River, Kassala State, Sudan. *M. Sc. Thesis*, Department of Zoology, Faculty of Science, University of Khartoum, Sudan.
- [18] Ali, A.N.; Dahanukar, R. and Raghavan (2013). Length-weight and length-length relationship of three species of snakehead fish, *Channadiplogramma*, *C.maruliusand*, and *C. striata* from the riverine reaches of Lake Vembanad, Kerala, India. *J. Threat., Taxa.*, 5(13): 4769–4773. DOI: <http://dx.doi.org/10.11609/JoTT.o3353.4769-73>.
- [19] Özaydın, O., Uçkun, D., Akalın, S., Leblebici, S. and Tosunoğlu, Z. (2007). Length-weight relationships of fishes captured from Izmir Bay, Central Aegean Sea. *J. Appl., Ichthyo.*, Vol., 23: 695–696. <https://doi.org/10.1111/j.1439-0426.2007.00853.x>.
- [20] Haimovici, M. and Velasco, G., 2000. Length-weight relationship of marine fishes from southern Brazil. *Naga, The ICLARM Quarterly*, 23(1), 19-23.

- [21] Petrakis, G. and Stergiou, K.I. (1995). Weight-length relationships for 33 fish species in Greek waters. 16 (2) *Fish., Res.*, 21(3-4), 465- 469.[https://doi.org/10.1016/0165-7836\(94\)00294-7](https://doi.org/10.1016/0165-7836(94)00294-7).
- [22] Ujjania, N.C., Kohli, M.P.S. and Sharma, L.L. (2012). Length-weight relationship and condition factors of Indian major carps (*Catlacatla*, *Labeorohita* and *Cirrhinus mrigala*) in Mahi Bajaj Sagar, India. *Res.,J.Bio.*, 2(1), 50-56.ISSN 2049-1727. Available online at: [www.scientific-journals.co.uk](http://www.scientific-journals.co.uk).
- [23] Gurkan, S. and Taskavak, E. (2007). Length-weight relationships for syngnathid fishes of the Aegean Sea, Turkey. *Belgian J. Zool.*, 137(2), 219.
- [24] Alam, M.M., Rahman, M.T. and Parween, S. (2014). Morphometric characters and condition factors of five freshwater fishes from Pagla River of Bangladesh. *Inter., J. Aquat., Bio.*, 2(1), 14- 19.<https://doi.org/10.22034/ijab.v2i1.18>.
- [25] Dan-Kishiya, A.S. (2013). Length-weight relationship and condition factor of five fish species from a tropical water supply reservoir in Abuja, Nigeria. *American J. Res.,Commun.*, 1(9), 175- 187.
- [26] Abdalla, M. Y. M., Abdelhalim, A. I. and Adam, A. E. (2020). Study of some biological aspects of the Nile carp, *Labeo niloticus* (Pisces, Cyprinidae) from Khashm El-Girba reservoir and Atbara River, Sudan: I Abundance; sex ratio; gonado-somatic index and breeding season. *Asian J. Res., Zool.*, Vol., 3(3)21-28.[DOI: 10.9734/AJRIZ/2020/v3i3330093](https://doi.org/10.9734/AJRIZ/2020/v3i3330093).
- [27] Shelton, A.O. and Mangel, M. (2012). Estimating von Bertalanffy parameters with individual and environmental variations in growth. *J. Biol., Dynamics*, 6(sup2), pp.3-30. <https://doi.org/10.1080/17513758.2012.697195>.
- [28] Liu, J. (2023). Spatiotemporal models for exploring variability in scallop condition across the Bay of fundy. Available online at: <http://hdl.handle.net/10222/83249>.
- [29] Olagbemide, P. T. and Owolabi, O. D. (2023). Length-weight relationship and condition factor of *Oreochromis niloticus* (Linnaeus, 1758) in selected tropical reservoirs of Ekiti State, southwest Nigeria. DOI: [10.18006/2023.11\(4\).707.719](https://doi.org/10.18006/2023.11(4).707.719).
- [30] Imorou, R. S., Adite, A., Sossoukpe, E., Abou, Y. (2019).Length-weight models and condition factors of fishes from Okpara stream, Oueme River, Northern-Basin. *Inter., J. of Forest, Anim., and Fish., Res.*, Vol., 3(3): 65-80. <http://dx.doi.org/10.22161/ijfaf.3.3.1>.

- [31] Adeosun, F. I., Abdulraheem, I., Adesina, B. T. and Amrevuawho, O. M. (2017). Food and feeding habits and allometric relationship of *Synodontis schall* in Lower Ogun River, Akomoje, Ogun state, Nigeria. *Pertanika J. Trop. Agri., Sci.*, 40(4): 601 – 610. ISSN: 1511-3701.
- [32] Ahmed, Egbal, O., Ali, M. E., Aziz, Afra, A. (2011). Length-weight relationships and condition factors of six fish species in Atbara river and Khashm El-Girba reservoir, Sudan. *Inter., J. Agri., Sci.*, Vol., 3(1): 65-70. Online available at: <http://www.bioinfo.in/contents.php?id=26>.
- [33] Akombo, p. M., Akange, E., T., Adikwu, I., A. and Araoye, P. A. (2014). Length-weight relationship, condition factor and feeding habits of *Synodontis schall* (Bloch and Schneider, 1801) in River Benue at Makurdi, Nigeria. *Inter., J. Fish., Aqua., Stud.*, 1(3): 42-48.
- [34] Dadebo, E. (2016). Length-weight relationship, breeding season, sex ratio, maturity and fecundity of the Nile catfish *Synodontis schall* (Bloch and Schneider, 1801) (Pisces: Mochokidae) in Lake Chamo, Ethiopia. *Ethiopian J. Sci., Technol.*, 9(2), pp.87-102. DOI: [10.4314/ejst.v9i2.2](https://doi.org/10.4314/ejst.v9i2.2).
- [35] Abd-Elhalim, A. I., (1983). Investigations into the growth and reproduction of *Synodontis schall* (Bloch – Schneider, 1801) from the White Nile. *M. Sc. Thesis. Dep. Of Zoology, Faculty of Science, University of Khartoum*.
- [36] Arame, H., Adite, A., Adjibade, K.N., SidiImorou, R., Sonon, P.S. (2020). Length-weight relationships and condition factors of Mochokidae (Pisces: Teleostei: Siluriformes) from Niger River, Northern Benin. *Aquat., Res.*, 3(2), 72-84. <https://doi.org/10.3153/AR20007>.
- [37] Georges, B. K., N'Doua, E. R., Theophile, B. A., Silvain, Y. S. and Valentin, N. (2017). Seasonal variation in the length-weight relationships and condition factor of *Synodontis schall* (Bloch and Schneider, 1801) (Siluriformes: Mochokidae) in man-made Lake Ayame 2 (Cote d'Ivoire). *Inter., J. Fish., Aquat., Stud.*, Vol., 5(1): 173-177.
- [38] Kumolu-Johnson, C. A., and Ndimele, P.E., (2010). Length-weight relationships and condition Factors of twenty-one fish species in Ologe lagoon, Lagos, Nigeria. *Asian J. of Agri., Sci.*, 2(4): 174 – 179. ISSN: 2041-3890.
- [39] Anene, A., (2005). Condition factors of four cichlid species of a man-made lake in Imo State, Southeast, Nigeria. *Turk. J. Fish. Aquat. Sci.*, 5: 43-47.

[40] Shalloof, K., A., SH. and El-Far, A., M. (2017). Length-weight relationship and condition factor of some fishes from the River Nile in Egypt with special reference to four Tilapia species. *Egy., J. Aqua., Biol., Fish.*, Vol., 21(2): 33-46.DOI: 10.21608/ejabf.2017.3296.

UNDER PEER REVIEW