

## Original Research Article

# COMPARATIVE EVALUATION OF THE EFFECT OF DIFFERENT MEDIA IN AQUAPONIC SYSTEM ON THE PERFORMANCE OF PLANT GROWTH AND YIELD FOR TOMATO

---

### ABSTRACT

**Aims:** To evaluate the performance of tomato crops grown in four different media in a bell siphon arranged aquaponics system.

**Study design:** completely randomized design.

**Place and Duration of Study:** Hi-tech Research and Training Unit (HTR&TU), Instructional Farm (IF), Kerala Agricultural University, Vellanikkarabetween 03-08-2021 to 03-01-2022 and from 01-02-2023 to 30-06-2023.

**Methodology:** An aquaponic system was equipped with twenty-four media beds with bell siphons, three filter tanks (sedimentation tank, mechanical filter and bio filter), three fish tanks having 1000L capacity each, a submersible pump, and two aerators. Two hundred genetically improved farmed tilapia (GIFT) fingerlings were grown in the fish tanks. Four types of media viz. 20mm gravel (M<sub>1</sub>), 8-20mm broken tiles (M<sub>2</sub>), 8-15mm hydroton (M<sub>3</sub>), and 8-10mm gravel (M<sub>4</sub>) were filled in the media beds. The two varieties of tomato, Manulakshmi and Yakamoz RZ were raised. Plant height, plant spread, days to first flowering, days to first harvest, total yield and average fruit weight were observed to compare the crop performance in each media and identify the best-performing media in an aquaponics system.

**Results:** During the first and second season of Manulakshmi in the aquaponics system, plants grown in 8-10mm gravel exhibited significantly superior results than all other media. The maximum plant spread (66.833cm), plant height (140.556cm), total yield (1.730Kg), and average weight of fruit (38.433g) were observed in 8-10mm gravel during the first season. In the second season of Manulakshmi, the observed plant spread, plant height, total yield and average weight of fruit were 140.778cm, 66.556cm, 1.729Kg and 38.611g respectively. In the first season of Yakamoz RZ in the aquaponics system, maximum plant height (145.222cm) and total yield (3.871Kg) were observed in 8-10mm gravel whereas, maximum plant spread (50.500cm) was observed in 20mm gravel. The days to first flowering and days to first harvest were comparatively shorter in 8-10mm gravel in both varieties.

**Conclusion:** Plants grown in 8-10mm gravel exhibited superior results than all other media. Variation of plant characteristics among different media in the aquaponics system reveal that 8-10mm gravel is more favorable for the growth and yield of tomato plants than other media.

*Keywords: Aquaponics, soilless media, tomato, crop growth, yield*

## 1. INTRODUCTION

India is one of the largest agricultural commodities producing country on a global basis utilising more than half of its land area for crop production. There has been remarkable increase in agricultural productivity over the last decades [1]. On the contrary, Indian agriculture is riddled with the challenge of feeding an escalating human population in the foreseeable future. Intensity of population in Kerala is 35 million (World population review), which is 2.5% of India's population. Kerala's land is three times as densely settled as the rest of India. The unavailability of land, unpredictable climatic conditions, disinterest of youth in agriculture, scarcity of labour, high labour wages, and inadequate market facilities prevents the state from realization of its production potential. Improved agricultural production systems can be a sustainable way of addressing this issue. To sustain a balance between food security and crop production, aquaponics within greenhouse technology could be exploited. The use of aquaponics methods holds the potential to significantly contribute to food provision and address global challenges, including water scarcity, food security, water pollution, high energy consumption, and the extensive transportation of food over long distances [2]. The aquaponics system establishes a symbiotic relationship between the fish and plants, where the fish provide nutrients to the plants, and in return, the plants purify the wastewater by utilizing these nutrients within the environment where the fish reside [3], [4], [5]. Aquaponics has been recognized as a forward-thinking farming approach combining aquaculture system, filtration system and hydroponics system, which involves crop production in a soilless substrate. In hydroponics systems plants growing in the soilless medium may comprise of perlite, vermiculite, rock wool, peat moss, coir, composted pine bark, sawdust, sand, or gravel [6], [7]. Studies indicate that the choice of soilless media significantly influences nitrifying bacteria counts [8]. Graber and Junge (2009) have revealed that the integration of aquaculture and hydroponics systems can be utilized for producing fish and vegetables like aubergine, tomato, and cucumber suitable for human consumption. Hence a study was conducted to evaluate the performance of two varieties of tomato in an aquaponic system in two different seasons and find the best media giving better result with each variety.

## 2. MATERIAL AND METHODS




Fig.1. The schematic representation of system

Study was carried out at Hi-tech Research and Training Unit (HTR&TU), Instructional Farm (IF), Kerala Agricultural University, Vellanikkara. The experiment was conducted in a polyhouse located at 10°32'50" North latitude and 76°16'18" East longitude.

The applied method encompassed an aquaponic system consisting of twenty-four media beds with bell siphons, three filter tanks (sedimentation tank, mechanical filter and bio filter), three fish tanks having 1000 lit capacity each, a submersible pump, and two aerators. The system was designed in such a way that nutrient-rich water circulates throughout the day and night.

Field experiment was laid out in a completely randomized design. A naturally ventilated polyhouse with a floor area of 150 m<sup>2</sup> was selected for the study [9]. Three intermediate bulk container (IBC) tanks (1.1m x 0.92m x 1m) served as nutrient water reservoir as well as fish culture tank. An aerator with six knobs, an air output of 10 liters/min was used to provide oxygenation in both systems. The water was channelled from the fish tank to the sedimentation tank, mechanical filter and the bio filter and finally to the hydroponic beds where plants were cultivated. A bypass arrangement was provided so that the excess flow from the pump was redirected back to the fish tank. The filtered water from the sedimentation tank was transferred through an inlet connection to the bottom of the mechanical filter and then to the bio-filter. After completing the filtration process in sedimentation tank, mechanical filter and biofilter (removing soluble and insoluble unwanted impurities), water was allowed to flow into hydroponics system through the main pipe and laterals. Two inlets were provided in each bed. The outflow from each bed were allowed to flow in to the drainage pipe. Two hundred genetically improved farmed tilapia (GIFT) fingerlings were introduced into the fish tanks in proportions of 70, 70, and 60 in the first, second, and third tanks, respectively. Used barrels were cut horizontally into two equal halves and were used for making two media beds. Four types of media viz. 20mm gravel (M<sub>1</sub>), 8-20mm broken tiles (M<sub>2</sub>), 8-15mm hydroton (M<sub>3</sub>), and 8-10mm gravel (M<sub>4</sub>) were filled in the beds. The comparative evaluation of the media was conducted by raising two varieties of tomato, Manulakshmi and Yakamoz RZ, during the periods from 03-08-2021 to 03-01-2022 and from 01-02-2023 to 30-06-2023. Two tomato varieties were transplanted in groups of three in each bed with 50x50cm spacing. Plant height and plant spread were observed at 14 Days after planting (DAP), 28 DAP, 42 DAP, 56 DAP, 70 DAP, and 84 DAP. Days to first flowering and first harvest also made. The weight of a single fruit harvested at different stages and the total yield of the tagged plant were noted. The comparison of media within the system was statistically analysed using Analysis of Variance (ANOVA).

### 3. RESULTS AND DISCUSSION

The analysis of growth parameters of Manulakshmi in aquaponics system during first and second season (Table 1 and 2 respectively) showed that the plants grown in M<sub>4</sub> media were significantly higher than those of all other medias during the first season followed by M<sub>1</sub> media (20mm gravel). Among all the treatments plants grown in 8-10mm gravel exhibited highest plant height (140.556 cm, 140.778 cm), plant spread (66.833cm, 66.556 cm), while shortest days to flowering (53.889 days, 53.889 days) during the first and second season. The plant parameters in M<sub>2</sub> and M<sub>3</sub> were on par with each other. The variation of plant characteristics among different media in the aquaponics system reveal that 8-10mm gravel is more favorable for the growth and yield of tomato plants than other media. This is due to the size and shape of the particles in the 8-10mm gravel. Moreover, this media does not clog frequently, and algae formation was negligible compared to all other media. The aeration and root-holding strength are also higher in the case of 8-10mm gravel. The results are in line with the findings of Suseela [10].

Table 1. Growth parameters of Manulakshmi in aquaponics system during first season

Media	Plant height	Plant spread	Days to first flowering
M <sub>1</sub>	135.111± 3.756 <sup>b</sup>	63.167 ± 1.732 <sup>b</sup>	57.667± 2.179 <sup>b</sup>
M <sub>2</sub>	129.333± 3.606 <sup>c</sup>	60.333± 2.062 <sup>c</sup>	61.778± 1.093 <sup>c</sup>
M <sub>3</sub>	125.889± 3.333 <sup>c</sup>	58.500± 2.092 <sup>c</sup>	60.000± 2.000 <sup>c</sup>
M <sub>4</sub>	140.556± 3.575 <sup>a</sup>	66.833± 2.739 <sup>a</sup>	53.889± 1.167 <sup>a</sup>
F- Value	29.470	24.832	36.980
P- Value	< .001	< .001	< .001
CD (0.05)	3.434	2.103	1.617

Table 2. Growth parameters of Manulakshmi in aquaponics system during second season

Media	Plant height	Plant spread	Days to first flowering
M <sub>1</sub>	135.333± 1.871 <sup>b</sup>	62.778± 1.856 <sup>b</sup>	56.444± 1.236 <sup>b</sup>
M <sub>2</sub>	129.889± 1.833 <sup>c</sup>	59.667± 2.000 <sup>c</sup>	62.222± 0.667 <sup>c</sup>
M <sub>3</sub>	126.111± 1.900 <sup>c</sup>	58.111± 1.900 <sup>c</sup>	59.889± 1.167 <sup>d</sup>
M <sub>4</sub>	140.778± 2.438 <sup>a</sup>	66.556± 1.590 <sup>a</sup>	53.889± 0.782 <sup>a</sup>
F- Value	89.961	36.864	123.718
P- Value	< .001	< .001	< .001
CD (0.05)	1.948	1.772	0.955

The analysis of growth parameters of Yakamoz RZ in aquaponics system during first and second season are summarized in Table 3 and 4 respectively. It reveals that plants grown in M<sub>4</sub> media were significantly superior than all other medias during both seasons. The plant spread of Yakamoz RZ in M<sub>1</sub> media was significantly higher than that of all other medias.

Table 3. Growth parameters of Yakamoz RZ in aquaponics system during first season

Media	Plant height	Plant spread	Days to first flowering
M <sub>1</sub>	139.222± 3.528 <sup>b</sup>	50.500± 1.436 <sup>a</sup>	51.444± 3.087 <sup>a</sup>
M <sub>2</sub>	133.667± 2.646 <sup>c</sup>	48.111± 1.965 <sup>b</sup>	57.778± 1.787 <sup>b</sup>
M <sub>3</sub>	130.889± 2.261 <sup>c</sup>	46.222± 1.481 <sup>b</sup>	55.000± 3.000 <sup>b</sup>
M <sub>4</sub>	145.222± 4.410 <sup>a</sup>	46.111± 2.421 <sup>b</sup>	50.556± 2.603 <sup>a</sup>
F- Value	32.932	10.912	14.018
P- Value	< .001	< .001	< .001
CD (0.05)	3.189	1.798	2.567

Table 4. Growth parameters of Yakamoz RZ in aquaponicssystem during second season

Media	Plant height	Plant spread	Days to first flowering
M <sub>1</sub>	139.778± 3.114 <sup>b</sup>	54.667± 2.500 <sup>a</sup>	53.778± 2.048 <sup>ab</sup>
M <sub>2</sub>	132.222± 1.856 <sup>c</sup>	49.111± 0.928 <sup>b</sup>	57.556± 1.509 <sup>c</sup>
M <sub>3</sub>	131.556± 2.963 <sup>c</sup>	48.778± 1.481 <sup>b</sup>	56.111± 1.900 <sup>bc</sup>
M <sub>4</sub>	147.556± 2.506 <sup>a</sup>	47.556± 2.506 <sup>b</sup>	51.222± 3.232 <sup>a</sup>
F- Value	72.008	23.128	13.496
P- Value	< .001	< .001	< .001
CD (0.05)	2.553	1.898	2.179

The Statistical analysis of yield parameters of Manulakshmi in aquaponics system during first and second season are shown in Table 5 to Table 6 respectively. The analysis of yield in aquaponics system showed that the plants grown in M<sub>4</sub> media (1.730Kg and 1.729Kg) were significantly higher than that of all other medias during first and second season respectively. The results were in close agreement with the findings of Salam et al. [11] and Gaviria et al. [12]. There was no significant difference between the medias in case of average weight of fruit. The days to first harvest of Manulakshmi in M<sub>4</sub> were significantly lower than all other medias in both seasons.

Table 5. Yield parameters of Manulakshmi in aquaponicssystem during first season

Media	Days to first harvest	Total yield	Average weight of fruit
M <sub>1</sub>	86.444± 1.333 <sup>b</sup>	1.531± 0.061 <sup>b</sup>	38.067± 0.472 <sup>a</sup>
M <sub>2</sub>	88.889± 1.054 <sup>c</sup>	1.297± 0.087 <sup>c</sup>	38.022± 0.412 <sup>a</sup>
M <sub>3</sub>	87.778± 1.563 <sup>bc</sup>	1.330± 0.067 <sup>c</sup>	38.122± 0.393 <sup>a</sup>
M <sub>4</sub>	81.778± 1.563 <sup>a</sup>	1.730± 0.064 <sup>a</sup>	38.433± 0.346 <sup>a</sup>
F- Value	45.257	72.902	1.870
P- Value	< .001	< .001	0.155
CD (0.05)	1.341	0.068	0.393

Table 6. Yield parameters of Manulakshmi in aquaponicssystem during second season

Media	Days to first harvest	Total yield	Average weight of fruit
M <sub>1</sub>	86.111± 1.167 <sup>b</sup>	1.541± 0.063 <sup>b</sup>	38.322± 0.427 <sup>a</sup>
M <sub>2</sub>	89.444± 1.236 <sup>c</sup>	1.302± 0.078 <sup>c</sup>	38.200± 0.614 <sup>a</sup>
M <sub>3</sub>	88.000± 1.225 <sup>c</sup>	1.332± 0.059 <sup>c</sup>	38.200± 0.324 <sup>a</sup>

M <sub>4</sub>	81.667± 1.323 <sup>a</sup>	1.729± 0.050 <sup>a</sup>	38.611± 0.451 <sup>a</sup>
F- Value	67.011	89.515	1.560
P- Value	< .001	< .001	0.218
CD (0.05)	1.191	0.061	0.448

The Statistical analysis of yield parameters of Yakamoz RZ in aquaponics system during first and second season are shown in Table 7 to Table 8 respectively. Plants grown in M<sub>4</sub> media had the highest yield than that of all other medias in both seasons. There was no significant difference between the average fruit weight in four media in both seasons. Yakamoz RZ grown in 20mm gravel and 8-10 mm gravel flowered earlier than that in 8-20mm broken tiles, 8-15mm hydroton.

Table 7. Yield parameters of Yakamoz RZ during first season

Media	Days to first harvest	Total yield	Average weight of fruit
M <sub>1</sub>	78.889± 2.667 <sup>a</sup>	3.577± 0.062 <sup>b</sup>	88.689± 0.909 <sup>a</sup>
M <sub>2</sub>	85.778± 2.333 <sup>b</sup>	3.203± 0.162 <sup>d</sup>	87.911± 1.104 <sup>a</sup>
M <sub>3</sub>	83.778± 2.108 <sup>b</sup>	3.357± 0.092 <sup>c</sup>	87.833± 0.971 <sup>a</sup>
M <sub>4</sub>	78.000± 1.732 <sup>a</sup>	3.871± 0.136 <sup>a</sup>	88.211± 0.448 <sup>a</sup>
F- Value	25.504	52.887	1.697
P- Value	< .001	< .001	0.187
CD (0.05)	2.150	0.114	0.859

Table 8. Yield parameters of Yakamoz RZ during second season

Media	Days to first harvest	Total yield	Average weight of fruit
M <sub>1</sub>	77.778± 1.563 <sup>a</sup>	3.580± 0.063 <sup>b</sup>	88.278± 0.689 <sup>a</sup>
M <sub>2</sub>	84.000± 2.449 <sup>b</sup>	3.246± 0.125 <sup>d</sup>	87.978± 0.418 <sup>a</sup>
M <sub>3</sub>	83.778± 2.539 <sup>b</sup>	3.404± 0.061 <sup>c</sup>	87.533± 0.758 <sup>a</sup>
M <sub>4</sub>	76.667± 1.323 <sup>a</sup>	3.898± 0.125 <sup>a</sup>	88.133± 0.604 <sup>a</sup>
F- Value	32.516	72.175	2.354
P- Value	< .001	< .001	0.091
CD (0.05)	1.961	0.096	0.606

#### 4. CONCLUSION

The variation of plant characteristics among different media in the aquaponics system reveal that 8-10mm gravel is more favorable for the growth and yield of tomato plants than other media. This is due to the size and shape of the particles in the 8-10mm gravel. Moreover, this media does not clog frequently, and algae formation was negligible compared to all other media. The aeration and root-holding strength are also higher in the case of 8-10mm gravel.

#### REFERENCES

1. ChandR, Parappurathu S. Temporal and spatial variations in agricultural growth and its determinants. *Economic and Political Weekly*. 2012;.47(27): 55-64.
2. Graber A, Junge R. Aquaponic Systems: Nutrient recycling from fish wastewater by vegetable production. *Desalination*. 2009;246(1-3):147-156.
3. Roe BR, Midmore DJ. *Sustainable Aquaponics*. Center for Plant and Water Science, CQ University, Rockhampton, Queensland, Australia; 2008.
4. Martins CIM, Eding EH, Verdegem MC, Heinsbroek LT, Schneider O, Blancheton JP, d'Orbcastel ER Verreth, JAJ. New developments in recirculating aquaculture systems in Europe: A perspective on environmental sustainability. *Aquac Eng*2010;43(3): 83-93.
5. SaceCF, FitzsimmonsKM. Vegetable production in a recirculating aquaponic system using Nile tilapia (*Oreochromis niloticus*) with and without freshwater prawn (*Macrobrachiumrosenbergii*). *Academia J Agric Res*.2013;1(12):236-250.
6. ReshHM. *Hydroponic Food Production*. 6th ed. New Concept Press Publishing Co., Mahwah, NJ; 2004.
7. Tyson RV, Hochmuth R C, Lamb E M, Hochmuth GJ, Sweat MS. A decade of change in Florida's greenhouse vegetable industry. *Proc Fla St Hortic Soc*. 2001; 114:280–283.
8. Lang HJ, Elliott GS. Enumeration and inoculation of nitrifying bacteria in soilless potting media. *J Am Soc Hort Sci*.1997;122(5): 709–714.
9. Suseela P, Sam B. Selection of optimal design of polyhouse and suitable varieties of salad cucumber under humid tropical climate. *Ann Hortic*. 2018; 11(2): 126-130.
10. Suseela P. Study on structural design and management for high- tech horticulture in Kerala. *Instructional Farm, Vellanikkara*; 2019.
11. SalamM A, Jahan N, Hashem S,Rana KMS. Feasibility of tomato production in aquaponic system using different substrates. *Prog Agric*2014; 25: 54-62.
12. Gaviria L, Marín D, Martínez J and Muñoz A. The effect of media type on the growth and yield of tomatoes in aquaponic systems. *Aquaponics J*.2021;15(2): 123-134.