

Field efficacy of certain chemicals and neem products against shoot and fruit borer, *Earias vittella* (Fabricius.) on okra in Prayagraj, U.P.

Abstract

A field trial was conducted at the research plot of the Department of Entomology at Central Research Field, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during *kharif* season of 2023 where seven treatments were replicated thrice in Randomized Block Design against *Earias vittella* viz. (T₁) Fipronil 5%SC @ 1 ml/lit, (T₂) Spinosad 45%SC @ 1 ml/lit, (T₃) Azadiractin 1%EC @ 3 ml/lit, (T₄) Chlorantraniliprole 18.5%SC @ 0.3 ml/lit, (T₅) Imidacloprid 17.8%SL @ 0.4 ml/lit, (T₆) Emamectin benzoate 5%SG @ 0.3 ml/lit, (T₇) NSKE 5% @ 50 ml/lit and (T₀) Control and their influences on yield of okra. Each treatment was sprayed twice at 14 days interval. The observations of shoot and fruit infestation per plant were taken one day before, 3rd, 7th and 14th day after each spray. All the insecticides tested, significantly reduced the pest infestation compared to untreated plot. The lowest percent infestation of shoot and fruit borer was observed in Chlorantraniliprole 18.5%SC (8.96%, 9.29%) and Spinosad 45%SC (9.74%, 10.42%) followed by Imidacloprid 17.8%SL (10.58%, 11.04%) Emamectin benzoate 5%SG (11.89%, 12.27%), Fipronil 5%SC (12.46%, 13.09%), Azadiractin 1%EC (12.77%, 14.16%), NSKE 5% (13.42%, 14.63%). The mean crop yield ranged between 30q/ha to 160q/ha. The highest was recorded in Chlorantraniliprole 18.5%SC (160 q/ha) followed by Spinosad 45%SC (154 q/ha). When cost benefit ratio worked out, the best and the most economic treatment was Chlorantraniliprole 18.5%SC (1:5.7) followed by Spinosad 45%SC (1:5.2), Imidacloprid 17.8%SL (1:5.1), Emamectin benzoate 5%SG (1:4.3), Fipronil 5%SC (1:3.9), Azadiractin 1%EC (1:3.0), NSKE 5% (1:2.5) and Untreated (1:1.18).

Keywords – Chlorantraniliprole, *Earias vittella*, efficacy, insecticides okra, yield

Introduction –

Okra [*Abelmoschus esculentus* (L.)] is an annual vegetable belonging to *Malvaceae* family; it is also known by different names viz., ladies finger, bhindi, bamia, okra or gumbo in

different parts of the world. Okra is known as ‘Queen of vegetables’. Okra is valued for its tender green fruits. It is cooked in variety of ways and used as an ingredient in a wide variety of dishes. Its medicinal value has also been reported in curing ulcer and relief from haemorrhoids (**Kaveri and Kumar, 2020**). The tender fruits of okra are used as vegetables or in culinary preparations as sliced and dried pieces. It is commercially cultivated in India, Turkey, Iran, Western Africa, Yugoslavia, Bangladesh, Afghanistan, Burma, Japan, Malaysia, Brazil, Ghana, Ethiopia and Southern United States. Though okra finds its origin in South-Africa, India stands top in area and production. The major okra growing states includes Andhra Pradesh, Assam, Uttar Pradesh, Bihar, Orissa, West Bengal, Maharashtra and Karnataka (**Mulani et al., 2021**).

India ranks first in world with the production of 6371 million tons followed by Nigeria with 1837 million tons and then comes Mali with 659 million tons. Gujarat is the leading okra producing state with production of 1019.42 thousand tons followed by West Bengal with 893.96 thousand tons then comes Bihar with 794.10 thousand tonnes (**Food and Agriculture Organization, 2021**).

Okra has good nutritional value particularly the high content of vitamin C (13mg / 100g), fat (0.2g / 100g), carbohydrate (6.4g / 100g), iron (1.5mg / 100g), moisture (89.6g / 100g), protein (1.9g / 100g), fiber (1.2g / 100g), calories (35kcal / 100g), and other minerals (**Chandravanshi et al., 2018**).

Okra crop suffers damage by a number of insect pest viz., the jassids, *Amrasca biguttula biguttula*, Ishida; the aphid, *Aphis gossypii* Glover; the fruit borers, *Earias insulana* Boisduval and *Earias vittella* (Fab.); *Helicoverpa armigera* Hub.; whitefly, *Bemisia tabaci* Genn.; and red spider mite, *Tetranychus cinnabarinus* that appears occasionally. Among all pests, shoot and fruit borer, *Earias vittella* (Fabricius) is the most damaging pest of okra as young larva borers into tender shoots in early vegetative growth of plants (**Dash et al., 2020**).

Okra shoot and fruit borer is the most destructive pest of okra as the larva bores in to young shoots and fruits. The damage due to fruit borer accounts for nearly 22.5% in Uttar Pradesh

25.93% to 40.91% in Madhya Pradesh. The major notorious pest of okra fruit and shoot borer having endemic nature, inflect direct losses in yield of marketable fruits. According to an estimate this pest can cause a 76% loss in fruit yield of okra (**Rajput and Tayde, 2017**).

Justification –

Conventionally farmers are using various types of synthetic chemical insecticides to control shoot and fruit borer. But the unconscious and unjustified use of synthetic pesticides creates several problems in agro-ecosystem such as direct toxicity to beneficial insects, fishes and man. Among the various methods of pest management, the use of insecticides forms the first line of defence against the insect pests. Newer insecticide molecules are better alternative to conventional synthetic insecticides in the context of environmentally benign management tactics so also in order to mitigate the adverse effect on the total environment. In many cases, alternate or eco-friendly method of insect management offer adequate level of pest control with less hazards and safe to non-target organisms. With this background, the present study was undertaken to evaluate the efficacy of insecticides against *Earias vittella* on okra.

Material Methods –

The experiment was conducted at the experimental research plot of the Department of Entomology, Central Research Farm, Sam Higginbottom University of Agriculture Technology and Sciences, during the *kharif* season of 2023. The experimental design was laid in Randomized Block Design using Arka Anamika variety with 8 treatments, each replicated thrice. The plot size was 2m × 1m with a spacing of 45cm × 30cm. The treatments included – Fipronil 5% SC 1ml / lit, Spinosad 45% SC 1ml / lit, Azadirachtin 1% EC 3ml / lit, Chlorantraniliprole 18.5% SC 0.3ml / lit, Imidacloprid 17.8% SL 0.4ml / lit, Emamectin benzoate 5% SG 0.3ml / lit, NSKE 5% 50ml / lit and a control. Application of the two rounds of insecticidal treatments were applied at 15 days interval.

The numbers of infestation were counted on 5 randomly selected plants in each plot. The pre-treatment count was made a day before the spray whereas, the post-treatment counts were made on 3rd, 7th and 14th day after each spray. The percent Infestation over control against Shoot and fruit borer was calculated by considering the mean of three observations recorded at 3rd, 7th, and

14th day afterspray.

Result and Discussion –

Among all the treatments, Chlorantraniliprole 18.5%SC was found to be most effective in managing the shoot and fruit borer infestation on okra .The values obtained in the first and second spray were 8.96% and 9.29%. These results are supported by **Chandran *et al.*,(2020)** and **Reddy *et al.*, (2019)**. Spinosad 45% SC was also found to be very effective in reducing the infestation of shoot and fruit borer. The same results were observed by **Kaveri and kumar (2022)** and **Rajput and Tayde (2017)**, who reported that application of Spinosad 45%SC reduced the borer infestation and recorded the lowest percentage of shoot and fruit damage. Where the observations of first and second sprays obtained were 9.74% and 10.42% infestation over untreated plot. The next best treatment was found to be Imidacloprid 17.8%SL in which the efficacy values of first and second spray were 10.58% and 11.04% respectively, This results were supported by **Janu and Kumar (2022)** and **Pachole *et al.*, (2017)**. The efficacy of Emamectin benzoate 5% SG on shoot and fruit borer in first and second spray are 11.89% and 12.27% respectively. These results are as per the findings of **Manikanta and kumar (2022)** and **Dash *et al.*, (2020)**.The next best treatment was found to be Fipronil 5%SC in which the efficacy values of first and second spray were 12.46% and 13.09% respectively to the similar findings of **Rani and Kumar (2022)** and **Naidu and Kumar (2019)**. The least effective treatments were found to be Azadiractin 1%EC and NSKE 5% in which efficacy values of first and second sprays were (12.77%), (14.16%) and (13.42%), (14.63%) respectively. The results obtained were as similar to **Choudhury *et al.*, (2021)** and **Patil *et al.*, (2021)**.

Table 1. Field efficacy of certain chemicals and neem products against shoot and fruit borer, *Earias vittella* (Fabricius.) on okra in Prayagraj, U.P.

Percent shoot infestation	Percent Fruit infestation
---------------------------	---------------------------

Tr. No.	Treatments	Dose (ml/lit)	1 DBS	3 DAS	7 DAS	14 DAS	Mean	3 DAS	7 DAS	14 DAS	Mean	Yield (q/ha)	C:B ratio
T ₀	Control	-	14.02	15.02	15.17	15.84	15.34	15.99	16.36	17.05	16.46	30	1:1.18
T ₁	@Fipronil 5% SC	1	14.70	13.21	10.17	14.01	12.46	14.08	12.04	13.17	13.09	103	1:3.9
T ₂	@Spinosad 45% SC	1	15.80	10.07	7.73	11.43	9.74	11.54	9.03	10.69	10.42	154	1:5.2
T ₃	@Azadirachtin 1% EC	3	14.96	13.07	10.42	14.83	12.77	14.98	13.20	14.32	14.16	81	1:3.0
T ₄	@Chlorantraniliprole 18.5%SC	0.3	16.69	9.08	7.59	10.22	8.96	10.38	8.09	9.42	9.29	160	1:5.7
T ₅	@Imidacloprid 17.80 % SL	0.4	17.59	11.00	8.03	12.72	10.58	12.02	10.08	11.03	11.04	135	1:5.1
T ₆	@Emamectin benzoate 5%SG	0.3	17.50	12.72	9.14	13.81	11.89	13.27	11.06	12.49	12.27	115	1:4.3
T ₇	@NSKE 5%	50	16.75	13.49	11.43	15.35	13.42	15.42	13.80	14.67	14.63	69	1:2.5
F – test			NS	S	S	S	S	S	S	S	S	-	-
C.D.(0.05%)			-	1.19	1.65	1.32	1.31	1.68	3.07	2.20	0.81	-	-
S.Ed (±)			1.65	0.58	0.80	0.64	0.63	0.81	1.48	1.06	0.39	-	-

***Figures in parentheses are arc sin transformation values, DAS – day after spray, DBS – Day before spray, NS – non significant, S – significant, C:B – cost benefit ratio.**

Conclusion –

From the present study, the results showed that Chlorantraniliprole 18.5%SC and Spinosad 45%SC are most effective treatments against okra shoot and fruit borer (*Earias vittella*) which

produced maximum yield and recorded highest Cost-Benefit ratio compared to other treatments. Neem products such as Azadiractin 1%EC and NSKE 5% found to be least effective in managing *Earias vittella*. Hence, it is suggested that the effective insecticides may be alternated in harmony with the existing Integrated pest management programs in order to avoid the problems associated with insecticidal resistance, pest resurgence etc. Botanicals are the part of integrated pest management in order to avoid indiscriminate use of pesticides causing pollution in the environment and not much harmful to beneficial insects.

Reference –

- Kaveri, G. and Kumar, A.(2020).** Field efficacy of certain biopesticides against okra shoot and fruit borer, *Earias vittella* (Fabricius) on okra, *Abelmoschus esculentus* (Linn.) Moench. *Journal of Entomology and Zoology Studies*, **8**(6):1279-1281.
- Mulani, H. B., Bantewad, S. D. and Jayewar, N. E.(2021).** Biointensive management of Okra shoot and fruit borer *Earias vittella*. *The Pharma Innovation Journal*, **10**(8):681-690.
- Anonymous (2021).** Food Agriculture Organization (FAO STAT).
- Chandravanshi, D. K., Tomar, R. K. S., Awasthi, A. K. and Kerketta, A. (2019).** Field efficacy of different insecticides and bio-pesticides against okra shoot and fruit borer. *Journal of Pharmacognosy and Phytochemistry*, **8**(1):2623-2625.
- Dash, L., Ramalakshmi, V. and Padhy, D.(2020).** Bio-efficacy of Emamectin benzoate 5% SG against shoot and fruit borer *Earias vittella* (Fabricius) on okra. *The Pharma Innovation Journal*, **9**(12):144-146.
- Kumar, K. I. and Tayde, A. R (2018).** Screening of okra genotypes against shoot and fruit borer (*Earias vittella* Fab.) under field conditions in Allahabad. *Journal of Pharmacognosy and Phytochemistry*, **7**(1):657-659.
- Rajput, G. S. and Tayde, A.(2017).** Population dynamics and comparative efficacy of certain novel insecticides, botanicals and bioagents, against shoot and fruit borer (*Earias vitella* Fabricius) of Okra crop. *Journal of Entomology and Zoology Studies*, **5**(4):1667-1670.

- Chandran, R., Ramesha, B. and Sreekumar, K. M. (2020).** Efficacy of new insecticides against okra shoot and fruit borer, *Earias vittella* (Fab.) (Lepidoptera: Noctuidae). *Entomon*, **45**(4): 295-300.
- Reddy, G. N., Thara, K. T. and Deepak, S.(2019).** Field efficacy of selected bio-agent and insecticide against shoot and fruit borer, *Earias vittella* (Noctuidae: Lepidoptera) on okra. *Journal of Entomology and Zoological studies*, **7**(3):380-383.
- Manikanta, S. E. N. and Kumar, A (2022).** Efficacy of certain chemicals and essential oils against okra shoot and fruit borer [*Earias vittella* (Fabricius)]. *The Pharma Innovation Journal*, **11**(4):1385-1389.
- Rani, K. and Kumar, A.(2022).** Field efficacy of different chemicals against shoot and fruit borer [*Earias vittella* (Fabricius)] of okra [*Abelmoschus esculentus* (L.) Moench]. *The Pharma Innovation Journal* , **11**(4):1603-1607.
- Naidu, G. and Kumar, A.(2019).** Field efficacy of certain insecticides against shoot and fruit borer (*Earias vittella* Fab.) on rainy season okra in Prayagraj (UP). *Journal of Entomology and Zoology Studies*, **7**(6):1211-1213.
- Janu, R. and Kumar, A (2022).** Field efficacy of selected insecticides against okra shoot and fruit borer [*Earias vittella* (Fabricius)]. *The Pharma Innovation Journal*, **11**(4):1549-1551.
- Pachole, S. H., Thakur, S. and Simon, S.(2017).** Comparative bioefficacy of selected chemical insecticides and biorationals against shoot and fruit borer, (*Earias vittella* Fabricius) on okra [*Abelmoschus esculentus* (L.) Moench]. *Journal of Pharmacognosy and Phytochemistry*, **6**(5):1493-1495.
- Choudhury, M. A. R., Mondal, M. F., Khan, A. U., Hossain, M. S., Azad, M. O. K., Prodhan, M. D. H. and Naznin, M. T. (2021).** Evaluation of biological approaches for controlling shoot and fruit borer (*Earias vittella* F.) of okra grown in peri-urban area in Bangladesh. *Horticulturae*, **7**(1):7.

Patil, H. N., Tayde, A. R. and Chandar, A. S.(2022). Comparative efficacy of botanicals against shoot and fruit borers,(*Earias vittella*, Fabricius) on okra. *The Pharma Innovation Journal*, **11**(2):222-224.

UNDER PEER REVIEW